



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

> OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

Tack

SEP 2 3 2009

Petro-Canada c/o Robert G. Butz Ph.D. Carter Ledyard & Milburn LLP 701 8th Street, N.W., Suite 410 Washington, DC 20001-3893

Dear Dr. Butz:

Subject: Revised Labeling PC RTU Herbicide EPA Registration No. 69526-12 Your Submission Dated August 31, 2009

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

a. When placing an optional statement with a footnote on the final printed labeling assure the associated footnote is also on the labeling.

b. Delete the picture of the house being held in hands. This is an implied safety claim that the home is in safe hands and could conflict with potential copyrights.

c. Delete "General" in the Precautions and Restrictions heading. This implies that these restrictions do not all have to be followed as specified on the label

2. Submit one (1) copy of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

If you have any questions concerning this letter, please contact me at 703-305-6224.

Sincerely yours,

Joanne I. Miller Product Manager (23) Herbicide Branch Registration Division (7505P) Ċ

Enclosure

[BATCH CODE WILL APPEAR ON IMMEDIATE CONTAINER]

[FRONT PANEL]

PC RTU Herbicide

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt *		0.0340%
Mecoprop-p, dimethylamine salt	**	0.0220%
Dicamba, dimethylamine salt ***		0.0042%
OTHER INGREDIENTS:		<u>99.9398%</u>
	Total	100.0000%

* 2,4-D Acid Equivalent	0.028%
** Mecoprop-p Acid Equivalent	
*** Dicamba Acid Equivalent	

ACCEPTED with COMMENTS In EPA Letter Dated:

SEP 2 3 2009

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.



KEEP OUT OF REACH OF CHILDREN

See (back panel) (side panel) (inside) (booklet) for precautionary statements and directions for use.

FORMULATED IN CANADA FOR:

PETRO-CANADA 385 SOUTHDOWN ROAD MISSISSAUGA, ONTARIO L5K 1A8

EPA REG. NO. 69526-12

EPA EST. NO. 79693-CAN-001

NET CONTENTS: 6 fl. oz. (177mL); 24 fl. oz. (1pt, 8 fl.oz) / 710ml; 64 fl. oz. (1/2 galion) (1.89 L); 1 gal. (3.78 L); 1.33 gal. (5.03 L); 1.58 gal. (5.98 L); 1.75 gal. (6.62 L)

US Patent pending

Protect from Freezing

[One of the following phrases will be found on FPL] For use only on home lawns [OR] For use on home lawns and other grassy areas around the home.

 \circ

[Following notations may be found on back panel in reference to claims on front or back panels]

* When compared to other pesticides using these active ingredients.

[†]Important: Do not apply to Bentgrass and Floratam. [Spanish Translation:] / No se aplique al Bentgrass y a Floratam. See full list of controlled weeds and use directions on peel-away label.

[The Following are optional marketing statements and illustrations that may appear on front panel, front of booklet attached to back of container, back panel, or side panel:]

Microtechnology improves penetration of reduced active ingredient formulation *

Won't harm lawns when used as directed. [†]

Lawn Weed Remover[†] [Spanish Translation:] Eliminador de malas hierbas que no daña el césped[†]

Use as a spot treatment to control broadleaf weeds on a variety of lawns[†], in yards, vacant lots and other grassy areas around the home.

When used as directed, kills broadleaf lawn weeds without harming most lawn grasses^T.

KILLS BROADLEAF WEEDS INCLUDING: Dandelions, Plantains, Clover, Chickweed, Bedstraw, Black Medic, English Daisy, Curled Dock, Heal-all, Ground Ivy (Creeping Charlie), Knotweed, Ragweed, Shepherd's Purse, plus (over 60) (many) other lawn weeds listed (on) (side) (back) (panel) (inside) (inside booklet)

KILLS: Dandelions, Plantains, Clover, Chickweed, Bedstraw, Black Medic, English Daisy, Curled Dock, Heal-all, Ground Ivy (Creeping Charlie), Knotweed, Ragweed, Shepherd's Purse, plus over 60 other lawn weeds.

When used as directed, controls a wide range of broadleaf weeds in most home lawns^T.

Kills Weeds Dead

Kills listed broadleaf weeds

Kills over 60 broadleaf weeds without harming lawns[†]. See (inside booklet) (booklet) (back panel) (back) (side panel) (side) for details.

Kills down to the root

Kills listed broadleaf weeds in lawns - right down to the root.

Outdoor use only

Ready-To-Use. No mixing.

Ready to use! No mixing required! T

Reduced Active Ingredient Formulation^{*}

Low-Odor formulation

Low-Odor formula

Convenient Ready-to-Use Formula

Step 1: Shake Well S

Step 2: Apply Step 3: Let Dry

Container spot treats up to 50 weeds [for 6 fl. oz.]

2

ESL20080422 Notif EPA 3/6/2009

MASTER LABEL

Amend 8/31/2009

G

[Bracketed information will be removed from final printed label. Phrases in parenthesis are optional.]

Container spot treats more than 200 weeds [for 24 fl. oz.] Container spot treats more than 500 weeds [for 1/2 gal.] Container spot treats more than 1000 weeds [for 1 gal.] Container spot treats more than 1333 weeds [for 1.33 gal.] Container spot treats more than 1580 weeds [for 1.58 gal.] www.todaysclearchoice.com

PEEL HERE>

[The following are illustration that may appear on final printed label:]

[House and home lawn in hands]



[Dandelion with root highlighted]



[Dandelion with root highlighted showing spray and portion of spray bottle]



.

- .

[To Appear on Back or Side Panel or Inside Booklet or Base Panel]

PRECAUTIONARY STATEMENTS.

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all label directions.

GENERAL PRECAUTIONS AND RESTRICTIONS

- Do not apply this product in a way that will contact any person or pet, either directly or through drift.
 Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply near lakes, streams, rivers or ponds.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not apply in windy conditions.
- Do not apply as a fine mist because of potential injury to desirable plants.

WHEN TO USE

- Apply when weeds are young and actively growing.
- Do not water within 24 hours after spraying.
- Spray when daytime temperatures are above 45° F and below 90° F.
- Do not apply when the lawn is under severe stress from drought, insect attack or high temperatures.
- Do not use more product than specified on the label.
- Newly seeded lawns may be treated after the third mowing.
- If the lawn is dry, water 24 hours before using this product.

WHERE TO USE

4

This product is used as a spot treatment to control broad leaf weeds on a variety of lawns*, in yards, vacant lots and other grassy areas.

*IMPORTANT: Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain hybrid bermudagrasses vary in tolerance to this product. Temporary discoloration may occur on St. Augustine and Centipede.

LIST OF WEEDS: This product controls the following weeds:

Bedstraw
Bindweed (field)
Birdsfoot trefoil
Black medic
Blackseed plantain
Blueweed
Brambles
Broad-leaved plantain
Buckhorn plantain
Bull thistle
Burdock
Buttercup
Canada thistle
Carolina geranium
Chickweed
Chicory
Clover
Cocklebur
Common chickweed
Cudweed
Daisy fleabane
Dandelion
Devil's paint brush
(hawkweed)
Dock (curled)
English daisy
Evening primrose
Fall dandelion (fall hawkbit)
False dandelion
Fleabane
Florida pusley
Frenchweed poison oak

Grass leaved stitchwort Ground ivy (creeping Charlie) Hawkweed Heal-all Henbit Honeysuckle Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Purslane Ragweed	
Charlie) Hawkweed Heal-all Henbit Honeysuckle Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Grass leaved stitchwort
Charlie) Hawkweed Heal-all Henbit Honeysuckle Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Ground ivy (creeping
Heal-all Henbit Honeysuckle Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Henbit Honeysuckle Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Hawkweed
Honeysuckle Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Heal-all
Jimsonweed Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Henbit
Knotweed Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Honeysuckle
Kochia Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Jimsonweed
Kudzu Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Knotweed
Lambsquarters Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Kochia
Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Kudzu
Little starwort Mallow Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	Lambsquarters
Morningglory Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Mouse-ear chickweed Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Mustards Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Narrow-leaved plantain Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Oxalis (yellow woodsorrel) Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Ox-eye daisy Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Pennywort Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Peppergrass Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Pigweed Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Pineapple weed Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Plantains (narrow, buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
buckhorn, broadleaf) Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Poison ivy Poorjoe Povertyweed Prostrate pigweed Purslane	
Poorjoe Povertyweed Prostrate pigweed Purslane	
Povertyweed Prostrate pigweed Purslane	
Prostrate pigweed Purslane	
Purslane	
Ragweed	
	Ragweed

Russian thistle
Sandwort (thyme-leaved)
Sheep sorrel
Shepherd's purse
Smartweed (green)
Soliva
Sow thistle
Speedwell (purslane
leaved)
Spotted spurge
Spreadwell (annual)
Spurge
Stitchwort
Stonecrop (mossy)
Sumac
Veronica (thyme-leaved)
Vervain
Vetch
Violet
Wild aster
Wild blackberry
Wild carrot
Wild garlic
Wild geranium
Wild lettuce
Wild onion
Wild radish
Wild raspberry
Yarrow
Yellow hawkweed

HOW TO USE

- This product is ready-to-use. Do not mix or dilute.
- Shake well before use.
- Open spray nozzle.
- Adjust nozzle to provide desired spray pattern.
- Hold spray about 12 inches from weed.
- Aim at center of weed and spray to lightly cover. Do not overspray.
- Rotate nozzle to closed position.
- Weeds begin to die after 12 hours. It may take up to 4 weeks for a weed to completely die.
- If necessary, repeat application after 8 weeks.

5

STORAGE AND DISPOSAL

Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place, in a locked storage area.

Pesticide and Container Disposal:

If empty: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling unavailable, place container in trash.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

It is impossible to eliminate all risks inherently associated with the use of this product. Injury to desirable plants, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Petro-Canada, the manufacturer, or seller. All such risks shall be assumed by the buyer. To the extent consistent with applicable law, neither Petro-Canada, the manufacturer nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product. Except as provided by applicable law or except as warranted by this label, Petro-Canada, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Except as provided by applicable law, buyer's or user's exclusive remedy, and Petro-Canada's, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.