

69526-15

5/26/2009

10f 17



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

69526-15

26 MAY 2009

Term of Issuance:

Conditional

Name of Pesticide Product:

PC Herbicide Concentrate Plus

NOTICE OF PESTICIDE:

X Registration
Reregistration
(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Petro Canada
2489 Sheridan Way
Mississauga, Ontario LSK 1A8
Canada

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1) Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of this product under FIFRA section 4.
2) Add the phrase "EPA Reg. No. 69526-15" to the labeling.
3) Assure that the EPA Establishment Number and Net Contents are on the label.
4) You must generate one-year studies for guidelines 830.6317 (storage stability) and 830.6320 (corrosion characteristics) with observations being made at 0, 3, 6, 9, and 12 month intervals. The results being submitted to the Agency within one year of the date on this notice.
5) Per the Pesticide Registration (PR) Notice 93-7 Supplement Three-A and the combined RED label table, the Restricted-Entry Interval in the Agricultural Use Requirements box on page 3 must be changed from "24 hours" to "48 hours".

SEE COMMENTS ON NEXT PAGE

Signature of Approving Official:

Joanne I. Miller
Product Manager 23
Herbicide Branch
Registration Division (7505P)

Handwritten signature of Joanne I. Miller

Date:

26 MAY 2009

6) Per the Acute Toxicity Review, you have the option to add the signal word CAUTION to the label.

7) Per the Acute Toxicity Review, "Contains Petroleum Distillate." must be added to the label under Hazards to Humans and Domestic Animals.

8) Per the Acute Toxicity Review, First Aid statements are not required, though Category III statements may be used. If you chose to include a First Aid box on the label, the review recommends that a specific "if swallowed" statement be added based on the petroleum distillate concentration.

If Swallowed:

- Immediately call a poison control center or doctor.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give any liquid to the person.
- Do not give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: May pose an aspiration pneumonia hazard. Contains petroleum distillate.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-XXX-XXXX for emergency medical treatment information.

9) Per the Acute Toxicity Review and the combined RED label table requirements for liquid pesticides, the Personal Protective Equipment (PPE) section on page 2, should read as follows:

"Some materials that are chemical-resistant to this product are Barrier Laminate, Nitrile rubber, Butyl, Neoprene rubber or Viton. If you want more options, follow the instructions for Category E on an EPA chemical-resistance category selection chart.

All mixers*, loaders*, applicators, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical resistant gloves (except for applicators using groundboom equipment), and
- chemical resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See engineering controls for additional requirements."

10) Per the combined RED label table, the Environmental Hazard Statement for Occupational Use labels should read:

"This pesticide may be toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present,

or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.”

11) Per the combined RED label table, add the following language to the Environmental Hazards Statement for Residential Use labels:

“This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.”

12) Per the combined RED label table, the text on page 3 in the Agricultural Use box citing early entry PPE requirements must be replaced with:

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls worn over short-sleeve shirt and short pants,
- chemical-resistant footwear plus socks,
- chemical-resistant gloves made of any water-proof material,
- chemical-resistant headgear for overhead exposure,
- protective eyewear.

13) Per the combined RED label table, the following edits to label use rates must be made to the product labeling with any contradictory label language removed from the label:

- a) Throughout the label, change the maximum number of broadcast and spot applications per year to two (2).
- b) Throughout the label, the use rates must be changed to comply with those cited in the combined RED label table. Assure that use rates do not exceed the rates approved for those sites.

14) Note: The same typographical error seems to occur three times on page 12 in the Container Disposal statements for each of the three different container sizes on the label. As appropriate, add “is” between “recycling” and “unavailable” for each of the three container disposal statements to read “If recycling is unavailable, puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.”

15) Per the Label Review Manual, replace "Except as provided by applicable law" with "To the extent consistent with applicable law" in the last two sentences under Conditions of Sale and Limitations of Warranty and Liability statement on page 9.

16) Note: Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. Submit one copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

[Front Panel]

PC HERBICIDE CONCENTRATE PLUS

[For non-WPS occupational and homeowner uses] Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

[For WPS occupational use] For use on turf including turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

[For WPS occupational use and non-WPS occupational use] Follow Agricultural Use Requirements when using this product on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. †

[For homeowner use] For use only on home lawns †.

† Important: Do not apply to Bentgrass and Floratam.

Read the label and back panel before using. Protect from Freezing.

[Optional Marketing Statements:]

Specially formulated for broadleaf weeds in turf † (parks, lawns, golf courses, cemeteries, sports fields, and other grassy areas.).

Low-odor formulation.

[For homeowner use] When used as directed, kills broadleaf lawn weeds without harming most lawn grasses †.

[For all uses] KILLS BROADLEAF WEEDS INCLUDING: Dandelions, Plantains, Clover, Chickweed, Bedstraw, Black Medick, English Daisy, Curled Dock, Heal-all, Ground Ivy (Creeping Charlie), Knotweed, Ragweed, Shepherd's-Purse, plus over 35 other lawn weeds listed on side/back panel.

When used as directed, controls a wide range of broadleaf weeds in most home lawns †.

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt* 0.885%

Mecoprop-p, dimethylamine salt** 0.573%

Dicamba, dimethylamine salt*** 0.109%

OTHER INGREDIENTS: 98.433%

Total 100.000%

* 2,4-D Acid Equivalent 0.738%

** Mecoprop-p Acid Equivalent 0.478%

*** Dicamba Acid Equivalent 0.091%

ACCEPTED
with COMMENTS
in EPA Letter Dated

26 MAY 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

69526-15

KEEP OUT OF REACH OF CHILDREN

See [back] [side] panel for additional precautionary statements

FORMULATED IN CANADA FOR:
PETRO-CANADA
2489 NORTH SHERIDAN WAY
MISSISSAUGA, ONTARIO L5K 1A8

EPA REG. NO. 69526-X

EPA EST. NO. _____

NET CONTENTS: 6 fl. oz, 32 fl. oz (1 qt), 40 fl. oz, 64 fl. oz (1/2 gal), 1 Gal, 5 Gal, 30 Gal, 55 Gal or 275 Gal

Batch Code: _____

US Patent Pending

[Back/Side Panel]**PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco.

**[For WPS and non-WPS occupational uses]
Personal Protective Equipment (PPE)**

All mixers, loaders, applicators, flaggers, and other handlers must wear:

- long-sleeved shirt and long pants,
- shoes and socks, plus
- chemical-resistant gloves, when applying with any handheld nozzle or equipment, mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Some materials that are chemical-resistant to this product are Barrier Laminate, Nitrile rubber, Butyl, Neoprene rubber or Viton.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

ENGINEERING CONTROLS:

Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS
<ul style="list-style-type: none"> • Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet. • Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water. • Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. • As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. This product may adversely affect non-target plants. Do not apply to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply to areas where surface water is present, or to intertidal areas below the mean high water mark, except as noted on appropriate labels. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas, and to non-target plants. Do not apply when windy. To prevent product runoff, do not water to the point of runoff, or apply when raining or when rain is expected that day. Do not contaminate water when disposing of equipment wash waters or rinsate.

[For Homeowner use]

Rinse applicator over lawn or garden area only.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read and follow all label directions.

[WPS occupational use:]**AGRICULTURAL USE REQUIREMENTS**

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls,
- chemical-resistant gloves made of any water-proof material,
- shoes plus socks,
- protective eyewear.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated area.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

[Non-WPS occupational use:]**NON-AGRICULTURAL USE REQUIREMENT**

Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

GENERAL PRECAUTIONS AND RESTRICTIONS**[WPS and non-WPS occupational uses]**

Do not apply as a fine mist because of potential injury to desirable plants. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

[Homeowner use and non-WPS occupational uses]

People or pets may only enter treated area after sprays have dried.

[For homeowner use]

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.

Do not apply as a fine mist because of potential injury to desirable plants.

[WPS and non-WPS occupational use]**SPRAY DRIFT MANAGEMENT**

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, and relative humidity) and method of application (e.g., ground, airblast, and chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

Use only Medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles. When applying sprays that contain 2,4-D as the sole active ingredient, or when applying sprays that contain 2,4-D mixed with active ingredients that require a coarse or coarser spray, apply only as a coarse or coarser spray (ASAE standard 572) or a volume mean diameter of 385 microns or greater for spinning atomizer nozzles.

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety. This requirement does not apply to forestry or rights-of-way applications.

When applications are made with a crosswind, the swath will be displaced downwind.

The applicator must compensate for this by adjusting the path of the aircraft upwind.

Additional requirements for ground boom applications:

Do not apply with a nozzle height greater than 4 feet above the canopy.

[For WPS and non-WPS occupational uses and homeowner uses]

ADDITIONAL PRECAUTIONS AND RESTRICTIONS

- Do not apply through any type of irrigation system.
- Do not apply near lakes, streams, rivers ponds, or ditchbanks. Do not apply by air.
- Do not use more product than specified on the label.
- Do not apply when the lawn [and/or turf, and/or site] is under severe stress from drought, insect attack or high temperatures.
- Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees and other desirable plants. Do not apply in windy conditions.
- Newly seeded lawns [and/or turf, and/or site] may be treated after the third mowing.
- Do not irrigate within 24 hours of application.
- ***IMPORTANT:** Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain hybrid bermudagrasses vary in tolerance to this product. Temporary discoloration may occur, in particular on St. Augustine and Centipede.

APPLICATION DIRECTIONS

WEEDS CONTROLLED IN TURF [or, LAWN] AND ORNAMENTAL AREAS:

Bedstraw
Bindweed (field)
Birdsfoot trefoil
Black medic
Blackseed plantain
Blueweed
Brambles
Broad-leaved plantain
Buckhorn plantain
Bull thistle
Burdock
Buttercup
Canada thistle
Carolina geranium
Chickweed
Chicory
Clover
Cocklebur
Common chickweed
Crudweed
Daisy fleabane
Dandelion
Devil's paint brush (hawkweed)
Dock (curled)
English daisy
Evening primrose
Fall dandelion (fall hawkbit)
False dandelion
Fleabane
Florida pusley
Frenchweed poison oak

Grass leaved stitchwort
Ground ivy
Hawkweed
Heal-all
Henbit
Honeysuckle
Jimsonweed
Knotweed
Kochia
Kudzu
Lamb's-quarters
Little starwort
Mallow
Morningglory
Mouse-ear chickweed
Mustards
Narrow-leaved plantain
Oxalis (yellow woodsorrel)
Ox-eye daisy
Pennywort
Peppergrass
Pigweed
Pineapple weed
Plantains (narrow, buckhorn, broadleaf)
Poison ivy
Poorjoe
Povertyweed
Prostrate pigweed
Purslane
Ragweed

Russian thistle
Sandwort (thyme-leaved)
Sheep sorrel
Shepherd's purse
Smartweed (green)
Soliva
Sow thistle
Speedwell (purslane leaved)
Spotted spurge
Spreadwell (annual)
Spurge
Stitchwort
Stonecrop (mossy)
Sumac
Veronica (thyme-leaved)
Vervain
Vetch
Violet
Wild aster
Wild blackberry
Wild carrot
Wild garlic
Wild geranium
Wild lettuce
Wild onion
Wild radish
Wild raspberry
Yarrow
Yellow hawkweed

HOW TO USE

- To treat small areas, up to 1,000 sq. ft. or individual weeds use a hand or pressure sprayer.
- Apply 4 to 8 gal. of **PC Herbicide Concentrate Plus**/acre. Use higher rate when weed pressure is high, especially when there is heavy clover.
- Mix 1 part of **PC Herbicide Concentrate Plus** with 14 parts of water to make a 6.5% solution.
- Shake the spray solution before and during application.
- Aim at center of weed and spray to lightly cover. Do not overspray.
- Adjust nozzle to deliver small spray droplets for uniform coverage of the target weed.
- Weeds begin to die after 12 hours. It may take up to 4 weeks for a weed to completely die.

[Heading for WPS and non-WPS occupational uses]

TURF AND ORNAMENTAL (GOLF COURSES, CEMETERIES, PARKS, SPORTS FIELDS, TURFGRASS, LAWNS AND OTHER GRASS AREAS) (Postemergence)

[Optional heading for homeowner uses]

HOME LAWN AND ORNAMENTAL AREAS (Postemergence)

IMPORTANT: Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain hybrid bermudagrasses vary in tolerance to this product. Temporary discoloration may occur on St. Augustine and Centipede lawns.

WHEN TO USE

- Spray when weeds are young and actively growing.
- Do not water within 24 hours after spraying.
- Apply when daytime temperatures are above 45° F and below 90° F.
- Do not apply when the lawn [and/or turf] is under severe stress from drought, insect attack or high temperatures.
- Do not use more product than specified on the label.
- Newly seeded lawns may be treated after the third mowing.
- If the lawn [and/or turf] is dry, water 24 hours before using this product.
- For good turf [and/or lawn] management, normally four applications per year per treatment site are adequate. This does not include spot treatments.

[For WPS and non-WPS occupational uses]

MAXIMUM APPLICATIONS PER YEAR	APPLICATION MAXIMUM	APPLICATION INTERVAL
Do not apply more than four applications per year.	The maximum rate (8 gal./acre) for a single application is 0.747 lb ae/acre (0.896 lb DMA salt/acre). The maximum seasonal rate is 3.0 lb ae/acre, excluding spot treatments.	Repeat application at 30-day intervals if needed.

USE DIRECTIONS

For turf [lawn] other than Bentgrass and Floratam:

Dilute 1 part **PC Herbicide Concentrate Plus** with 14 parts of water to make a 6.5% solution. Spray the diluted solution at the rate of 60-120 gallons per acre. Avoid overlapping between spray patterns. Each gallon of this product, when diluted correctly, treats up to 10,900 sq. ft.

For spot treatment using hand-operated sprayers, use the maximum rate, 8 gal/acre diluted with 14 parts water to make a 6.5% solution. For small quantities use the following table:

Amount of PC Herbicide Concentrate Plus	Amount of Water to Add to Sprayer	Approximate Treated Area in Square Feet
1/2 gallon	7 gallons	2700
1 quart (1/4 gallon)	14 quarts (3.5 Gallons)	1360
1 pint (1/8 gallon)	7 quarts (1.75 gallons)	680
1 ounce	14 ounces	42

If needed repeat spot treatment after 30 days. Spot treatments are not subject to maximum seasonal rate of 3.0 lb ae/acre, but must not exceed 4 treatments per year.

Cool Season Grasses: Higher water volumes may be used when tank-mixed with a turf fertilizer. Follow fertilizer labels for proper amounts to add.

Warm Season Grasses: Centipede, St. Augustine

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses. These grasses are sensitive to 2,4-D and great care should be taken to prevent overdosing.

NOTE: Care should be taken to avoid overdosing Centipede and St. Augustine grasses or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent discoloration or injury.

Warm Season Grasses: Bahia, Common Bermuda, and Zoysia

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: Some hybrid bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.

[For homeowner use]

For turf [lawn] other than Bentgrass and Floratam:

Dilute 1 part **PC Herbicide Concentrate Plus** with 14 parts of water to make a 6.5% solution. Spray the diluted solution at the rate of 17.6 – 35.3 oz. of diluted solution on 100 sq. ft. Avoid overlapping between spray patterns.

Each gallon of this product, when diluted correctly, treats up to 10,900 sq. ft.

Warm Season Grasses: Centipede, St. Augustine

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses. These grasses are sensitive to 2,4-D and great care should be taken to prevent overdosing.

NOTE: Care should be taken to avoid overdosing Centipede and St. Augustine grasses or injury may result. Large volumes of spray water will aid in obtaining uniform coverage. If hand-type sprayers are used, it is preferable to use a single nozzle sprayer rather than a multiple nozzle boom as sideways application with a boom where the spray from more than one nozzle is allowed to fall on the same area will result in heavy local over-application and subsequent discoloration or injury.

Warm Season Grasses: Bahia, Common Bermuda, and Zoysia

Use reduced rates if grass is stressed from heat or drought. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Some temporary discoloration may occur on warm season grasses.

NOTE: Some hybrid bermudagrasses may be sensitive to this product. Contact your local extension service weed control specialist.

MASTER LABEL

14 of 17

[For WPS and non-WPS occupational uses]

TURF, GROWN FOR SEED OR SOD (Postemergence)

For selective broadleaf control in warm season and cool season turfgrass established for commercial sod production.

IMPORTANT: Do not use this product on Bentgrass and Floratam, a variety of St. Augustine grass. Certain hybrid bermudagrasses vary in tolerance to this product. Temporary discoloration may occur on St. Augustine and Centipede lawns.

MAXIMUM APPLICATIONS PER YEAR	APPLICATION MAXIMUM	APPLICATION INTERVAL
Do not apply more than four applications per year.	The maximum rate (8 gal./acre) for a single application is 0.747 lb ae/acre (0.896 lb DMA salt/acre).	Repeat application at no less than 30 days between applications.

USE DIRECTIONS

Mix 1 part of the product with 14 parts of water to make a 6.5% solution.

WHEN TO USE

- Spray when weeds are young and actively growing.
- Do not water within 24 hours after spraying.
- Apply when daytime temperatures are above 45° F and below 90° F.
- Do not apply when turf is under severe stress from drought, insect attack or high temperatures.
- Do not use more product than specified on the label.
- If the sod is dry, water 24 hours before using this product.

HOW TO USE

Follow-up applications may be required for dense infestations of perennial and biennial weeds. Do not apply this product to 'Floritam' St. Augustine grass.

NEWLY SEEDED AREAS: Applying this product to grass seedlings is recommended after the second mowing.

NEWLY SODDED, SPRIGGED, OR PLUGGED AREAS: Delay applications to newly sodded, sprigged, or plugged grasses 3 to 4 weeks after the sodding, sprigging, or plugging operations. Also, applications to dormant bermudagrass, dormant Zoysia grass, and dormant Bahiagrass are suggested.

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

USE PRECAUTIONS:

1. Do not apply this product through any type of irrigation system.
2. Avoid drift of spray mist onto vegetables, flowers, ornamental plants, shrubs, trees, and other desirable plants. Do not pour spray solutions near these plants.
3. Delay mowing 1 to 2 days before and after the application of this product.
4. Do not apply this product immediately before rainfall or irrigation. Do not irrigate or water the turfgrass within 24 hours after application.
5. Treated areas may be reseeded 3 to 4 weeks after application.

[For WPS and non-WPS occupational uses]

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place in a locked storage area. In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills. To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

[Rigid, nonrefillable containers, equal to or less than five gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If recycling unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Rigid nonrefillable containers, greater than 5 gallons:] Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If recycling unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

[Non-rigid, nonrefillable container (275-gallon bladder with frame):] Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. If recycling unavailable, puncture and dispose of in sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke. Return frame to dealer.

[For homeowner use]

STORAGE AND DISPOSAL

Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Pesticide Storage: Keep out of reach of children and animals. Keep from freezing. Keep pesticide in original container. Do not put into food or drink containers. Store in a cool, dry place, preferably in a locked storage area.

Pesticide and Container Disposal:

Nonrefillable container. Do not reuse or refill this container.

If empty: Offer for recycling if available, or place container in trash. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

It is impossible to eliminate all risks inherently associated with the use of this product. Injury to desirable plants, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Petro-Canada, the manufacturer or seller. All such risks shall be assumed by the buyer. To the extent consistent with applicable law neither Petro-Canada, the manufacturer, nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product. Except as provided by applicable law or except as warranted by this label, Petro-Canada, the manufacturer or seller makes no warranties, guarantees, or representations of any kind, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to, merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. Except as provided by applicable law, buyer's or user's exclusive remedy, and Petro-Canada, the manufacturer's or seller's total liability, shall be for damages not exceeding the cost of the product.