

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
CAUTION**

Avoid inhalation or contact with eyes or open wounds.

ENVIRONMENTAL HAZARDS

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or public waters unless this product is specifically identified and addressed in an NPDES permit. Do not discharge effluent containing this product to sewer systems without previously notifying the sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. Keep out of direct sunlight. Do not expose to organic compounds known to be toxic to metabolically active microorganisms. Avoid puncturing, rupturing, or rough handling of container such that moisture loss occurs. Keep lid tightly closed. Transfer fungus from damaged containers to prevent loss of viability. Do not use more than 2 weeks after purchase.

DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from use of this product may be disposed of on site or at an approved disposal facility. Triple rinse plastic container with detergent and water. Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or incineration, if allowed by state and local authorities, by burning.

**TECHNICAL
LAGINEX™
BIOLOGICAL LARVICIDE**

Active Ingredient: <i>Lagenidium giganteum</i> (California strain) mycelium	100%
Total	100%

Contains 10¹⁰ CFU per liter

**CAUTION:
KEEP OUT OF REACH OF
CHILDREN**

See additional precautionary statements on side panel

Registered by: **AgraQuest, Inc.**
1105 Kennedy Place, Suite 4
Davis, CA 95616

EPA Registration No.: 69592-3

EPA Establishment No.: 069592-CA-001

Net Contents: 2.2 LITERS

**FOR THE MANUFACTURE OF PRODUCTS TO
BE USED IN THE FOLLOWING AREAS**

DRAINAGE DITCHES

Activity may be reduced if water is stagnant or flowing faster than 5 meters per minute.

FLOOD PLAINS

Higher application rates should be used since these sites support high larval densities.

WILDLIFE PONDS

Excessive organic content in water may reduce activity.

TIRES, MISC. SMALL CONTAINERS

These sites should be treated with the lower rate range.

MARGINS OF RIVERS AND STREAMS

For optimal results treat quiescent areas of rivers and streams with flows not exceeding 5 meters/min.

MARSHES

Activity of this product may be reduced in water which is stagnant, organically rich, or high in salt content.

TREATMENT SITES -- CROP

**RICE
SOYBEAN
IRRIGATED PASTURE**

SUSCEPTIBLE GENERA

Aedes, Anopheles, Coquillettidea, Culex, Culiseta, Deinocerites, Eretmapodites, Haemagogus, Mansonia, Opifex, Orthopodomyia, Psorophora, Sabethes, Uranotaenia, and Wyeomyia.

ACCEPTED

JUL 16 1997

For the use of insecticide, fungicide, and
herbicide, as amended, for the
control of insects and
EPA Reg. No. 69592-3

69592-3

7-16-1997

1/1