

69592-10

08/12/2003

1/11



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division
(7511C) 401 M St., S.W.
Washington, D.C. 20460

EPA Reg.
Number:

69592-10

Date of Issuance:

8/12/03

NOTICE OF PESTICIDE:

 X Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional
Time-Limited

Name of Pesticide Product:

Rhapsody™ AS

Name and Address of Registrant (include ZIP Code):

AgraQuest, Inc.
1530 Drew Avenue
Davis, CA 95616-6320

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

The registration application referred to above, submitted in connection with registration under § 3(c)(7)(C) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you meet the following terms and conditions.

1. Comply with all requirements specified in Notice of Registration dated June 20, 2000 of EPA Reg. No 69592-4 Serenade Biofungicide Wettable Powder.
2. This registration will automatically expire on midnight July 1, 2004. EPA will reevaluate this registration before July 1, 2004 to determine whether to convert the registration to a non-expiring registration.

A stamped copy of the label is enclosed for your records.

3. Submit five(5) copies of the revised final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

Enclosure

Signature of Approving Official:

Janet L. Andersen
Janet L. Andersen, Ph.D.
Director
Biopesticides and Pollution Prevention Division (7511C)

Date:

8/12/03

Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 1

Rhapsody™ AS

AN AQUEOUS SUSPENSION Biofungicide

ACTIVE INGREDIENT

QST 713 strain of *Bacillus subtilis* . . . 1.34%

INERT INGREDIENTS 98.66%

Total 100.00%

Contains a minimum of 1×10^9 CFU/g

EPA Reg. No. 69592-XX

EPA Est. No.:

1	2	3	4	5
69592-	67545-	66728-	37429-	69592-
MEX-1	AZ-1	GA-2	GA-2	CA-1

Superscript corresponds to last digit of lot number stamped on container

U.S. Patent Nos. 6,060,051, 6,103,228, 6,291,426 and 6,417,163 on QST 713 strain of *Bacillus subtilis*

Net contents: [Agricultural: 1.0 gallon, 2.5 gallons, 3.0 gallons; Home and Garden: 1 quart]

KEEP OUT OF REACH OF CHILDREN**CAUTION**

FIRST AID

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS – Agricultural Use HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash before reuse. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- NIOSH approved respirator with any N, R, P or HE filter

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions are available, use detergent and hot water for washables. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

Wash hands before eating, drinking, chewing gum; using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS – Agricultural Use

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from treated areas.


For Agricultural Use

DIRECTIONS FOR USE – Agricultural Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For emergencies such as leaks or spills, call 24-hour toll-free CHEMTREC hotline at 1.800.424.9300



Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 2

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: coveralls, waterproof gloves, shoes plus socks.

STORAGE, DISPOSAL and SPILLS – Agricultural Use
Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in a dry area inaccessible to children. Store in original containers only. Keep container closed when not in use

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water when disposing of equipment rinsate.

CONTAINER DISPOSAL: Agricultural Use: For 1.0 gallon or 2.5-gallon or 3.0 gallon plastic containers — Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE INFORMATION – Agricultural Use

Rhapsody AS is an effective broad spectrum, preventative biofungicide for the control of many important ornamental diseases. Rhapsody AS is an ideal resistance management tool given its unique, multiple modes of action. It may be applied alone, in alternating spray programs or in tank mixes with other registered crop protection products. For maximum effectiveness, apply Rhapsody AS prior to or in the early stages of disease development

INTEGRATED PEST MANAGEMENT (IPM)

Integrate Rhapsody AS into an overall disease and pest management strategy whenever fungicides are necessary. Follow practices known to reduce disease development. Consult local agricultural authorities for specific IPM strategies developed for your crop(s) and location.

Be sure use of this product conforms to resistance management strategies, which may include rotating and/or tank-mixing with other products with different modes of action.

USE RATE DETERMINATION – Agricultural Use

Carefully read and follow all label directions, use rates and restrictions. Use maximum label rates and shortened spray intervals for conditions conducive to rapid disease development. For proper application, determine the area to be treated, the recommended label use rate and select appropriate spray volume to give good canopy penetration and coverage of plant parts to be protected. Prepare only the amount of spray solution required to treat the desired area. Accurate spray equipment calibration is essential prior to use.

PREHARVEST INTERVAL – Agricultural Use

Rhapsody AS can be applied up to and including the day of harvest.

APPLICATION INSTRUCTIONS – Agricultural Use

GROUND: Be sure to maintain agitation during mixing and application to assure uniform product suspension. Thorough coverage of all foliage is essential for effective disease control.

Rhapsody AS can be applied in commonly used ground equipment, pressurized greenhouse, and hand-held sprayers. To achieve good coverage use proper spray pressure, spray volume, nozzles, nozzle spacing and ground speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

AERIAL: Do not apply this product by air.

CHEMIGATION: Do not apply this product through any type of irrigation system.

MIXING INSTRUCTIONS – Agricultural Use

MIXING: Rhapsody AS is intended for dilution with water for spray applications. Partially fill the spray tank with clean water and begin agitation. Add the specified amount of Rhapsody AS to the tank. Finish filling the tank to the desired volume to obtain the proper spray concentration. It is critical that the spray solution be agitated during mixing and application to assure a uniform suspension. Do not allow the spray mixture

Draft 12/23/02 revised 050603, 073003, and 081103

to stand overnight or for prolonged periods. Maintain a spray solution pH between 4.5 and 8.5.

Rhapsody AS may be tank-mixed with other registered fungicides to enhance control of plant diseases. Do not exceed recommended dosage rates. Rhapsody AS cannot be mixed with any product with prohibition against such mixing. Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions.

COMPATIBILITY: To ensure compatibility do not combine Rhapsody AS in the spray tank with pesticides, surfactants or fertilizers unless prior experience has shown the combination physically compatible, effective and non-injurious under your use conditions. Do not use with penetrant-type adjuvants.

Rhapsody AS is compatible with many commonly used pesticides, fertilizer and adjuvants. However, tank-mix partners should be evaluated in a jar test as follows: Using a suitable container add proportional amounts of products to water. Add wettable powders first, followed by water dispersible granules, then liquid flowables, and lastly, emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible.

ADDITIVES: Rhapsody AS is compatible with a wide range of additives. Since the product is primarily a protectant – and thorough coverage of all above-ground plant parts is required – it is recommended that an adjuvant known to be non-phytotoxic to the plant be added to the spray tank. Rhapsody AS must not be mixed with any type of penetrant.

APPLICATION RATES

FOR USE ON ORNAMENTALS – Agricultural Use

Rhapsody AS is a protectant fungicide for control of certain foliar diseases in the field, greenhouse, interiorscape, residential and commercial landscapes, nurseries, and shade house environments.

Rhapsody AS controls foliar diseases of annual and perennial bedding plants, potted flowers, cut flowers, tropical foliage, and container grown trees and shrubs.

Apply Rhapsody AS at rates ranging from 2 to 8 quarts of product per 100 gallons of spray per acre on a 3- to 10-day schedule. Begin applications when conditions favor disease development but prior to the onset of disease. Under normal conditions apply Rhapsody AS at a rate of 2 - 4 quarts of product per 100 gallons of spray per acre on a 7-day schedule. When conditions favor severe disease development shorten the spray interval or use a higher rate. Thorough coverage is essential for effective disease control.

Rhapsody AS™: page 3

For post-harvest dip applications on cut flower crops, dip cut flowers/buds in a solution containing 6 to 25 fluid ounces of Rhapsody AS in 10 gallons of water soon after cutting. Immerse flowers for a period sufficient to provide thorough contact between cut flower/bud and the treatment solution. Use higher rates under conditions of heavy disease pressure.

Rhapsody AS has been tested for phytotoxicity on the ornamental species listed below. However, it is impossible to test this product with all species and cultivars under all conditions. A small-scale preliminary trial should be conducted to check the cultivar sensitivity before using this product on a large number of plants.

TABLE 1

PLANTS EVALUATED FOR PHYTOTOXICITY

Annual and Perennial Flowering Plants:

Alyssum	Asters	Azalea
Begonia	Calla lily	
Chrysanthemum	Cyclamen	Dianthus
Dwarf Bee-Balm		Easter lily
Garden phlox	Geraniums	Gerbera
Golden star	Hydrangea	Impatiens
Kalanchoe	Lanaria	Lisianthis
Lobelia	Marigolds	Orchids
Pansies	Petunia	Poinsettia
Portulaca	Ranunculus	Roses
Salvia spp.	Snapdragons	Stock
Verbena spp.	Vinca	Violas
Zinnias		

Tropical foliage.

Aglonemea	Dieffenbachia
Dracaena spp.	English Ivy
Hibiscus	Spathiphyllum

Trees and Shrubs:

Azalea	Boxwood
Crape myrtle	Dogwood
Gumbo azalea	Indian Hawthorn
Japanese maple	Legustrum japonicum
Lilac	Loropetalum
Photinia	Rhododendron
Rosaceae spp.	Soft Touch Holly
Spirea	

Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 4

TABLE 2ORNAMENTAL DISEASES CONTROLLED BY
RHAPSODY ASAnthracnose -- *Colletotrichum* spp.Bacteria -- *Erwinia*, *Pseudomonas*, *Xanthomonas* spp.Black spot of rose -- *Diplocarpon rosea*Botrytis -- *Botrytis cinerea*Leaf spots -- *Alternaria*, *Cercospora*, *Entomosporium*,
Helminthosporium, *Myrothecium*, *Septoria* spp.Powdery mildew -- *Erysiphe*, *Oidium*, *Podosphaera*,
Sphaerotheca spp.Scab -- *Venturia* spp

Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 5

Recommended Application Rates for Selected Greenhouse Crops –Agricultural Use
(Rhapsody AS has a 0-Day PreHarvest Interval for all crops contained on this label)

Green-house Crops	Diseases	Rate/Acre Qts./100 gallons spray mix	Use Recommendations
Broccoli	Pin Rot <i>Alternaria/Xanthomonas</i> complex	3 - 6	For suppression, begin applications when environmental conditions in the greenhouse are conducive to rapid disease development. Continue sprays at 7- day intervals. Thorough coverage is essential. Do not apply more than 6 qts. Rhapsody AS per acre per application.
Carrot	Black Root Rot/Black Crow <i>Alternaria</i> spp.	2 - 4	Begin applications when environmental conditions in the greenhouse are conducive to rapid disease development. Continue sprays at 7- day intervals or as needed. Thorough coverage is essential. Do not apply more than 4 qts. Rhapsody AS per acre per application.
	Bacterial Leaf Blight <i>Xanthomonas campestris</i>	6 - 10	Begin applications when environmental conditions in the greenhouse are conducive to rapid disease development. Continue sprays at 7-day intervals or as needed. Use high rates and shorter intervals when conditions are conducive to rapid disease development. Thorough coverage is essential. Do not apply more than 10 qts. Rhapsody AS per acre per application.
Cucurbits	Powdery Mildew <i>Erysiphe</i> spp. <i>Sphaerotheca</i> spp.	4 - 8	Begin applications soon after emergence or transplant and continue on a 7-day interval or as needed. When environmental conditions in the greenhouse and plant stage are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a rotational program with other registered fungicides. Continue sprays at 7- day intervals or as needed. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.
Leafy Vegetables (except Brassica)	Downy Mildew <i>Bremia lactucae</i> <i>Peronospora</i> spp.	3 - 6	For suppression, use as a preventative treatment in early crop stages in a tank mix or alternating spray program with other registered fungicides. Apply as a foliar spray and begin applications when conditions favor disease development. Continue sprays at 7- to 10- day intervals or as needed. Apply in sufficient water to ensure complete coverage of entire plant. When conditions are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a rotational program with other registered fungicides for downy mildew and powdery mildew control. Do not apply more than 6 qts. Rhapsody AS per acre per application.
	Powdery Mildew <i>Erysiphe cichoracearum</i>		
	Sclerotinia head and leaf drop <i>Sclerotinia</i> spp.	4 - 6	<p><u>For control of early Sclerotinia head and leaf drop:</u> Apply at planting or immediately following planting but prior to crop emergence as a 4- to 6-inch seed line treatment. Make a second application as a directed spray with multiple nozzles per each seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Additional applications should be made on 10- to 14-day intervals if conditions for disease development persist. Use higher rates under conditions conducive to moderate to severe disease pressure. Light irrigation after application to incorporate the product may improve disease control. Thorough coverage is essential. Do not apply more than 6 qts. Rhapsody AS per acre per application.</p> <p align="center">OR</p> <p><u>For control of Sclerotinia head and leaf drop:</u> Apply as a directed spray with multiple nozzles per each seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning or transplanting. Additional applications should be made on 10- to 14-day intervals if conditions for disease development persist. Use higher rates under conditions conducive to moderate to severe disease pressure. Light irrigation after application to incorporate the product may improve disease control. Thorough coverage is essential. Do not apply more than 6 qts. Rhapsody AS per acre per application.</p>

Green-house Crops	Diseases	Rate/Acre Qts./100 gallons spray mix	Use Recommendations
Onion	Botrytis neck rot <i>Botrytis</i> spp. Botrytis Leaf Blight <i>Botrytis squamosa</i> Onion Purple Blotch <i>Alternaria porri</i> Onion Downy Mildew <i>Peronospora destructor</i>	6 - 10	Begin applications when environmental conditions in the greenhouse are conducive to rapid disease development. Continue sprays at 7-day intervals or as needed. When conditions in the greenhouse are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a rotational program with other registered fungicides for Botrytis neck rot control. Thorough coverage is essential. Do not apply more than 10 qts. Rhapsody AS per acre per application.
Garlic	Botrytis Neck Rot <i>Botrytis</i> spp.	6-10	Begin applications when environmental conditions in the greenhouse are conducive to rapid disease development. Continue sprays at 7-day intervals or as needed. Under moderate to severe disease pressure, it is recommended that Rhapsody AS be used in a rotational program with other registered fungicides for Botrytis neck rot control. Thorough coverage is essential. Do not apply more than 10 qts. Rhapsody AS per acre per application.
	Rust <i>Puccinia porri</i>	2 - 6	For suppression, it is recommended that Rhapsody AS be used as a part of a preventative disease control program in alternation or tank mix with other registered fungicides. Continue applications as needed. Thorough coverage is essential. Do not apply more than 6 qts. Rhapsody AS per acre per application.
Pepper	Gray mold <i>Botrytis cinerea</i>	4 - 8	Begin applications soon after emergence or transplant and continue on a 7-day interval or as needed. When environmental conditions in the greenhouse are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a rotational program with other registered fungicides. Continue sprays at 7-day intervals or as needed. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.
	Powdery mildew <i>Oidiopsis taunca</i>	6 - 8	For suppression, begin applications soon after emergence or transplant and continue on a 7-day interval or as needed. When environmental conditions in the greenhouse are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a rotational program with other registered fungicides. Continue sprays at 7-day intervals or as needed. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.
	Bacterial Spot <i>Xanthomonas</i> spp.	2 - 8	Begin applications soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a 5- to 7-day interval or as needed. When conditions are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a tank mix program with copper-based bactericides registered for control of bacterial spot at labeled rates. Continue sprays at 7-day intervals or as needed. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.

8/11

Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 7

Green-house Crops	Diseases	Rate/Acre Qts./100 gallons spray mix	Use Recommendations
Tomato	Early Blight <i>Alternaria solani</i> Late Blight <i>Phytophthora infestans</i>	4 - 8	For suppression, begin applications when plants are 4- to 6-inches high. Repeat applications at 5- to 7-day intervals or as needed. It is recommended that Rhapsody AS be used in a rotational program with other registered fungicides for foliar disease control. Use shorter spray intervals under conditions conducive to rapid disease development. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.
	Powdery Mildew <i>Leveillula taurica</i>	4 - 8	Begin applications soon after emergence or transplant and when environmental conditions in the greenhouse are conducive to rapid disease development. Continue applications on 5- to 7- day intervals or as needed. Use maximum label rates under conditions conducive to rapid disease development. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.
	Bacterial Spot <i>Xanthomonas</i> spp.	2 - 8	Begin applications soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a 5- to 7-day interval or as needed. When conditions are conducive to rapid disease development, it is recommended that Rhapsody AS be used in a tank mix program with copper-based bactericides registered for control of bacterial spot at labeled rates. Continue sprays at 7-day intervals or as needed. Thorough coverage is essential. Do not apply more than 8 qts. Rhapsody AS per acre per application.

Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 8

For Home and Garden Use

PRECAUTIONARY STATEMENTS— Home and Garden

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water. Get medical attention if irritation persists. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals

ENVIRONMENTAL HAZARDS — Home and Garden

Do not apply directly to water. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE — Home and Garden

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE INFORMATION — Home and Garden

Rhapsody AS is a broad spectrum, preventative biofungicide recommended for the control or suppression of many important plant diseases (see Tables 2 and 3).

Rhapsody AS can be applied in commonly used ground equipment, hose-end, and hand-held sprayers.

MIXING AND APPLICATION INSTRUCTIONS — Home and Garden

Prepare only the amount of spray solution required to treat the desired area. Accurate spray equipment calibration is essential prior to use. Mix thoroughly and keep spray solution agitated during application. Do not allow the spray mixture to stand overnight or for prolonged periods.

MIXING: —

For Apples/Pears, Broccoli, Carrot, Cherries, Cucurbits, Grapes, Leafy Vegetables, Onions/Garlic, Pepper, Tomato, and Walnuts, mix 4 fluid oz of Rhapsody AS per gallon of water.

For Ornamental Plants (such as those in Table 1), mix 1 - 2 fluid ounces of Rhapsody AS per gallon of water. Use the higher rate under conditions of high disease pressure

APPLICATION: Use 15 fluid ounces of spray solution per 100 square feet. Spray to ensure thorough coverage of the plant. For best results, treat prior to disease development or at the first sign of disease infection. Repeat at 7-day intervals or as needed. Under conditions of high disease pressure, shorten the spray interval or, for ornamental plants, use the higher

rate.

Rhapsody AS can be applied up to and including the day of harvest.

Hose-End Sprayer Application Instructions:

For Apples/Pears, Broccoli, Carrot, Cherries, Cucurbits, Grapes, Leafy Vegetables, Onions/Garlic, Pepper, Tomato, and Walnuts, apply 4 fluid ounces of Rhapsody AS per 1000 square feet.

For Ornamental Plants — Annual and Perennial Flowering Plants, Tropical Foliage, Trees and Shrubs, apply 2 fluid ounces of Rhapsody AS per 1000 square feet.

TABLE 1

PLANTS EVALUATED FOR PHYTOTOXICITY

Annual and Perennial Flowering Plants:

Alyssum	Asters	Azalea
Begonia	Calla lily	
Chrysanthemum	Cyclamen	Dianthus
Dwarf Bee-Balm		Easter lily
Garden phlox	Geraniums	Gerbera
Golden star	Hydrangea	Impatiens
Kalanchoe	Lanaria	Lisianthus
Lobelia	Marigolds	Orchids
Pansies	Petunia	Poinsettia
Portulaca	Ranunculus	Roses
Salvia spp.	Snapdragons	Stock
Verbena spp.	Vinca	Violas
Zinnias		

Tropical foliage:

Aglonemea	Dieffenbachia
Dracaena spp	English Ivy
Hibiscus	Spathiphyllum

Trees and Shrubs:

Azalea	Boxwood
Crape myrtle	Dogwood
Gumbo azalea	Indian Hawthorn
Japanese maple	Legustrum japonicum
Lilac	Loropetalum
Photinia	Rhododendron
Rosaceae spp.	Soft Touch Holly
Spirea	

10/11

Draft 12/23/02 revised 050603, 073003, and 081103

Rhapsody AS™ page 9

TABLE 2**ORNAMENTAL DISEASES CONTROLLED BY
RHAPSODY AS**Anthracnose – *Colletotrichum* spp.Bacteria – *Erwinia*, *Psuedomonas*, *Xanthomonas* spp.Black spot of rose – *Diplocarpon rosea*Botrytis – *Botrytis cinerea*Leaf spots – *Alternaria*, *Cercospora*, *Entomosporium*,
Helminthosporium, *Myrothecium*, *Septoria* spp.Powdery mildew – *Erysiphe*, *Oidium*, *Podosphaera*,
Sphaerotheca spp.Scab – *Venturia* spp.**TABLE 3****VEGETABLE AND FRUIT DISEASES CONTROLLED
OR SUPPRESSED BY RHAPSODY AS**Bacterial Leaf Blight (*Xanthomonas campestris*)Bacterial Speck (*Psuedomonas syringae* pv. Tomato)Bacterial Spot (*Xanthomonas* spp.) — suppressionBlack Mold (*Alternaria alternata*)Black Root Rot/Black Crown Rot (*Alternaria* spp.)Botrytis Neck Rot (*Botrytis* spp.)Downy Mildew (*Bremia lactucae*, *Peronospora* spp.)
and (*Plasmopara viticola* — suppression)Early Blight (*Alternaria solani*)Fire Blight (*Erwinia amylovora*) — suppressionGray Mold (*Botrytis cinerea*)Late Blight (*Phytophthora infestans*)Pin Rot (*Alternaria/Xanthomonas* complex) —
suppressionPowdery Mildew (*Uncinula necator*, *Erysiphe* spp.,
Sphaerotheca spp., *Oidiopsis taunca*, *Leveillula taurica*,
Podosphaera leucotricha)Scab (*Venturia* spp.) — suppressionWalnut Blight (*Xanthomonas campestris*)

STORAGE AND DISPOSAL – Home and Garden

STORAGE: Store in a dry area inaccessible to children. Store in original container only. Keep container closed when not in use

CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS FOR SALE AND WARRANTY

AgraQuest warrants to those persons lawfully purchasing this product that at the time of the first sale of this product by Seller that this product conformed to its description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions. Buyers and users of this product assume the risk of any use contrary to such directions. EXCEPT AS PROVIDED ELSEWHERE IN WRITING CONTAINING AN EXPRESS REFERENCE TO THIS WARRANTY AND LIMITATION OF DAMAGES, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OR GUARANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. To the extent permitted by state law, the Seller's liability for any breach of warranty shall not exceed the purchase price of the material as to which a claim is made.

To the extent permitted by state law, Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, or without the fault or negligence of the Seller, or from failure to follow the label.

Rhapsody AS is a registered trademark of AgraQuest, Inc.

© Copyright AgraQuest, Inc., 2003

AgraQuest, Inc.
1530 Drew Avenue
Davis, California 95616
www.agraquest.com

