

69592-11

9/9/2004

Page 1 of 2

07 September 2004

SERENADE® MAX™ MASTER LABEL

Serenade® MAX™

A Wettable Powder Biofungicide

[Optional/Alternate Statement:

“NOP Logo: For Organic Production”]

[Optional/Alternate Statement: “NOP Logo:

Can be Used for Organic Production”]

ACTIVE INGREDIENT

QST 713 strain of dried *Bacillus subtilis* 14.6%

OTHER INGREDIENTS 85.4%

Total 100.0%

Contains a minimum of 7.3×10^9 cfu/g

EPA Reg. No. 69592-X

EPA Est. No.:

| 1 | 2 | 3 | 4 | 5 |
|--------|--------|--------|--------|--------|
| 69592- | 67545- | 66728- | 37429- | 69592- |
| MEX-1 | AZ-1 | GA-2 | GA-2 | CA-1 |

Superscript corresponds to last digit of lot number stamped on container

U.S. Patent Nos. 6,060,051, 6,103,228, 6,291,426, and 6,417,163 on QST 713 strain of *Bacillus subtilis*

Net weight:

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID – Agricultural Use

IF ON SKIN: Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice. Have the product label with you when calling a poison control center or doctor.

[See attached label booklet for First Aid, Precautionary Statements and Directions for Use.]

ACCEPTED

SEP 09 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 69592-11

PRECAUTIONARY STATEMENTS– Agricultural Use

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water. Get medical attention if irritation persists. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. May be harmful if inhaled. Avoid breathing spray mist. Remove contaminated clothing and wash clothing before use.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- NIOSH approved respirator with any N, P, R or HE filter.

Follow manufacturer’s instructions for cleaning and maintaining PPE. If no instructions are available, use detergent and hot water for washables. Keep and wash PPE separately from other laundry.

[OPTIONAL STATEMENT: When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides, the handler PPE requirements may be reduced or modified as specified in the WPS.]

USER SAFETY RECOMMENDATIONS

Users should:
Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
Remove clothing/PPE immediately if pesticides get inside. Then wash thoroughly and put on clean clothing.
Remove PPE immediately after handling this product.
Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS– Agricultural Use

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Do not apply when weather conditions favor drift or runoff from treated areas.

For Agricultural Use

DIRECTIONS FOR USE – Agricultural Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For emergencies such as leaks or spills, call 24-hour toll-free CHEMTREC hotline at 1.800.424.9300.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is: Long sleeved shirts and long pants pants, waterproof gloves, shoes plus socks.

STORAGE, DISPOSAL & SPILLS – Agricultural Use

Do not contaminate water, food, or feed by storage and disposal.

STORAGE: Store in a dry area inaccessible to children. Store in original containers only. Keep container closed when not in use.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Do not contaminate water when disposing of equipment rinsate.

CONTAINER DISPOSAL:

For 1000 lb. bulk bag with liner: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Dispose of liner in a sanitary landfill or by incineration if allowed by State and local authorities. If burned stay out of smoke. Return empty bulk bag to manufacturer for reuse. If bulk bag is contaminated and cannot be reused, dispose of in the same manner as liner.

For all other agricultural use containers: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

GENERAL USE INFORMATION– Agricultural Use

Serenade MAX is a broad spectrum, preventative product for the control or suppression of many important plant diseases. Serenade MAX may be applied as a foliar spray alone, in alternating spray programs or in tank mixes with other registered crop protection products. When conditions are conducive to heavy disease pressure, use Serenade MAX in a rotational program with other registered fungicides. Serenade MAX may be applied with spray equipment commonly used for making ground or aerial applications and sprinkler/irrigation systems commonly used for chemigation. Serenade MAX can be used for organic production.

INTEGRATED PEST MANAGEMENT (IPM)

Integrate Serenade MAX into an overall disease and pest management strategy whenever fungicide use is necessary. Follow practices known to reduce disease development. Consult local agricultural authorities for specific IPM strategies developed for your crop(s) and location.

USE RATE DETERMINATION – Agricultural Use

Carefully read and follow all label directions, use rates and restrictions. Serenade MAX should be applied prior to or in the early stages of disease development. Use maximum label rates and shortened spray intervals for conditions conducive to threatening or rapid disease development. For proper application, determine the number of acres to be treated, the recommended label use rate and select appropriate gallonage to give good canopy penetration and coverage of plant parts to be protected. Prepare only the amount of spray solution required to treat the measured acreage. Accurate spray equipment calibration is essential prior to use.

PREHARVEST INTERVAL – Agricultural Use

Serenade MAX can be applied up to and including the day of harvest.

APPLICATION INSTRUCTIONS – Agricultural Use

GENERAL: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower/treatment coordinator are responsible for considering all of these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

GROUND: Thorough coverage is essential for optimum disease control. To achieve good coverage use proper spray pressure, gallonage per acre, nozzles, nozzle spacing and ground speed. Consult spray nozzle and accessory catalogues for specific information on proper equipment calibration.

AERIAL: This product can be applied by aerial application. Refer to the Aerial Drift Reduction Advisory Information section of this label for general directions and precautions. Use the application rate indicated for the appropriate crop in sufficient water to achieve thorough coverage, or a minimum of 3 gallons of water per acre.

CHEMIGATION DIRECTIONS FOR USE

CHEMIGATION: This product can be applied through sprinkler or drip type irrigation systems, including a center pivot, lateral move, end tow, side wheel roll, traveler, solid set, and hand move. Refer to the Chemigation Directions for Use section of this label for general directions and precautions. Use the application rate indicated for the appropriate crop as specified in the Use Recommendations section of this label.

MIXING INSTRUCTIONS – Agricultural Use

MIXING: Serenade MAX must be diluted with water for Spray applications and may be used in spray equipment commonly used for making ground applications. Partially fill the spray tank with clean water and begin agitation. Add the specified amount of Serenade MAX to the tank. Finish filling the tank to the desired volume to obtain the proper spray concentration. Maintain agitation continuously while spraying. Do not allow spray mixture to stand overnight or for prolonged periods. [Optional Statement: Maintain a spray solution pH between 4.5 and 8.5].

Serenade MAX may be tank mixed with other registered pesticides to enhance plant disease control. Do not exceed recommended dosage rates. This product cannot be mixed with any product with prohibition against such mixing. Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions.

COMPATIBILITY: Do not combine Serenade MAX in the spray tank with pesticides, surfactants or fertilizers if there has been no previous experience or use of the combination to show it is physically compatible, effective and non-injurious under your use conditions.

Serenade MAX is compatible with many commonly used pesticides, fertilizers, adjuvants and surfactants but has not been fully evaluated with all of these. To ensure compatibility of tank-mix combinations they should be evaluated prior to use, as follows: Using a suitable container add proportional amounts of product to water. Add wettable powders first, followed by water dispersible granules, then by liquid flowables and lastly, emulsifiable concentrates. Mix thoroughly and let stand for at least five minutes. If the combination stays mixed or can be remixed, it is physically compatible. Test the combination on a small portion of the crop to be treated to ensure that a phytotoxic response will not occur as a result of application. [OPTIONAL STATEMENT: Do not use with penetrant-type adjuvants.]

ADDITIVES: Serenade MAX is compatible with a wide range of additives. Since the product is primarily a protectant, thorough coverage of all above-ground plant parts is required for effective product performance. To improve plant surface coverage, add a nonphytotoxic adjuvant [such as Biotune™] to spray tank.

General Requirements:

- 1) Apply this product through sprinkler or drip type irrigation systems including center pivot, lateral move, end tow, side wheel roll, traveler, solid set or hand move systems. Do not apply this product through any other type of irrigation system.
- 2) Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- 3) Ensure that the irrigation system used is properly calibrated and if you have questions, call the State Extension Service specialists or the equipment manufacturer.
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.

Equipment Requirements:

- 1) Public water supply means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of 25 individuals daily at least 60 days throughout the year.
- 2) Chemigation systems connected to the public water systems must contain a functional, reduced-pressure zone (RPZ), backflow preventer or the functional equivalent in the water supply upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top of the overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- 3) The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- 4) The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back towards the injection pump.
- 5) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 6) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected

- 7) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 8) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.
- 9) Do not apply when wind speed favors drift beyond the area intended for treatment

Application Instructions:

- 1) Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues may cause product to lose effectiveness or strength.
- 2) Do not combine Serenade MAX with pesticides, surfactants or fertilizers for application through chemigation equipment unless prior experience has shown the combination physically compatible, effective and non-injurious under conditions of use. Serenade MAX has not been fully evaluated for compatibility with all adjuvants or surfactants. It is advisable to conduct a spray compatibility test if mixture with adjuvants or surfactants is planned.

Center-pivot, Lateral Move, End Tow, and Traveler Irrigation Equipment (Use only with electric or oil hydraulic drive systems which provide a uniform water distribution):

- Determine size of area to be treated.
- Determine the time required to apply no more than 1/4 inch of water (6,750 gallons water per acre) over the area to be treated when the system and injection equipment are operated at normal pressures recommended by the equipment manufacturer. Run system at 80 to 95% of manufacturer's rated capacity.
- Using only water, determine the injection pump output when operated at normal line pressure.
- Determine the amount of Serenade MAX fungicide required to treat area.
- Add required amount of Serenade MAX fungicide and sufficient water to meet the injection, time requirements of the solution tank.
- Maintain constant solution tank agitation during the injection period.
- Stop injection equipment after treatment is completed. Continue to operate the system until Serenade MAX fungicide solution has cleared the sprinkler head.

Solid-set, Slide (wheel) Roll, and Hand Move Irrigation Equipment:

- Determine acreage covered by sprinkler.
- Fill injector solution tank with water and adjust flow rate to use contents over a 10- to 30-minute interval.
- Determine the amount of Serenade MAX fungicide required to treat area.

- Add the required amount of Serenade MAX fungicide into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.
- Inject Serenade MAX fungicide at the end of the irrigation cycle or as a separate application to maximize foliar fungicide retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until Serenade MAX fungicide solution has cleared the last sprinkler head.

AERIAL DRIFT REDUCTION ADVISORY INFORMATION

General: Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed. Note: This section is advisory in nature and does not supersede the mandatory label requirements.

INFORMATION ON DROPLET SIZE: The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE: Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets. Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When high flow rates are needed, use higher flow rate nozzles instead of increasing pressure. # of Nozzles - Use the minimum number of nozzles that provide uniform coverage. Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential. Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM WIDTH: For aerial applications, the boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3 – 10 mph as measured by an anemometer. Use medium or coarser spray according to

07 September 2004

SERENADE® MAX™ MASTER LABEL

5 8 25

ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or the crop canopy.

recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

APPLICATION HEIGHT: Do not make application at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT: When applications are made with a crosswind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.).

WIND: Drift potential is lowest between wind speeds of 2 - 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. **NOTE:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY: When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS: Do not apply during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS: The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from the sensitive areas). Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and

Recommended Application Rates for Selected Crops – Agricultural Use

Serenade MAX has a 0-Day PreHarvest Interval for all crops contained on this label

Under moderate to severe disease pressure, for improved performance, increase rates and reduce spray intervals or use Serenade MAX in a tank mix or rotational program with other registered fungicides.

| Crops | Disease | Rate lbs./acre* | Application Instructions |
|---|---|--------------------|--|
| Artichoke | Powdery Mildew <i>Leveillula taurica</i> , <i>Erysiphe cichoracearum</i> | 1 - 3 | Begin application when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. Serenade MAX may be applied up to and including the day of harvest. |
| Asparagus | Rust <i>Puccinia asparagi</i> Botrytis Blight <i>Botrytis cinerea</i> | 1 - 3 | Begin application soon after emergence and when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. Serenade MAX may be applied up to and including the day of harvest. |
| Bananas Plantains | Sigatoka <i>Mycosphaerella spp.</i> | 1 - 3 | Begin application when leaves first appear and repeat on <u>7 to 21</u> day intervals or as needed. The addition of an approved emulsifiable oil to spray solutions will improve performance. |
| Bushberry, Caneberry and other berry crops Blueberries Blackberry Raspberry Loganberry Huckleberry Cranberry Gooseberry Elderberry Currant Caneberry and other berry crops | Mummy Berry <i>Monilinia vaccinii-corymbosi</i> Anthracnose Fruit Rot <i>Colletotrichum gloeosporiodes</i> Botrytis Blight <i>Botrytis cinerea</i> Leaf Rust <i>Pucciniastrum vaccinii</i> Powdery Mildew <i>Microsphaera alni</i> Sooty Mold Misc. fungi Alternaria Fruit Rot <i>Alternaria tenuissima</i> Bacterial Canker <i>Pseudomonas spp.</i> | 1 - 3 | Mummy Berry - For suppression, begin application at the bud break stage of development and repeat on a <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for mummy berry control. Bacterial Canker – Apply before fall rains and again during dormancy before spring growth. For all other diseases – Begin application prior to disease development and repeat on a <u>7 to 10</u> day interval or as needed. Cranberries – Make application to non-flooded fields only. Serenade MAX may be applied to fruit up to and including the day of harvest. |

* Rate presented in lbs./acre unless otherwise noted.

| Crops | Disease | Rate lbs./acre [#] | Application Instructions |
|--|--|--------------------------------|---|
| Brassica vegetables (Cole Crops) Broccoli Cabbage Cauliflower Brussels Sprouts Collards Kale Mustard Greens Kohlrabi and other brassica crops | Pin Rot Complex <i>Alternaria/Xanthomonas</i> Bacterial Leaf Spot <i>Pseudomonas syringae</i> Bacterial Soft Rot <i>Erwinia / Pseudomonas</i> Black Rot <i>Xanthomonas campestris</i> Xanthomonas Leaf Spot <i>Xanthomonas campestris</i> Alternaria Leaf Spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum higginsianum</i> Cercospora Leaf Spot <i>Cercospora brassicaicola</i> Downy Mildew <i>Peronospora parasitica</i> <i>Peronospora</i> spp. Powdery Mildew <i>Erysiphe polygoni</i> Southern Blight <i>Sclerotium rolfsii</i> | 1 - 3 | Pin Rot - For suppression, begin application when environmental conditions are conducive to disease development and repeat on <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for pin rot control. For all other diseases - Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed. |
| Bulb Vegetables Onion Garlic Shallots and other bulb vegetables | Botrytis Neck Rot <i>Botrytis</i> spp. Botrytis Leaf Blight <i>Botrytis squamosa</i> Onion Purple Blotch <i>Alternaria porri</i> Onion Downy Mildew <i>Peronospora destructor</i> Downy Mildew <i>Peronospora</i> spp. Powdery Mildew <i>Erysiphe</i> spp. | 1 - 3 | Begin application when environmental conditions are conducive to disease development and repeat on a <u>7 to 10</u> day interval or as needed. Apply sufficient water to provide complete coverage of plants. |
| | Rust <i>Puccinia porri</i> | 1 - 3 | For suppression, begin application when conditions are conducive to disease development and repeat on a <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for rust control. |

Rate presented in lbs./acre unless otherwise noted.

| Crops | Disease | Rate lbs./acre [#] | Application Instructions |
|--|--|--------------------------------|--|
| Cereal Grains Barley Corn Millets Oat Rice Rye Sorghum Triticale Wheat and other cereal grain crops | Powdery Mildew <i>Erysiphe graminis</i> Rust <i>Puccinia</i> spp. Blast <i>Pyricularia oryzae</i> Sheath Spot and Blight <i>Rhizoctonia oryzae</i> <i>Thanatephorus kernel</i> Smut <i>Tilletia barclayana</i> Bacterial Blight and Streak <i>Xanthomonas</i> spp Stem Rot <i>Sclerotium oryzae</i> <i>Magnaporthe</i> spp Brown Rot, Leaf Spots and Smuts <i>Cerccospora</i> spp <i>Entyloma</i> spp <i>Dreschlera</i> spp <i>Cochliobolus</i> spp <i>Ceratobasidium</i> spp | 1 -3 | Begin applications when environmental conditions and plant stage are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure. |
| Citrus Orange Grapefruit Lemon Tangerine Tangelo Pummelo and other citrus crops | Greasy spot <i>Mycosphaerella citri</i> Post Bloom Fruit Drop <i>Colletotrichum acutatum</i> Scab <i>Elsinoe fawcetti</i> Melanose <i>Diaporthe citri</i> Alternaria Leaf Spot <i>Alternaria alternata</i> | 1 -3 | <p>Greasy spot - For suppression, begin applications at first new foliar flush, and repeat with subsequent new flushes. When conditions are conducive to rapid disease development, Serenade must be used in a tank mix program with other registered products, such as spray oil or copper-based fungicides, at labeled rates.</p> <p>Post bloom fruit drop – For suppression, begin applications at early bloom and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed. Utilize the shorter spray interval between applications if warm, wet conditions persist.</p> <p>Citrus scab – For suppression, begin applications at first new foliar flush and repeat at petal fall and at ½ inch diameter fruit.</p> <p>Melanose – For suppression, begin applications at petal fall and repeat on a <u>14 to 21</u> day interval until fruit becomes resistant.</p> <p>Alternarial Leaf Spot – Begin applications when environmental conditions and plant stage are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed.</p> <p>For improved performance on post bloom fruit drop, scab and melanose, use Serenade in a tank mix or rotational program with other registered fungicides.</p> |
| Crops | Disease | Rate lbs./acre [#] | Application Instructions |

| | | | |
|--|---|--------------|--|
| <p>Corn Sweet Corn Popcorn Seed Corn and other corn crops</p> | <p>Common rust <i>Puccinia sorghi</i> Northern Leaf Blight <i>Exserohilum turcicum</i> Helminthosporium turcium Southern Leaf Blight <i>Bipolaris maydis</i> <i>Helminthosporium maydi</i> <i>Cochliobolus heterostrophus</i></p> | <p>1 - 3</p> | <p>Begin applications when environmental conditions are conducive to disease development. Continue applications on <u>7 to 10</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure.</p> |
| <p>Cucurbits Cucumber Cantaloupe Melon Muskmelon Squash Watermelon and other cucurbit crops</p> | <p>Powdery Mildew <i>Erysiphe</i> spp. <i>Sphaerotheca</i> spp. Gummy Stem Blight <i>Didymella bryoniae</i> <i>Phoma cucurbitacearum</i> Angular Leaf Spot <i>Pseudomonas syringae</i> Anthracnose <i>Colletotrichum lagenarium</i> Downy Mildew <i>Pseudoperonospora cubensis</i> Bacterial Fruit Blotch <i>Acidovorax avenaesubsp</i></p> | <p>1 - 3</p> | <p>Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day interval or as needed.</p> |
| <p>Fruiting Vegetables Pepper Tomato Eggplant Ground Cherry Tomatillo Okra and other fruiting vegetables</p> | <p>Bacterial Spot <i>Xanthomonas</i> spp.</p> | <p>1 - 3</p> | <p>Begin application soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a <u>5 to 7</u> day interval or as needed. When conditions are conducive to rapid disease development, for improved control, use Serenade in a tank mix program with copper-based bactericides registered for control of bacterial spot at labeled rates.</p> |
| | <p>Bacterial Speck <i>Pseudomonas syringae pv tomato</i></p> | <p>1 - 3</p> | <p>Begin application soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a <u>5 to 7</u> day interval or as needed. Use higher rates when conditions are conducive to rapid disease development.</p> |
| | <p>Early Blight <i>Alternaria solani</i> Late Blight <i>Phytophthora infestans</i></p> | <p>1 - 3</p> | <p>For suppression, begin application when plants are 4- to 6-inches high. Repeat applications on a <u>5 to 7</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for late blight control.</p> |
| | <p>Powdery Mildew <i>Leveillula taurica</i> <i>Oidiopsis taurica</i> <i>Erysiphe</i> spp. <i>Sphaerotheca</i> spp. Downy Mildew <i>Pseudoperonospera cubensis</i></p> | <p>1 - 3</p> | <p>For suppression, begin application soon after emergence or transplant and continue on a <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for powdery mildew control.</p> |
| | <p>Gray Mold <i>Botrytis cinerea</i></p> | <p>1 - 3</p> | <p>Begin application soon after emergence or transplant and repeat on a <u>7 to 10</u> day interval or as needed.</p> |

Rate presented in lbs./acre unless otherwise noted.

| Crops | Disease | Rate lbs./acre# | Application Instructions |
|-------|---------|-----------------|--------------------------|
|-------|---------|-----------------|--------------------------|

| | | | |
|---------------|---|-----------------------|--|
| Grape | Gray Mold <i>Botrytis cinerea</i> Sour Rot [a complex of pathogens <i>Aspergillus niger, Alternaria tenuis, Botrytis cinerea, Cladosporium herbarum, Rhizopus arrhizus, Penicillium sp.</i> , and others] | 1 - 3 | Begin application at bloom, before bunch closure, at veraison and preharvest. Serenade may be applied to fruit up to and including the day of harvest. |
| | Powdery Mildew <i>Uncinula necator</i> | 1 - 3 | Begin application when new shoots are ½- to 1½- inches long. Repeat when shoots are 3- to 5-inches long, when shoots are 8- to 10-inches long and then at 7 to 10 day intervals until disease conditions no longer exist. |
| | Downy Mildew <i>Plasmopara viticola</i> | 1 - 3 | For suppression, apply at 10-inch shoot, then at 7-day intervals until bunch closure (berry touch). For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for downy mildew control. |
| | Phomopsis <i>Phomopsis viticola</i> | 1 - 3 | Begin applications when shoots are ½ to 1 inch long and repeat when shoots are 6-8 inches long. |
| | Eutypha <i>Eutypha lata</i> | 2 - 5% | Apply solution to pruning wounds. Sanitation is critical. All wood from infected plants must be removed from the vineyard and destroyed (either buried or burned). |
| Herbs/ Spices | Bacterial Blight <i>Pseudomonas syringae</i> Anthracnose <i>Colletotricum spp.</i> Alternaria Leaf Blight <i>Alternaria spp.</i> | 1 - 3 | Begin application when environmental conditions are conducive to disease development. Repeat on 7 to 10 day interval or as needed. |
| Hop | Powdery Mildew <i>Sphaerotheca macularis</i> Downy Mildew <i>Peronospora spp.</i> | 2 - 4 lbs./100 gal | Use the higher rates when moderate to high disease pressure is present or expected. Begin applications when environmental conditions are conducive to rapid disease development. Continue sprays at 7-day intervals or as needed. Apply at a rate of 2-4 lbs. per 100 gallons of water using ground equipment. Minimum spray volume recommendations for hop growth stages are as follows: Emergence to training: Use 2-4 lbs. of product per 100 gallons of water. Apply using a minimum spray volume of 20 gallons per acre. Coverage will vary with the size of the vines and the type of spray equipment. Apply adequate spray volume to achieve complete spray coverage. Training to wire: Use 2-4 lbs. of product per 100 gallons of water. Apply using a minimum spray volume of 50 gallons per acre. Coverage will vary with the size of the vines and the type of spray equipment. Apply adequate spray volume to achieve complete spray coverage. Wire touch through harvest: Use 2-4 lbs. of product per acre. Apply in a minimum spray volume of 100 gallons per acre. Higher water volumes may be necessary to achieve thorough coverage after side arms develop. Apply adequate spray volume to achieve complete spray coverage. |

Rate presented in lbs./acre unless otherwise noted.

| Crops | Disease | Rate lbs./acre# | Application Instructions |
|-------|---------|--------------------|--------------------------|
|-------|---------|--------------------|--------------------------|

| | | | |
|---|--|---------------------------------------|--|
| <p>Leafy Vegetables Lettuce Celery Spinach Parsley Radicchio and other leafy vegetable crops</p> | <p>Downy Mildew <i>Bremia lactucae</i> <i>Peronospora</i> spp. Powdery Mildew <i>Erysiphe cichoracearum</i> Pink Rot <i>Sclerotinia sclerotiorum</i></p> | <p>1 - 3</p> | <p>Pink rot – Begin application approximately 8 weeks before harvest and repeat on a <u>14-day</u> interval. Apply Serenade as a directed spray in sufficient water to ensure thorough coverage of the base of the plants and the surrounding soil surface. Light irrigation following application to incorporate Serenade may improve disease control.</p> <p>Downy mildew / powdery mildew - For suppression, begin application when conditions are conducive to disease development and repeat on <u>7 to 10</u> day intervals or as needed. For improved performance, use Serenade MAX in a tank mix or rotational program with other registered fungicides for downy mildew and powdery mildew control.</p> |
| <p>Leafy Vegetables Lettuce Celery Spinach Parsley Radicchio and other leafy vegetable crops</p> | <p>Sclerotinia Head and Leaf Drop <i>Sclerotinia</i> spp.</p> | <p>1-3</p> | <p><u>For control of early Sclerotinia head and leaf drop:</u> Apply at planting or immediately following planting but prior to crop emergence as a 4 to 6 inch seed line treatment. Make a second application as a directed spray with multiple nozzles per each seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on <u>10 to 14</u> day intervals if conditions for disease development persist. Use higher rates under conditions conducive to moderate to severe disease pressure. Light irrigation after application to incorporate the product may improve disease control.</p> <p style="text-align: center;">OR</p> <p><u>For control of Sclerotinia head and leaf drop:</u> Apply as a directed spray with multiple nozzles per each seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning or transplanting. Repeat applications on <u>10 to 14</u> day intervals if conditions for disease development persist. Use higher rates under conditions conducive to moderate to severe disease pressure. Light irrigation after application to incorporate the product may improve disease control.</p> |
| <p>Legumes/ Vegetables (succulent and dried beans and peas)</p> | <p>Rust <i>Uromyces appendiculatus</i></p> | <p>1 - 3</p> | <p>For suppression, begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for rust control.</p> |
| <p>Beans Green beans Snap beans Shell beans</p> | <p>Rust <i>Puccinia</i> spp Powdery Mildew <i>Erysiphe</i> spp</p> | <p>1 - 3</p> | <p>Begin applications when environmental conditions and plant stage are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure.</p> |
| <p>Dry Beans Garbanzo beans</p> | <p><i>Aphonomyces</i> spp</p> | <p>1 - 3</p> | <p>Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed.</p> |
| <p>Lima beans Peas Chick peas Split peas Lentils and other legume/ vegetable crops</p> | <p>White Mold <i>Sclerotinia sclerotiorum</i></p> | <p>1 - 3</p> | <p>Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed.</p> |
| <p>Crops</p> | <p>Disease</p> | <p>Rate lbs./acre[#]</p> | <p>Application Instructions</p> |

| | | | |
|---|--|-------|--|
| Mint and other herb/spices | Rust <i>Puccinia menthae</i> Powdery Mildew <i>Erysiphe spp</i> Downy Mildew <i>Peronospora spp.</i> | 1 - 3 | Begin application soon after emergence and when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure. |
| Oil Seed Crops Canola Castor Coconut Cotton Flax Oil Palm Olive Peanut Rapeseed Safflower Sesame Sunflower Soybeans and other oilseed crops | Bacterial Speck <i>Pseudomonas syringae pv. glycinea</i> Brown Spot <i>Septoria glycines</i> Pod and Stem Blight <i>Diaporthe phaseolorum var. sojae</i> <i>Phomopsis longicola</i> Downy Mildew <i>Peronospora mansherica</i> Rust | 1 - 3 | Begin application soon after emergence and when conditions are conducive to disease development. Repeat on <u>7 to 10</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure. |
| Olive | Olive Knot <i>Pseudomonas savastanoi</i> | 1 - 3 | Apply before fall rains and again during dormancy before spring growth. Under conditions conducive to heavy disease pressure for improved control, use Serenade in a tank-mix or rotational program with a copper-based bactericide registered for control of olive knot. |
| Peanut | Early Leaf Spot <i>Cercospera arachidicola</i> Late Leaf Spot <i>Cercosporidium personatum</i> Rust <i>Puccinia arachidis</i> White Mold <i>Sclerotinia sclerotiorum</i> | 1 - 3 | Begin application when environmental conditions are conducive to disease development. Repeat applications on <u>14-day</u> intervals or as needed. For improved control, use Serenade MAX in a tank mix program with copper-based fungicides registered for control of peanut leaf spot. . Peanut hay may be fed to livestock. |

| Crops | Disease | Rate lbs./acre [#] | Application Instructions |
|-------|---------|--------------------------------|--------------------------|
|-------|---------|--------------------------------|--------------------------|

| | | | |
|--|--|---------------------------------------|--|
| Pome Fruit Apple Crabapple Pear Quince Mayhaw and other pome fruit | Fire Blight <i>Erwinia amylovora</i> | 2 - 4 | For suppression begin application at 1 - 5% bloom and repeat as necessary to protect open, untreated blossoms when conditions favoring disease development are likely to occur. For maximum control, use Serenade MAX prior to and as close as possible to fire blight infection events. During periods of rapid bloom development and frequent infection periods, spray intervals of <u>3 - 7</u> days may be required. After petal fall, continue applications on a <u>7-day</u> interval while environmental conditions favor disease development. Apply in sufficient water to provide full coverage. For improved performance, use Serenade MAX in a rotational program with antibiotics registered for fire blight control such as but not limited to oxytetracycline or streptomycin. Proper orchard cultural practices are essential to eliminate fire blight-infected tissue from the orchard to assure good performance of any crop protection product. Care must be taken to remove and destroy dead and diseased wood from the orchard prior to and during the growing season. Use of Serenade Max alone has not been shown to affect fruit finish. Use caution when selecting spray adjuvants. Select only those adjuvants which through prior experience do not affect fruit finish when combined with Serenade MAX. |
| | Scab <i>Venturia spp.</i> | 1 - 3 | For suppression, begin application at green tip or when environmental conditions become favorable for primary scab development and repeat on a <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for scab control. |
| | Powdery Mildew <i>Podosphaera leucotricha</i> | 1 - 3 | Begin application at tight cluster, or sooner, if conditions are conducive to disease development. Repeat applications through the second cover spray on a <u>7 to 10</u> day interval. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. |
| Root / Tuber and Corm Vegetables | Black Root Rot/ Black Crown Rot <i>Alternaria spp.</i> | 1 - 3 | Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed. |
| Carrot Potato Sweet Potato Beets Ginger Horseradish Radish Gingseng Turnip and other root/ tuber and corm crops | Bacterial Leaf Blight <i>Xanthomonas campestris</i> Downy Mildew <i>Peronospora spp.</i> Powdery Mildew <i>Erysiphe spp.</i> White Mold <i>Sclerotinia sclerotiorum</i> Gray Mold <i>Botrytis spp.</i> | 1 - 3 | Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed. |
| | Early Blight <i>Alternaria solani</i> Late Blight <i>Phytophthora infestans</i> | 1 - 3 | For suppression, begin application soon after emergence and when conditions are conducive to disease development. Repeat on a <u>5 to 7</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for late blight control. |
| Roses, Field | Powdery Mildew <i>Sphaerotheca spp.</i> Rust <i>Puccinia spp..</i> | 2 - 4 | Begin applications when environmental conditions and plant stage are conducive to disease development. Continue applications on <u>7 to 14</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure. |
| Crops | Disease | Rate lbs./acre[#] | Application Instructions |

| | | | |
|--|---|--|--|
| <p>Stone Fruit Apricot Cherry Nectarine Peach Plum Prune and other stone fruit crops</p> | <p>Anthracnose <i>Colletotrichum</i> spp. Powdery Mildew <i>Sphaerotheca parnosa</i> <i>Podosphaera clandestine</i> <i>Podosphaera</i> spp Rusty Spot <i>Podosphaera leucotricha</i> Bacterial Canker <i>Pseudomonas</i> spp. Alternaria Spot / Fruit Rot <i>Alternaria alternata</i> Scab <i>Cladosporium carpophilum</i> Brown Rot Blossom Blight <i>Monillinia laxa</i> Fruit Brown Rot <i>Monillinia fruticola</i> Gray mold <i>Botrytis cinerea</i> Shot Hole <i>Wilsonomyces carpophilus</i> <i>Xanthomonas pruni</i> <i>Bhumeriella gaapi</i> <i>Cercospora</i> spp.</p> | <p>1 - 3</p> | <p>Brown Rot Blossom Blight – Begin application at early bloom and repeat through petal fall on a <u>7-day</u> interval or as needed.</p> <p>Scab – Begin application at petal fall and repeat on a <u>7 to 10</u> day interval or as needed.</p> <p>Bacterial Blight – Apply post harvest before fall rains and again during dormancy before spring growth.</p> <p>Powdery Mildew - For suppression, begin application at popcorn stage and repeat on a <u>7-day</u> interval or as needed. For improved performance, use Serenade MAX in a tank mix or rotational program with other registered fungicides for powdery mildew control.</p> <p>For all other diseases – Begin application prior to disease development when environmental conditions and plant stage are conducive to rapid disease development and repeat on a <u>7 to 10</u> day interval or as needed.</p> <p>Post harvest disease protection – To aid in the control of post harvest infections of <i>Botrytis</i> and <i>Monolinia</i> apply Serenade prior to harvest with sufficient water to thoroughly cover fruit. Apply on a 7-day schedule or as needed up until the time of harvest.</p> <p>Serenade MAX may be applied to fruit up to and including the day of harvest.</p> |
| <p>Strawberry</p> | <p>Powdery Mildew <i>Sphaerotheca macularis</i> Erysiphe spp. Anthracnose <i>Colletotrichum acutatum</i> Botrytis <i>Botrytis cinerea</i> Gray Mold <i>Botrytis</i> spp.</p> | <p>1 - 3</p> | <p>Botrytis/Powdery mildew - For suppression, begin application at or before flowering and repeat on <u>7 to 10</u> day intervals or as needed through harvest. Use higher rates and shorter application intervals under heavy disease pressure. For improved performance, use Serenade MAX in a tank mix or rotational program with other registered fungicides for powdery mildew and botrytis control.</p> <p>Anthracnose – Begin application prior to disease development and repeat on <u>7 to 10</u> day intervals or as needed.</p> <p>Serenade may be applied up to and including the day of harvest.</p> |
| <p>Sugar Beets</p> | <p>Powdery Mildew <i>Erysiphe betae</i> <i>Erysiphe polygoni</i> Leaf Spot <i>Cercospora beticola</i> Ramularia <i>Ramularia</i> spp Rust <i>Uromyces betae</i></p> | <p>1 - 3</p> | <p>Begin applications when environmental conditions are conducive to disease development. Continue applications on <u>7 to 10</u> day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure.</p> |
| <p>Tobacco</p> | <p>Blue Mold <i>Peronospora hyoscyami</i></p> | <p>1 - 3</p> | <p>Begin applications when conditions are conducive to disease development. Continue applications on <u>7 to 10</u> day intervals or as needed.</p> |
| <p>Crops</p> | <p>Disease</p> | <p>Rate lbs./acre[#]</p> | <p>Application Instructions</p> |

| | | | |
|--|--|--------------|---|
| <p>Tree Nuts Almond Pistachio Pecan Walnut Filberts Chestnut Cashew Beechnut Butternut and other treenut crops</p> | <p>Walnut Blight <i>Xanthomonas campestris</i> Alternaria Leaf Spot <i>Alternaria alternata</i> Anthracnose <i>Colletotrichum acutatum</i> Bacterial Canker <i>Pseudomonas syringae</i> Scab <i>Cladosporium carpophilum</i> Botryosphaeria Blight <i>Botryosphaeria dothidea</i> Shot Hole <i>Wilsonomyces carpophilus</i> <i>Xanthomonas pruni</i> <i>Blumeriella gaapi</i> <i>Cercospora</i> spp. Brown Rot <i>Monilinia</i> spp.</p> | <p>2 - 4</p> | <p>Walnut Blight – Begin application no later than pistillate bloom and repeat on 7 to 10 day intervals or as needed. Apply in advance of rain for maximum protection. Under conditions conducive to heavy disease pressure, for improved control, use Serenade in a tank-mix or rotational program with a copper-based bactericide registered for control of walnut blight.</p> <p>For all other diseases – Begin application prior to disease development and repeat on 7 to 10 day intervals or as needed.</p> |
| <p>Tropical Fruits Avocado Mango Papaya and other tropical fruits</p> | <p>Anthracnose <i>Colletotrichum gloeosporioides</i> Scab <i>Sphaceloma perseae</i></p> | <p>1 - 3</p> | <p>Avacado/Mango - Begin application at budbreak and repeat on a 14 to 21 day interval or as needed through harvest.</p> <p>Papaya - Begin application at flowering and repeat on a 14 to 21 day interval or as needed through harvest.</p> <p>Serenade MAX may be applied to fruit up to and including the day of harvest.</p> |
| <p>Kiwi</p> | <p>Botrytis Fruit Rot <i>Botrytis cinerea</i> Bacterial Blight <i>Pseudomonas viridiflava</i> and <i>Pseudomonas syringae</i> Sclerotinia <i>Sclerotinia sclerotiorum</i></p> | <p>2 - 4</p> | <p>Kiwi - Begin application at early bloom and repeat on 7 to 10 day intervals or as needed. Serenade MAX may be applied to fruit up to and including the day of harvest.</p> |
| <p>Watercress</p> | <p><i>Cercospora</i> leafspot <i>Cercospora</i> spp.</p> | <p>1 - 3</p> | <p>Begin applications when conditions are conducive to disease development. Continue applications on 7 to 10 day intervals or as needed.</p> |
| <p>Seed Production Crops blue grass rye grass fescue orchard grass and other crops grown to produce seeds</p> | <p>Powdery Mildew <i>Erysiphe</i> spp. Rust <i>Puccinia</i> spp.</p> | <p>1 - 3</p> | <p>Begin applications when environmental conditions and plant stage are conducive to disease development. Repeat on 7 to 10 day intervals or as needed. Use higher rates and shorter application intervals under heavy disease pressure.</p> |

Rate presented in lbs./acre unless otherwise noted.

Recommended Application Rates for Selected Greenhouse Crops
(Serenade has a 0-Day PreHarvest Interval for all crops contained on this label)

Under moderate to severe disease pressure, for improved performance, increase rates and reduce spray intervals or use Serenade MAX in a tank mix or rotational program with other registered fungicides.

| Green- | Diseases | Rate | Application Instructions |
|--------|----------|------|--------------------------|
|--------|----------|------|--------------------------|

| house Crops | | lbs./100 gallons spray mix | |
|--|---|---|---|
| Brassica Broccoli Cabbage Cauliflower Brussels Sprouts Collards Kale Mustard Greens Kohlrabi and other brassica crops | Pin Rot Complex <i>Alternaria/Xanthomonas</i> Bacterial Leaf Spot <i>Pseudomonas syringae</i> Bacterial Soft Rot <i>Erwinia / Pseudomonas</i> Black Rot <i>Xanthomonas</i> <i>campestris</i> Xanthomonas Leaf Spot <i>Xanthomonas</i> <i>campestris</i> Alternaria Leaf Spot <i>Alternaria</i> spp. Anthracnose <i>Colletotrichum</i> <i>higginsianum</i> Cercospora Leaf Spot <i>Cercospora brassicaicola</i> Downy Mildew <i>Peronospora parasitica</i> <i>Peronospora</i> spp. Powdery Mildew <i>Erysiphe polygoni</i> Southern Blight <i>Sclerotium rolfsii</i> | 1- 3 | <p>Pin Rot - For suppression, begin application when environmental conditions in the greenhouse are conducive to disease development and repeat on a <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for pin rot control.</p> <p>For all other diseases - Begin application soon after emergence or transplant and when conditions in the greenhouse are conducive to disease development. Repeat on a 7- to 10-day interval or as needed.</p> |
| Bulb Vegetables Onion Garlic Shallots and other bulb vegetables | Botrytis neck rot <i>Botrytis</i> spp. Botrytis Leaf Blight <i>Botrytis squamosa</i> Onion Purple Blotch <i>Alternaria porri</i> Onion Downy Mildew <i>Peronospora destructor</i> Downy Mildew <i>Peronospora</i> spp. Powdery Mildew <i>Erysiphe</i> spp. | 1- 3 | <p>Begin application when environmental conditions in the greenhouse are conducive to disease development and repeat on a 7- to 10-day interval or as needed.</p> |
| | Rust <i>Puccinia porri</i> | 1 - 3 | <p>For suppression, begin application when conditions are conducive to disease development and repeat on a 7- to 10-day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for rust control. Do not apply more than 6 lb. per acre per application.</p> |
| Green- house Crops | Diseases | Rate lbs./100 gallons spray mix | Application Instructions |

| | | | |
|---|---|------|---|
| Cucurbits Cucumber Cantaloupe Melon Muskmelon Squash Watermelon and other cucurbits | Powdery Mildew Erysiphe spp. Sphaerotheca spp. Gummy Stem Blight <i>Phoma cucurbitacearum</i> <i>Didymella bryoniae</i> Angular Leaf Spot <i>Pseudomonas syringae</i> Anthracnose <i>Colletotrichum lagenarium</i> Downy Mildew <i>Pseudoperonospora cubensis</i> Bacterial Fruit Blotch <i>Acidovorax avenaesubsp</i> | 1 -3 | Begin application soon after emergence or transplant and when environmental conditions in the greenhouse are conducive to disease development. Repeat on <u>7 to 10</u> day interval or as needed. |
| Fruiting Vegetables Pepper Tomato Eggplant and other fruiting vegetables | Gray mold <i>Botrytis cinerea</i> | 1 -3 | Begin application soon after emergence or transplant and repeat on a <u>7 to 10</u> day interval or as needed |
| | Powdery mildew <i>Leveillula taurica</i> <i>Oidiopsis taurica</i> Downy Mildew <i>Pseudoperonospora cubensis</i> | 1 -3 | For suppression, begin application soon after emergence or transplant and continue on a <u>7 to 10</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for powdery mildew control |
| | Bacterial Spot <i>Xanthomonas</i> spp. | 1 -3 | Begin application soon after emergence or transplant and when environmental conditions are conducive to disease development. Continue applications on a <u>5 to 7</u> day interval or as needed. When conditions are conducive to rapid disease development, for improved control, use Serenade in a tank mix program with copper-based bactericides registered for control of bacterial spot at labeled rates |
| | Early Blight <i>Alternaria solani</i> Late Blight <i>Phytophthora infestans</i> | 1 -3 | For suppression, begin application when plants are 4- to 6-inches high. Repeat applications on a 5- to 7-day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for late blight control. |
| Herbs/ Spices | Bacterial Blight <i>Pseudomonas syringae</i> Anthracnose <i>Colletotrichum</i> spp. Alternaria Leaf Blight <i>Alternaria</i> spp. | 1 -3 | Begin application when environmental conditions in the greenhouse are conducive to disease development. Repeat on a <u>7- to 10-day</u> interval or as needed. |
| Leafy Vegetables Lettuce Celery Spinach Parsley Radicchio and other leafy vegetables | Downy Mildew <i>Bremia lactucae</i> <i>Peronospora</i> spp. Powdery Mildew <i>Erysiphe cichoracearum</i> <i>Erysiphe</i> spp. Pink Rot <i>Sclerotinia sclerotiorum</i> | 1 -3 | Pink rot – Begin application approximately 8 weeks before harvest and repeat on a 14-day interval. Apply Serenade as a directed spray in sufficient water to ensure thorough coverage of the base of the plants and the surrounding soil surface. Light irrigation following application to incorporate Serenade may improve disease control. Downy mildew / powdery mildew - For suppression, begin application when conditions are conducive to disease development and repeat on a 7- to 10-day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for downy mildew and powdery mildew control. |

18 7 22

| | | | |
|---|---|-------|---|
| | Sclerotinia Head and Leaf Drop <i>Sclerotinia</i> spp. | 1 - 3 | For control of early Sclerotinia head and leaf drop: Apply at planting or immediately following planting but prior to crop emergence as a 4 to 6 inch seed line treatment. Make a second application as a directed spray with multiple nozzles per each seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning. Repeat applications on <u>7 to 10</u> day intervals if conditions for disease development persist. Use higher rates under conditions conducive to moderate to severe disease pressure. Light irrigation after application to incorporate the product may improve disease control. OR For control of Sclerotinia head and leaf drop: Apply as a directed spray with multiple nozzles per each seed line in sufficient water to ensure thorough coverage of lower plant leaves and surrounding soil surface within 7 days of thinning or transplanting. Repeat applications on <u>7 to 10</u> day intervals if conditions for disease development persist. Use higher rates under conditions conducive to moderate to severe disease pressure. Light irrigation after application to incorporate the product may improve disease control. |
| Root / Tuber Carrot Potato Sweet Potato Beets Ginger Horseradish Radish Gingseng Turnip and other root/tuber crops | Black Root Rot/ Black Crown Rot <i>Alternaria</i> spp. | 1 - 3 | Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed. |
| | Bacterial Leaf Blight <i>Xanthomonas campestris</i> White Mold <i>Sclerotinia sclerotiorum</i> | 1 - 3 | Begin application soon after emergence or transplant and when conditions are conducive to disease development. Repeat on a <u>7 to 10</u> day interval or as needed. |
| | Early Blight <i>Alternaria solani</i> Late Blight <i>Phytophthora infestans</i> | 1 - 3 | For suppression, begin application soon after emergence and when conditions are conducive to disease development. Repeat on a <u>5 to 7</u> day interval or as needed. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for late blight control. |
| Strawberry | Powdery Mildew <i>Sphaerotheca macularis</i> <i>Erysiphe</i> spp Anthracnose <i>Colletotrichum acutatum</i> Botrytis <i>Botrytis cinerea</i> Gray Mold <i>Botrytis</i> spp. | 1-3 | Botrytis/Powdery mildew - For suppression, begin application at or before flowering and repeat on a <u>7 to 10</u> day interval or as needed through harvest. For improved performance, use Serenade in a tank mix or rotational program with other registered fungicides for powdery mildew and botrytis control. Anthracnose – Begin application prior to disease development and repeat on a <u>7 to 10</u> day interval or as needed. Serenade may be applied up to and including the day of harvest. |

POST HARVEST DISEASE PROTECTION

| Crops | Diseases | Rate oz/Ton | Application Instructions |
|-------------------------------------|--|--------------------|---|
| Potatoes (post harvest application) | Silver Scurf <i>Helminthosporium solani</i> | 1.0-3.0 oz/ Ton | For the post harvest application to aid in the control of silver scurf. Sanitation and other cultural practices should also be employed to aid in control and minimize the potential for disease. Prepare the equivalent of 3 1/4 to 9 3/4 lbs of Serenade MAX in 25 gallons of water. Spray 2 quarts of the suspension per ton of potatoes. Potatoes must rotate along the conveyor line into the storage area to ensure complete coverage. |

For Home and Garden Use

Serenade® MAX™

[Alternate Name: Serenade® Garden Disease Control]

[Alternate Name: Serenade® Garden Disease Control Wettable Powder]

A Wettable Powder Biofungicide

[Optional/Alternate Statement: "NOP Logo: For Organic Production"]

[Optional/Alternate Statement: "NOP Logo: Can Be Used for Organic Production"]

ACTIVE INGREDIENT

QST 713 strain of dried *Bacillus subtilis*14.6%

OTHER INGREDIENTS 85.4%

Total 100.0%

Contains a minimum of 7.3 x 10⁹ cfu/g

EPA Reg. No. 69592-X

EPA Est. No.:

| | | | | |
|--------|--------|--------|--------|--------|
| 1 | 2 | 3 | 4 | 5 |
| 69592- | 67545- | 66728- | 37429- | 69592- |
| MEX-1 | AZ-1 | GA-2 | GA-2 | CA-1 |

Superscript corresponds to last digit of lot number stamped on container

U.S. Patent Nos. 6,060,051, 6,103,228, 6,291,426, and 6,417,163 on QST 713 strain of *Bacillus subtilis*

Net weight:

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS— Home and Garden

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove contaminated clothing and wash clothing before use.

ENVIRONMENTAL HAZARDS – Home and Garden

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE – Home and Garden

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

GENERAL USE INFORMATION – Home and Garden

Serenade MAX is a broad spectrum, preventative biofungicide recommended for the control or suppression of many important plant diseases. For control or suppression, use Serenade MAX as a preventative treatment when seedlings first emerge, and alternate treatment of plants with other appropriately labeled registered fungicides or bactericides.

This product can be used for organic and non-organic crop production.

As a general precaution, when exposed to high concentrations of a living microbial product such as this, wear a dust particle mask when mixing or applying this product.

MIXING AND APPLICATION INSTRUCTIONS – Home and Garden

MIXING:

Mix 1 /8 to 1 /2 cups of Serenade Max per gallon of water. Mix thoroughly and keep spray solution agitated during application. Use higher rates when environmental conditions are conducive to rapid disease development (high humidity, excessive rain, extreme moisture conditions).

Application:

Use 15 fluid ounces of spray solution per 100 square feet. Spray plant to runoff, covering both top and bottom surface of foliage, to insure thorough coverage. For best results, treat prior to disease development or at the first sign of disease infection. Repeat at 7-day intervals or as needed. Serenade MAX can be applied up to and including the day of harvest.

PLANTS [CROPS, SITES]:

HOME and GARDEN PLANTS:

Artichoke, Asparagus,

Berries (Blueberries, Blackberry, Raspberry, Loganberry, Huckleberry, Cranberry, Gooseberry, Elderberry, Currant, Caneberry, and other berry crops)

Brassica (Broccoli, Cabbage, Cauliflower, Brussels Sprouts, Collards, Kale, Mustard Greens, Kohlrabi and other brassica crops)

Bulb Vegetables (Onion, Garlic, Shallots and other bulb vegetables)

Citrus (Orange, Grapefruit, Lemon, Tangerine, Tangelo, Pummelo and other citrus crops)

Cucurbits (Cucumber, Cantaloupe, Melon, Muskmelon, Squash, Watermelon and other cucurbit crops)

Fruiting Vegetables (Pepper, Tomato, Eggplant and other fruiting vegetables)

Grape, Herbs/ Spices, Hop,

Leafy Vegetables (Lettuce, Celery, Spinach, Parsley, Radicchio and other leafy vegetable crops)

Legumes/vegetables (Beans, Green beans, Snap beans, Shell beans, Dry Beans, Garbanzo beans, Lima beans, Peas, Chick peas, Split peas, Lentils and other legume/ vegetable crops)

Mango, Mint, Olive, Papaya, Peanuts,

Pome Fruit (Apple, Crabapple, Pear, Quince, Mayhaw and other pome fruit)

Root / Tuber (Carrot, Potato, Sweet Potato, Beets, Ginger, Horseradish, Radish, Gingseng, Turnip and other root/ tuber crops)

Roses

Stone Fruit (Apricot, Cherry, Nectarine, Peach, Plum, Prune, and other stone fruit crops)

Strawberry, Sweet Corn, Tobacco, Watercress,

Tree Nut (Almond, Pistachio, Pecan, Walnut, Filberts, Chestnut, Cashew, Beechnut, Butternut and other tree nut crops)

GREENHOUSE PLANTS:

Brassica (Broccoli, Cabbage, Cauliflower, Brussels Sprouts, Collards, Kale, Mustard Greens, Kohlrabi and other brassica crops)

Bulb Vegetables (Onion, Garlic, Shallots and other bulb vegetables)

Cucurbits (Cucumber, Cantaloupe, Melon, Muskmelon, Squash, Watermelon and other cucurbits)

Fruiting Vegetables (Pepper, Tomato, Eggplant and other fruiting vegetables)

Herbs/ Spices

Leafy Vegetables (Lettuce, Celery, Spinach, Parsley, Radicchio, and other leafy vegetables)

Root / Tuber (Carrot, Potato, Sweet Potato, Beets, Ginger, Horseradish, Radish, Gingseng, Turnip and other root/ tuber crops)

Strawberry

DISEASES CONTROLLED OR SUPPRESSED

Bacterial Leaf Blight (*Xanthomonas campestris*)

Bacterial Spot (*Xanthomonas* spp.) - suppression

Bean Rust (*Uromyces appendiculatus*) - suppression

Black Root Rot/Black Crown Rot (*Alternaria* spp.)

Botrytis Leaf Blight (*Botrytis squamosa*)

Botrytis Neck Rot (*Botrytis* spp.)

Downy Mildew (*Bremia lactucae*, *Peronospora* spp., and *Plasmopara viticola*) - suppression

Early Blight (*Alternaria solani*) - suppression

Fire Blight (*Erwinia amylovora*) - suppression

Gray Mold (*Botrytis cinerea*)

Greasy spot (*Mycosphaerella citri*) - suppression

Late Blight (*Phytophthora infestans*) - suppression

Onion Downy Mildew (*Peronospora destructor*)

Onion Purple Blotch (*Alternaria porri*)

Pin Rot (*Alternaria/Xanthomonas* complex) - suppression

Powdery Mildew (*Uncinula necator*, *Erysiphe* spp., *Sphaerotheca* spp., *Oidiopsis taurica*, *Leveillula taurica*, *Podosphaera leucotricha*)

22 8 22

Scab (*Venturia* spp.) - suppression

Sclerotinia head and leaf drop (*Sclerotinia* spp.)

Sour Rot

Walnut Blight (*Xanthomonas campestris*)

White Mold (*Sclerotinia sclerotiorum*) – suppression

ORNAMENTALS, TREES, SHRUBS, FLOWERING PLANTS, TROPICAL PLANTS

DISEASES CONTROLLED OR SUPPRESSED

Anthracnose -- *Colletotrichum* spp.

Bacteria – *Erwinia* spp, *Pseudomonas* spp, *Xanthomonas* spp

Black spot of rose – *Diplocarpon rosea*

Botrytis -- *Botrytis cinerea*

Downy Mildew -- *Peronospora* spp.

Leaf spots – *Alternaria* spp, *Cercospora* spp, *Entomosporium* spp, *Helminthosporium* spp, *Myrothecium* spp, *Septoria* spp.

Powdery mildew – *Erysiphe* spp, *Oidium* spp, *Podosphaera* spp, *Sphaerotheca* spp.

Phytophthora spp --

Rust – *Puccinia* spp.

Scab –
Venturia spp.

STORAGE AND DISPOSAL – Home and Garden

STORAGE: Store in a dry area inaccessible to children. Store in original containers only. Keep container closed when not in use.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS FOR SALE AND WARRANTY

AgraQuest warrants to those persons lawfully purchasing this product that at the time of the first sale of this product by Seller that this product conformed to its description and was reasonably fit for the purposes stated on the label when used in accordance with Seller's directions. Buyers and users of this product assume the risk of any use contrary to such directions. EXCEPT AS PROVIDED ELSEWHERE IN WRITING CONTAINING AN EXPRESS REFERENCE TO THIS WARRANTY AND LIMITATION OF DAMAGES, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OR GUARANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO. To the extent permitted by state law, the Seller's liability for any breach of warranty shall not exceed the purchase price of the material as to which a claim is made.

To the extent permitted by state law, Buyers and users of this product are responsible for all loss or damage from use or handling of this product which results from conditions beyond the control of Seller, or without the fault or negligence of the Seller, or from failure to follow the label.

Serenade MAX is a registered trademark of AgraQuest, Inc.

© Copyright AgraQuest, Inc., 2003