12/17/200

OFFICE OF PREVENTION, PESTICIDES AND

TOXIC SUBSTANCES

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

December 17, 2009

Christine A. Dively Director of Regulatory Affairs Certis USA, LLC 9145 Guilford Road, Suite 175 Columbia, MD 21046

> RE: Product Name: Neem Oil 70% EPA Reg. No: 70051-2 Application for Label Notification Dated October 22, 2009 to update storage and disposal statements and add 250g refillable containers to sub label A per PR Notice 2007-4.

Dear Ms.Dively:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under Pesticide Registration (PR) Notice 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PR Notice 98-10 and it has been determined that the action(s) requested falls within the scope of PR Notice 98-10. Our records have been duly noted, and the label submitted with this application has been stamped "Notification received and reviewed" and will be placed accordingly in our records.

If you have any questions concerning this action, please feel free to contact Ms. Menyon Adams at (703) 347-8496 or email at <u>adams.menyon@epa.gov.</u>

Sincerely,

Linda Hollis

Linda Hollis, Chief Biochemical Pesticides Branch Biopesticides and Pollution Prevention Division (7511P)

Please read Instructions o	n reverse before con.,	ting form.	Form Apr	noved. Sum No. 2	070-0060	Print Form
\$EPA	Environmental	Inited States Protection Ag ngton, DC 20460	jency	Registr Amend X Other		OPP Identifier Number
	······································	Application for	Pesticide - Sec	ction I		
. Company/Product Num 70051-2	ber		2. EPA Product Ma Biochemical Pest		3. Pro	posed Classification
4. Company/Product (Nan Neem Oil 70%	1e)		PM# 91		\neg	None Restricted
Certis U.S.A., L.L.C. 9145 Guilford Road, S Columbia, MD 21046	Applicant <i>(Include ZIP Co</i> uite 175 <i>his is a new address</i>			is similar or iden	tical in cor	FIFRA Section 3(c)(3) nposition and labeling
		Se	ction - II			CATION
Explanation: Use eddi Notification o the requireme been made to U.S.C. Sec. 10 with the requi		y. (For section I and tice 2007-4. This notif at 40 CFR §§ 156.10, 1 lential Statement of Fc false statement to EP, 6.10, 156.140, 156.14 ment action and penal	Section II.) Teation is consistent with 156.140, 156.144, 156.1 rmula for this product. A. I further understand to 4, 156.146, and 156.156	h the guidance in Pl 46, and 156.156. No I understand that it that if the amended 5, this product may b	R Notice 200 o other chan is a violatior abel is not c	7: D7-4 and ges have to of 18 consistent
1. Material This Product	Will Be Packaged In:		·····			
Child-Resistent Peckeging Yes* No * Certification must be submitted	Yes X No	No. per container Pac	Yes" No. per kage wgt contain	er	f Container Metal Plastic Glass Paper Other (S	
3. Location of Net Conter	ts Information	4. Size(s) Roteil Gon 8 16 20 24	12 . , 21/2 250 cal	5. Location of La X On Lab On Lab	e)	ns ນັ້ ວ່ານວ່າ panying product
6. Manner in Which Labe	•	Lithograph Paper glued Stenciled	Oth		<u> </u>	<u>- ມ</u> ງາງ ບ
		and the second data was not a second data was not second as a second data was not second as a second data was n	ction - IV		د د د ر	
1. Contact Point (Compl	ete items directly below			d, if necessary, to p	progess shis	application.b
Name Christine Dively		Title	tor, Reg. Affairs		Telephoni	a No. (Include Area Code) -3806
-		misleading statement	may be punishable by I	ine of imprisonmen	t or	6. Date Application Received (Stamped)
2 Signature			1 2 17	1 - 1 - 1 - 1 - 1		
2 Signature	ino A. Dui	ely_ Dir	ector of Regu October 2	latory Aff	air	

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

.

/



Certis USA 9145 Guilford Road Suite 175 Columbia, MD 21046

(301) 604-7340

FAX: 301-604-7015 www.certisusa.com

October 26, 2009

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency 1200 Pennsylvania Ave., NW Washington DC 20460-0001

Re: Notification of label change per PR Notice 2007-4

Certis USA, L.L.C. respectfully submits the following label change for Neem Oil 70% (EPA Registration Number 70051-2) to include language recommended in PR Notice 2007-4 that was erroneously omitted from the original submission.

Neem Oil 70% is offered in 250-gallon refillable containers in addition to the nonrefillable containers previously identified. Container reuse statements and cleaning instructions appropriate for these refillable containers were added to the Storage and Disposal section of Sublabel A. No other changes were made to the label.

Enclosed in this submission are the following:

- Application for Registration EPA Form 8570-1
- A copy of draft labeling with the changes highlighted

Should you have any questions about this submission, I can be contacted by telephone at 301-483-3806 or by email at <u>cdively@certisusa.com</u>.

Respectfully,

Christin A. Dively -

Christine A. Dively Director of Regulatory Affairs

MASTER LABEL

This master label bears directions for agricultural, commercial and homeowner use with various application rates specific to the associated container sizes

Neem Oil 70% Fungicide/Miticide/Insecticide

Sublabel A: Agricultural Uses Sublabel B: Commercial Greenhouse, Nurseries, Ornamentals Sublabel C: Homeowner Use

Alternate Brand Names: Trilogy, Triact 70EC, Trilogy PHC, Fertilome Rose, Flower & Vegetable Spray, Ferti-lome Triple Action Plus, Garden Safe Fungicide 3 Concentrate, UltraStop Fruit Tree 3-in-1 Concentrate, Concern From the Earth For the Earth Garden Defense Multi-Purpose Spray Concentrate

Active Ingredient:

Clarified Hydrophobic Extract of Neem Oil	70.00%
Other Ingredients	30.00%
Total	100.00%

EPA Registration Number: 70051-2 EPA Establishment Number: 70051-CA-001

Manufactured by:

Certis U.S.A., L.L.C. 9145 Guilford Road, Suite 175 Columbia, Maryland 21046 Master Label / Sublabel A - Commercial AG



For use on vegetables, fruits, nuts, melons, outdoor ornamental plants and agronomic crops. For foliar control of fungal disease and insect pests.

Can be Used Up to Day of Harvest

FOR ORGANIC PRODUCTION

ACTIVE INGREDIENT:

Clarified Hydrophobic Extract of Neem Oil	70%
OTHER INGREDIENTS	
TOTAL	

This product contains 5.46 lb. of clarified hydrophobic extract of neem oil per US gallon.

This container will treat up to 0.5 acres when used at the highest application rate (only for containers less than 5 gallons).

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

FIRST AID

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.



Net Contents: EPA Reg. No. 70051-2 EPA Est. No.70051-CA-001 Composition covered by US Patents 5,298,251; 5,356,629; 5,372,817; 5,405,612; and 5,409,708.

Lot No.:

ددررر

Manufactured by Certis USA, L.L.C. 9145 Guilford Road Suite 175 Columbia, MD 21046



Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or Viton.
- Shoes plus socks.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is hazardous to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 4C CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency' assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that invowes⁹ contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinylchloride (PVC) or Viton.
- Shoes plus socks.

GENERAL INFORMATION:

- Thorough coverage is necessary to provide good disease and mite control. Neem Oil 70% prevents fungal attack of plant tissue and contact activity to mite pressure on the plant. Avoid excessive runoff of spray material for maximum control.
- For best results, maintain constant agitation in spray equipment.
- For optimal performance, do not mix with cold water (less than 45°F).
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted materials or crops prior to root establishment.
- Do not apply to known sensitive plant species, such as impatiens flowers, fuchsia flowers, hibiscus flowers, some rose flowers, ornamental olive trees, or some carnation varieties without prior testing.
- When used in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale.
- As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use with care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.

LANDSCAPE AND NURSERY USE INSTRUCTIONS

FUNGICIDE USE .

ORNAMENTAL DISEASE CONTROL

Neem Oil 70% is an effective fungicide for the prevention and control of various fungal diseases including black spot on roses, powdery mildew, downy mildew, anthracnose, rust, leaf spot, botrytis, needle rust, scab, flower, twig, and tip blight, and alternaria.

Landscape and Nursery Ornamentals

Neem Oil 70% will provide broad spectrum control of many fungal diseases affecting ornamental flowering plants, trees, shrubs, and foliage when used at the rate of 1 gallon per 100 gallons of water. Apply Neem Oil 70% at a rate of 1:100 when spraying as a preventative for outdoor applications. Under extreme outdoor conditions, apply Neem Oil 70% at the 1:50 application rate (see table for detailed use rates). This will provide the necessary control under the extreme outdoor conditions. When using Neem Oil 70% as a preventative, make application on a 14-day schedule. If disease is present, use a 7-day application schedule to control the disease and prevent further damage from occurring.

MIXING INSTRUCTIONS

Add Neem Oil 70% to one-half (1/2) full tank under agitation containing water of 45°F or greater before filling to desired level. If water temperature is below 45°F, premix Neem Oil 70% at a 1:1 ratio with tepid water to ensure good emulsification. Then dilute to final volume. When combining with other products, such as wettable powder insecticides or fungicides, add these items first when the tank is approximately 1/3 full. Ensure that there is good agitation while mixing for complete emulsification. Maintain agitation during spray application. Do not use if this oil does not form a uniform, cloudy emulsion.

000000

APPLICATION INSTRUCTIONS

Apply until plant surface is uniformly wet with minimal runoff at 25-40 psi with hand sprayer or 100-200 psi with power sprayer as a fine spray to both leaf surfaces. Avoid excessive application.

SPRAY TARGET	CONCENTRATION	SPRAY INTERVAL	PRECAUTIONS
Outdoor Flowering Plants	1:50 - 1:100 (2.0% - 1.0%)	7-14 days	Do not spray impatiens flowers.
Outdoor Woody Plants	1:50 - 1:100 (2.0% - 1.0%)	14 days	



SPRAY TARGET	CONCENTRATION	SPRAY INTERVAL	PRECAUTIONS
Snow Mold	1:100-1:500 (1.0%-2.0%)	14 days	In Fall before first snow
Dollar Spot	1:100-1:200 (1.0%-0.5%)	7-14 days	At first sign of disease
Brown Patch	1:100-1:200 (1.0%- 0.5%)	7-14 days	At first sign of disease

INSECTICIDE/MITICIDE USE

For use to control whiteflies, mealybugs, aphids, leafhoppers, mites and scales on ornamentals, trees, and shrubs in and around commercial nurseries and other commercial structures.

Use Neem Oil 70% at the rate of 1-2 gallons per 100 gallons of water. The low rate (1.0%) can be used when pest pressure is low. Use the high rate (2.0%) for high pest infestations or to control more troublesome pests, such as mites and scales. Determine the best rate for your particular situation.

Make sprays on a 7-14 day interval depending on the severity of the pest problem. Use the 7-day schedule until the pest population is reduced. Then use a 14-day interval for control.

AGRICULTURAL USE DIRECTIONS

Neem Oil 70% is a broad spectrum fungicide of certain diseases and controls mites in citrus, deciduous fruits and nuts, vegetable crops, cereal grains, and other miscellaneous crops.

- Do not use Neem Oil 70% after bloom on table grapes or following bunch closure on wine grapes. Do not use Neem Oil 70% after pit hardening on stone fruit.
- Use care when applying near streams, ponds, lakes, or other bodies of water. Do not apply Neem Oil 70% when weather conditions favor drift or when the likelihood of runoff is high.
- Apply Neem Oil 70% by conventional, low volume, and aerial application equipment.

APPLICATION INSTRUCTIONS

Ground Application: Apply Neem Oil 70% by ground application in a minimum of 25 gallons of water / acre.

Chemigation: Do not apply this product through any type of irrigation system.

Aerial Application: For those crops where aerial applications are indicated, apply a minimum of 5 gallons of water / aere unless otherwise specified. Avoid application under conditions when uniform coverage cannot be obtained or when excessive drift may occur. Do not apply directly to humans, animals or aquatic animals.

	(5-30 gallons per acre)	00000 00000	:د د د
Gallons of Water	Neem Oil 70%	DILUTION	0.1
5	1 Quart*	5%	ິວວ
10	1 Quart*	2.5%	
15	1 Quart*	1.67%	00
20	l Quart*	1.25%	
25	1 Quart*	1.0%	
30	1 Quart*	0.83%	

*For citrus, stone fruit, tropical fruit, pome fruit, and nuts, refer to specific application instructions.

0 000000

		(25-250 gallons per acr	· ·	
GallonsWater	Gallons of Neem Oil 70% for 0.5% Solution	Gallons of Neem Oil 70% for 1.0% Solution	Gallons of Neem Oil 70% for 1.5% Solution	Gallons of Neem Oil 70% for 2.0% Solution
25	Use 1.0% solution*	0.25 (32 fl. oz.)	0.375 (48 fl. oz.)	0.5 (64 fl. oz.)
50	0.25 (32 fl. oz.)	0.5 (64 fl. oz.)	0.75 (96 fl. oz.)	1.0
75	0.375 (48 fl. oz.)	0.75 (96 fl. oz.)	1.125	1.5
100	0.5	1.0	1.5	2.0
125	0.625 (80 fl. oz.)	1.25	1.875	2.5
150	0.75	1.5	2.25	3.0
175	0.875 (112 fl. oz.)	1.75	2.625	3.5
200	1.0	2.0	3.0	4.0
225	1.125	2.25	3.375	4.5
250	1.25	2.5	3.75	5.0

*For Vegetable, Cereal Grains, and Other Miscellaneous Crops, refer to specific use directions.

AGRICULTURAL USE DIRECTIONS (continued)

- For Diseases: Apply Neem Oil 70% as a 1.0% solution in sufficient amounts of water to achieve complete coverage of foliage. (See the following detailed directions per crop groupings). When used as a foliar fungicide, begin applications when conditions are favorable for disease but before infection. Apply Neem Oil 70% every 7-14 days until threat of disease is over. Use Neem Oil 70% in rotation with other fungicides to increase the spectrum of control.
- For Mites: For best results, apply Neem Oil 70% in sufficient amounts of water to achieve complete coverage. Neem Oil 70% is most effective when applied before mites or eggs are present in large numbers. (See the following detailed directions per crop groupings). Apply at a rate of 1.0-2.0% Neem Oil 70%. Reapply Neem Oil 70% every 7-21 days until pressure is over.
- For Insects: For best results, apply Neem Oil 70% at 1.0-2.0% in sufficient amounts of water to achieve complete coverage. Neem Oil 70% is most effective when applied before insects or eggs are present in large numbers? Repeat application based on pest pressure and field scouting for best results. Use the dilution tables to calculate dilution rates.

MIXING INSTRUCTIONS

Add Neem Oil 70% to one-half (1/2) full tank under agitation containing water of 45°F or greater before filling to desired level. If water temperature is below 45°F, premix Neem Oil 70% at a 1:1 ratio with tepid water to ensure good emulsification. Then dilute to final volume. When combining with other products, such as wettable powder insecticides or fungicides, add these items first when the tank is approximately 1/3 full. Ensure that there is good agitation while mixing for complete emulsification. Maintain agitation during spray application. Do not use if this oil does not form a uniform, cloudy emulsion.

TANK MIXING: Neem Oil 70% has been found to be compatible with most commonly used fungicides, insecicides and fertilizers. Check physical compatibility first with other products before mixing by using a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. After thoroughly mixing, let mixture stand for at least 5 minutes. Growers must tank mix combinations for phytotoxicity on a sample of plants prior to use. Due to the wide variation in climatic conditions, cultural practices, and other factors, the User assumes full responsibility for any crop damage or other liability resulting from the use of Neem Oil 70% in a tank mix combination. Avoid tank mixes with Morestan, Captan, Sulfur, Bravo, Echo, chlorothalonil or other chemically similar products as unpredictable results or leaf burn may occur.

CITRUS

Use Neem Oil 70% on citrus to control diseases to include Anthracnose, Post-Bloom Fruit Drop, and Scab. Apply as a foliar application at the rate of 1.0% in 25-200 gallons of water for high volume applications or a minimum of 0.5 gallons Neem

Oil 70% in a minimum of 5 gallons of water per acre for low volume applications. Do not apply more than 2.5 gallons of Neem Oil 70% per acre, per application.

Reapply Neem Oil 70% every 7-14 days until threat of disease is over.

For control of Alternaria and Greasy Spot, apply Neem Oil 70% at a 1.0% solution in 25-125 gallons of water per acre in combination with Copper @ 2.0 lbs a.i. per acre (as found in EPA registered products such as Kocide[®] and Champ[®]) as a preventative disease spray tank mixture.

Use Neem Oil 70% on citrus to control mites at a rate of 1.0-2.0% in 25-200 gallons of water for high volume application or 0.5 gallon Neem Oil 70% in a minimum of 5 gallons of water per acre for low volume application. As a dormant oil spray, use Neem Oil 70% as a 2% solution or 1 quart Neem Oil 70% minimum per acre. Do not apply more than 5 gallons of Neem Oil 70% per acre, per application.

STONE AND TROPICAL FRUITS

FUNGICIDE USE: Apply Neem Oil 70% at a rate of 1.0% in 25-200 gallons of water for high volume applications or 0.5 gallons Neem Oil 70% in a minimum of 5 gallons of water per acre for low volume application. Reapply Neem Oil 70% every 7-14 days until threat of disease is over.

MITICIDE USE: Use Neem Oil 70% at first sign of pests or pest damage at the appropriate rate listed below as a foliar spray in sufficient water for thorough coverage. Repeat applications as needed for adequate control every 7-21 days. Apply a 1.0-2.0% solution of Neem Oil 70% for high volume application or a minimum of 1 quart Neem Oil 70% in a minimum of 5 gallons of water per acre for low volume application. Do not apply more than 5 gallons of Neem Oil 70% per acre, per application.

Do not use Neem Oil 70% after pit hardening on stone fruit.

VEGETABLE, CEREAL GRAINS AND OTHER MISCELLANEOUS CROPS **FUNGICIDE USE**

Apply Neem Oil 70% at the rate of 0.5% - 1.0% in 25 - 100 gallons of water per acre or 2 pints Neem Oil 70% in a minimum of 5 gallons of water per acre for low volume applications. Repeat applications every 7-14 days. Apply no more than 2 gallons Neem Oil 70% per acre, per application. Larger plants will require higher amounts of water to obtain good coverage. For mushrooms, apply Neem Oil 70% at a rate of 0.5% - 2.0% in 25 -100 gallons of water per 8,000 square feet. Repeat applications every 7 - 14 days.

MITICIDE USE

Apply Neem Oil 70% at a rate of 0.5% - 2.0% in 25 - 100 gallons of water per acre or 2 pints Neem Oil 70% in a minimum of 5 gallons of water per acre for low volume application. Apply no more than 2 gallons Neem Oil 70% per acre in solution per application. Apply in a solution of sufficient volume to achieve complete coverage of foliage. Reapply Neem 01/70% every 7 - 21 days as needed for adequate control. Larger plants will require higher amounts of water to obtain good soverage.

POME (DECIDUOUS FRUITS) AND NUTS

ςς γ ο___ FUNGICIDE USE: Apply Neem Oil 70% at a rate of 1.0% in 25 - 100 gallons of water per acre or 0.5 quarts Neem. Oil 70% in a minimum of 5 gallons of water per acre for low volume application. Apply no more than 235 gallons of Neem Oil 70% per acre in solution per application. Repeat applications every 7 - 14 days.

00000

MITICIDE USE: Use Neem Oil 70% at a rate of 1.0 - 2.0% in 25 - 100 gallons of water per acrearity 5 quarts Neem Oil 70% in a minimum of 5 gallons of water per acre for low volume application. Apply no more than 2 gallons of Neem Oil 70% per acre per solution. Apply at first sign of pests or pest damage. Repeat application as needed for adequate control every 7 - 21 days. Do not apply more than 5 gallons Neem Oil 70% per acre, per application. ວັວັັວ ່ວວວດ

Walnuts: Avoid spraying after husk split.

CITRUS, POME, STONE, AND TROPICAL FRUITS

Crops (including but not limited to)

Apples Apricots Avocados Bananas Calamondin Cherries Barbados cherries Sour cherries Sweet cherries Cherimoya Citrus citron Citrus hybrids Coconuts Crabapples Dates Feijoa

Figs Grapefruits Guavas Jujubes Kiwifruit Kumquats Lemons Limes Loquats Lychee Mangoes Mandarins Satsuma mandarins Nectarines Olives Oranges

Papayas Passion fruit Peaches Pears Oriental pears Pineapples Plantains Plums Pomegranates Prunes Pummelos Quinces Tangelos Tangerines

CUCURBITS

Refer to specific application instructions found directly below the heading Agricultural Use Directions.

Crops (including but not limited to)

Balsam pears (bitter melons) Honey balls Cantaloupes Melons (incl. hybrids) Casaba Honeydew melons Chinese waxgourds Muskmelons Citron melons Persian melons Crenshaw Pumpkins Cucumbers Squashes Gherkins Watermelons (incl. hybrids) Gourds Zucchini **SMALL FRUITS AND BERRIES** Refer to specific application instructions found directly below the heading Agricultural Use Directions. 55555 Crops (including but not limited to) ,,,, Blackberries зs Grapes ວັ້ງ ງວວງ Blueberries Huckleberries Boysenberries Loganberries Olallie berries Cranberries Currants Raspberries (black & red) Dewberries Strawberries Elderberries Youngberries Gooseberries

Crops (including but not limited to)

Almonds Beech nuts Brazil nuts Butternuts Cashews Chestnuts Chinquapins Filberts Hickory nuts Japanese horse chestnuts Macadamia nuts Pecans Pistachios Walnuts

BULB, COLE AND LEAFY VEGETABLES

Refer to specific application instructions found directly below the heading Agricultural Use Directions.

Crops (including but not limited to)

Arugula
Broccoli
Broccoli raab
Chinese broccoli (Gai Ion)
Brussels sprouts
Cabbage (head & leaf)
Chinese cabbage
(Bok choy, Napa)
Chinese mustard cabbage
(Gai choy)
Cauliflower
Celery
Celtuce
Chervil
Chicory
Red chicory
Cilantro
Collards
Corn salad
Cress
Watercress
Daikon
Endive
Fennel

Garlic	
Greens	
Collard greens	
Mustard greens	
Rape greens	
Kale	
Kohlrabi	
Leek	
Lettuce (head & leaf)	
Low-bok	
Onions	
Orach	
Parsley	
Purslane	
Raddichio	
Rappini	
Rhubarb	
Shallots	
Spinach	ر ز ز ز ز ر ب ر ب
Chinese Spinach	,
(Amaranth, Tampala)	د د د د د ر د ب
Swiss chard	Ĵ.
Turnip tops	د د د د د
	-
	ودددد

LEGUME AND FRUITING VEGETABLES

Refer to specific application instructions found directly below the heading Agricultural Use Directions.

Crops (including but not limited to)

Beans Broad beans Calabaza Chickpeas Eggplants Ground cherries Guar Lentils Peas Peanuts Pepinos Peppers Pigeon peas Snow peas Soybeans Tomatillos Tomatoes

ROOT AND TUBER VEGETABLES

Refer to specific application instructions found directly below the heading Agricultural Use Directions.

Crops (including but not limited to)

Artichokes Beets Cardone Carrots Cassava Ginger Ginseng Horseradish Parsnips

Potatoes Sweet potatoes Radishes Rutabaga Taro (Dasheen) Turmeric Turnips Yams Yam beans

	HERBS AND SPICES		
			<u>))))))))))))))))))) </u>
Refer to specific application in	structions found directly below the heading Agric		כנוכ נ
	Crops (including but not limited to)	ر ر د د د د د د د	د د د د
Anise	Marigold	5 5 5 C (
Balm	Marjoram	ι ε ε	ر زر ن ز ر
Basil	Mint	د د د د د د د	. د د د
Borage	Nasturtium	ວ່ວ ວັ ວ່າວວ່າ	2
Camomile	Pennyroyal		دددر د د
Caraway	Peppermint		ردرر
Catnip	Rosemary		ور ر د د
Celery	Rue		0000
Chives	Sage		
Coriander	Savory		
Cumin	Spearmint		
Curry leaf	Sweet bay		
Dandelion	Tarragon		
Dill	Thyme		
Fennel	Wintergreen		

VEGETABLE, CEREAL GRAINS AND OTHER MISCELLANEOUS CROPS

14/2

Crops (including but not limited to)

	Crops (molaung out not n	
Acerola	K	Linep
Alfalfa		Aushrooms
Araacaga (Apio)	N	lispero
Asparagus		Dats
Barley		Dkra
Breadfruit		Juenepa
Buckwheat		lice
Canola		afflower
Coffee		leagrapes
Coriander		lesame
Corn		Sorghum
Popcorn		Goursup (Quanabanas)
Sweet corn		tar apples
Cotton		ugarcane
Hops	S	Sunflowers
Jicama		Tobacco .
Jojoba		Vheat
Pest Control	Suppression	Diseases
Aphids	Whiteflies	Alternaria
Mealybugs	· Thrips	Anthracnose
Aites		Early blight
Rust mites		Leaf blight
Spider mites		Botrytis
Scales (Soft scale)		Greasy spot
		Leaf Spot
		Downy mildew
		Post-bloom fruit drop
		Post-bloom fruit drop Powdery mildew
		Post-bloom fruit drop Powdery mildew Molds
		Powdery mildew Molds Scabs
		Powdery mildew Molds
		Powdery mildew Molds Scabs Rusts Shothole
		Powdery mildew Molds Scabs Rusts Shothole
	• •	Powdery mildew Molds Scabs Rusts Shothole
		Powdery mildew Molds Scabs Rusts Shothole
	• .	Powdery mildew Molds Scabs Rusts Shothole
	· · · · · · · · · · · · · · · · · · ·	Powdery mildew Molds Scabs Rusts Shothole
	• .	Powdery mildew Molds Scabs Rusts Shothole
	• .	Powdery mildew Molds Scabs Rusts Shothole
·		Powdery mildew Molds Scabs Rusts Shothole
		Powdery mildew Molds Scabs Rusts Shothole
		Powdery mildew Molds Scabs Rusts Shothole

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep in original container. Store in a dry place, away from direct sunlight, feed, or foodstuffs. Keep container tightly sealed when not in use. Do not store below 40°F (4°C).

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

or for 250-gallon refillable containers:

Container Handling: Refillable container. Refill this container with pesticide only. Do no reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

ັວຍ ວ່ານວ່ອ ວ່າວວ່າວ

> a ar Jogoog

Master Label / Sublabel B - Ornamental leem FUNGICIDE/MITICIDE/INSECTICIDE

For Indoor/Outdoor Use On Ornamental Flowering Plants, Trees, Shrubs, and Foliage Grown on Commercial Farms and Nurseries, Greenhouses and Landscapes. Controls Fungal Diseases Including Black Spot, Mildews, Rusts, and Scab. Can be Used Up to Day of Harvest

FOR ORGANIC PRODUCTION

ACTIVE INGREDIENT:

Clarified Hydrophobic Extract of Neem Oil	
OTHER INGREDIENTS	<u>30%</u>
TOTAL	100%

This product contains 5.46 lb. of clarified hydrophobic extract of neem oil per US gallon.

> **KEEP OUT OF REACH OF CHILDREN CAUTION**

Net Contents: 2.5 Gallons

EPA Reg. No. 70051-2 EPA Est. No.70051-CA-001 Composition covered by US Patents 5,298,251; 5,356,629; 5,372,817; 5,405,612; and 5,409,708.

Lot No .:

.....

Manufactured by Certis USA, L.L.C. 9145 Guilford Road Suite 175 Columbia, MD 21046



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Removo and wash contaminated clothing before reuse. 5555

FIRST AID

ිಂ If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after วว์ออ the first 5 minutes, then continue rinsing eye. 0000

5552 Call a poison control center or doctor for treatment advice. Have the product container or label with you when calling, ົວວ່ວວ poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.

Personal Protective Equipment:

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or Viton.
- Shoes plus socks.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is hazardous to fish and aquatic invertebrates.

For Terrestrial Uses: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product through any type of irrigation system.

General Information:

- Do not use Triact 70 EC on food crops.
 Kills eggs, larvae, and adult insects.
 Shake well before using.
 For best results, maintain constant agitation in spray equipment.
 For optimal performance, do not mix with cold water (less than 45°F).
 Prevents and controls black spot on roses, anthracnose, rust and powdery mildew.
 - Do not apply to wilted or otherwise stressed plants, or to newly transplanted materials prior to root establishment.
 - Do not apply to known sensitive plant species, such as impatiens flowers, fuchsia flowers, hibiscus flowers, some rose flowers, ornamental olive trees, or some carnation varieties without prior testing.
 - Formulated for interiorscape use.
 - When used in conjunction with beneficial insects, conduct a small trial to assure compatibility before using on a large scale.
 - As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.

• Use with care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate or butyl rubber or nitrile rubber or neoprene rubber or polyvinyl chloride (PVC) or Viton.
- Shoes plus socks.

TANK MIXING: Neem Oil 70% has been found to be compatible with most commonly used fungicides, insecticides and fertilizers. Check physical compatibility first with other products before mixing by using a jar test. Using a quart jar, add the proportionate amounts of the products to 1 quart of water. After thoroughly mixing, let mixture stand for at least 5 minutes. Growers must tank mix combinations for phytotoxicity on a sample of plants prior to use. Due to the wide variation in climatic conditions, cultural practices, and other factors, the User assumes full responsibility for any crop damage or other liability resulting from the use of Neem Oil 70% in a tank mix combination. Avoid tank mixes with Morestan, Captan, Sulfur, Bravo, Echo, chlorothalonil or other chemically similar products as unpredictable results or leaf burn may occur.

GREENHOUSE, LANDSCAPE, AND NURSERY USE INSTRUCTIONS

0000 0000

, , ,

FUNGICIDE USE

ORNAMENTAL DISEASE CONTROL

Neem Oil 70% is an effective fungicide for the prevention and control of various fungal diseases including black opotion roses, powdery mildew, downy mildew, anthracnose, rust, leaf spot, botrytis, needle rust, scab, flower, twig, and tip blight, and alternaria.

Landscape, Greenhouse and Nursery Ornamentals

Neem Oil 70% will provide broad spectrum control of many fungal diseases affecting ornamental flowering plants, trees, shrubs, and foliage when used at the rate of 1 gallon per 100-200 gallons of water, depending on the level of disease pressure. Use the lower rate (1:200) when spraying as a preventative in a greenhouse while the higher rate (1:100) is needed for outdoor applications. Determine the best rate for your particular situation.

Greenhouse

When Neem Oil 70% is used as a preventative, the lower rate (1:200) will provide adequate control for most greenhouse situations. If disease pressure is evident, use the higher rate (1:100). Make applications for disease prevention on a 7-14 day schedule depending on the anticipated severity of the disease pressure. To control existing disease, apply Neem Oil 70% on a 7-Day schedule until the disease pressure is eliminated. Use a 14-Day schedule to prevent disease pressure from returning. NOTE: Spray greenhouse roses only in a rotation program, which includes Neem Oil 70% once every 4 weeks.

Landscape Ornamentals

In outdoor landscape applications, apply Neem Oil 70% at rates of 1:50 - 1:100 rate (see tables for detailed use rates). This will provide the necessary control under the extreme outdoor conditions. When using Neem Oil 70% as a preventative, make application on a 14-day schedule. If disease is present, use a 7-day application schedule to control the disease and prevent further damage from occurring.

APPLICATION INSTRUCTIONS

Apply until plant surface is uniformly wet with minimal runoff at 25-40 psi with hand sprayer or 100-200 psi with power sprayer as a fine spray to both leaf surfaces. Avoid excessive application.

MANAGEMENT OF TURF DISEASES					
SPRAY TARGET	CONCENTRATION	SPRAY INTERVAL	PRECAUTIONS		
Snow Mold	1:100-1:500 (1.0%-2.0%)	14 days	In Fall before first snow		
Dollar Spot	1:100-1:200 (1.0%-0.5%)	7-14 days	At first sign of disease		
Brown Patch	1:100-1:200 (1.0%- 0.5%)	7-14 days	At first sign of disease		

MANAGEMENT OF LANDSCAPE AND ORNAMENTAL DISEASES

SPRAY TARGET	CONCENTRATION	SPRAY INTERVAL	PRECAUTIONS	
Potted Plants (Greenhouse) Flowering Bench Crops (Greenhouse)	1:100 - 1:200 (1.0% - 0.5%) 1:100 - 1:200 (1.0% - 0.5%)	14 days 7 - 14 days	Trial first on open blooms Trial first on open blooms -not suggested for cut roses. Do not use on poinsettias.	
Outdoor Woody Plants	1:50 - 1:100 (2.0% - 1.0%)	14 days	00000000000000000000000000000000000000	

INSECTICIDE/MITICIDE USE

For use to control whiteflies, mealybugs, aphids, leafhoppers, mites and scales on ornamentals, trees, and shrubs in and around greenhouses, commercial nurseries and other commercial structures. >>

Use Neem Oil 70% at the rate of 1-2 gallons per 100 gallons of water. The low rate (1.0%) can be used, when pest pressure is low. Use the high rate (2.0%) for high pest infestations or to control more troublesome pests, such as mites and scales. Determine the best rate for your particular situation. 2222

00,00,

Make sprays on a 7-14 day interval depending on the severity of the pest problem. Use the 7-day schedule until the pest population is reduced. Then use a 14-day interval for control.

MIXING INSTRUCTIONS

Add Neem Oil 70% to one-half (1/2) full tank under agitation containing water of 45°F or greater before filling to desired level. If water temperature is below 45°F, premix Neem Oil 70% at a 1:1 ratio with tepid water to ensure good emulsification. Then dilute to final volume. When combining with other products, such as wettable powder insecticides or fungicides, add these items first when the tank is approximately 1/3 full. Ensure that there is good agitation while mixing for complete emulsification. Maintain agitation during spray application. Do not use if this oil does not form a uniform, cloudy emulsion.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Keep in original container. Store in a dry place, away from direct sunlight, feed, or foodstuffs. Keep container tightly sealed when not in use. Do not store below 40°F (4°C).

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on-site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Then offer for recycling, if available or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Certis USA, L.L.C. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

> 082407 091007 092607

Kocide[®] is a registered trademark of Griffin L.L.C. Champ[®] is a registered trademark of Agtrol Chemical Products Morestan[®] is a registered trademark of Bayer Inc. Captan[®] is a registered trademark of Drexel Chemical Company Bravo[®] is a registered trademark of Syngenta Crop Protection, Inc. Echo[®] is a registered trademark of Sipcam Agro USA, Inc.

Master Label / Sublabel C - Homeowner Neem C FUNGICIDE/MITICIDE/INSECTICIDE

For use on roses, flowers, houseplants, ornamental trees and shrubs, fruits, nuts, and vegetables, home lawns

For use in and around the home and home garden.

Controls blackspot, powdery mildew, rust, needle rust, downy mildew, spider mites, aphids, whiteflies, twig and tip blight, anthracnose, scab and flower, botrytis, alternaria, turf diseases, and other insect pests. Can be Used Up to Day of Harvest

For Organic Gardening

FOR ORGANIC PRODUCTION

ACTIVE INGREDIENT:

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside panels for additional precautionary statements.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.



Net Contents: 32 ounces (1 quart) (0.946 liters)

EPA Reg. No. 70051-2 EPA Est. No.70051-CA-001

Lot No.:

Manufactured by Certis USA, L.L.C. 9145 Guilford Road Suite 175 Columbia, MD 21046



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

BEE HAZARD

This product is toxic to bees exposed to direct treatment. Do not apply this product while bees are actively visiting the treatment area.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

GENERAL INFORMATION

- Do not add adjuvants (spreaders, stickers, extenders, etc.) to Neem Oil 70%.
- Stops powdery mildew in 24 hours.
- Kills eggs, larvae, and adult insects.
- Shake well before using.
- For optimal performance, do not mix with cold water (less than 45°F).
- Prevents and controls black spot on roses, anthracnose, rust and powdery mildew.
- Do not apply to wilted or otherwise stressed plants, or to newly transplanted materials prior to root establishment.

3 0 0 3 0 0

ردژدد

- Do not apply to known sensitive plant species, such as impatiens flowers, fuchsia flowers, hidiscus flowers, some rose flowers, ornamental olive trees, or some carnation varieties without prior testing.
- As with other oil-based products, exercise care in timing applications to early morning/late evening to minimize the potential for leaf burn.
- Use with care on plants with tender tissue. Check for leaf burn in small scale trials prior to use.

2000

, 5579 2, 55

33339.

FUNGICIDE USE

For use on Fruits, Vegetables, Herbs, Spices, Roses, Houseplants, Flowers, Trees and Shrubs, Home Lawns

Management of Turf Diseases

Neem Oil 70% may be used on all turf such as Kentucky bluegrass, perennial ryegrass, turf type tall fescue, Bermudagrass, zoysiagrass, and St. Augustinegrass around the home. Refer to table for detailed application rates.

MANAGEMENT OF TURF DISEASES					
Spray Target	Concentration	Spray Interval	Precautions		
Snow Mold	1:100-1:500 (1.0%-2.0%)	14 days	In Fall before first snow		
Dollar Spot	1:100-1:200 (1.0%-0.5%)	7-14 days	At first sign of disease		
Brown Patch	1:100-1:200 (1.0%- 0.5%)	7-14 days	At first sign of disease		

Application Instructions

Neem Oil 70% is an effective fungicide for the prevention and control of various fungal diseases including powdery mildew, black spot, downy mildew, anthracnose, rust, leaf spot, botrytis, needle rust, scab and flower, twig and tip blight, and alternaria. As a preventative, apply Neem Oil 70% on a 7 to 14 day schedule until disease development is no longer present. To control disease already present, apply Neem Oil 70% on a 7 day schedule until disease pressure is eliminated. Then continue spraying on a 14 day schedule to prevent the disease from reoccurring. To prevent rust leaf spot diseases, antracnose and scab, begin applications at the first sign of spring budding. To prevent powdery mildew, apply in midsummer or when disease is first detected. Continue applications until disease pressure no longer exists.

Mixing Instructions

Mix Neem Oil 70% at the rate of 2 tablespoons (1 fluid ounce) per gallon of water. Thoroughly mix solution and spray all plant surfaces (including undersides of leaves) until completely wet. Frequently mix solution as you spray.

Hose End Sprayer Use Instructions

The sprayer attached to the container is ready-to-use. Simply attach to your garden hose and follow these instructions;

- 1. Insure the large, round "on/off" knob is set to OFF.
- 2. Turn on water.
- 3. PUSH IN small knob near front of sprayer. This allows water and product to mix.
- 4. Hold sprayer and container level and point towards area to be sprayed.
- 5. Turn large, round "on/off" knob to ON.
- 6. Begin spraying product evenly over area you wish to treat.
- 7. Spray to visibly wet.
- 8. To stop spraying, turn large, round "on/off" knob to OFF.
- 9. PULL OUT small knob near front of sprayer. This prevents product from mixing with water.
- 10. Turn off water at faucet. Relieve water pressure in the hose by turning large, round "on/off" knob ON until water pressure is reduced.
- 11. Turn the knob to OFF for storage of unused product/and/or disposal of empty container.

Covers 5,000 sq. ft. or equivalent per 32 oz. container size.

INSECTICIDE/MITICIDE USE: For Use on FRUITS, VEGETABLES, HERBS, SPICES, ROSES, HOUSEPLANTS, FLOWERS, TREES, AND SHRUBS

APPLICATION INSTRUCTIONS:

Apply Neem Oil 70% at first sign of insects/mites. For control of aphids, spider mites, scale, whiteflies, beetles, leafrollers, and other insect pests. Neem Oil 70% is most effective when applied every 7 to 14 days. For heavy insect populations, spray on a 7-day schedule.

MIXING INSTRUCTIONS:

Mix Neem Oil 70% at the rate of 2-4 tablespoons (102 fluid ounces) per gallon of water. Thoroughly mix solution and spray all plant surfaces (including underside of leaves) until completely wet. Frequently mix solution as you spray.

STORAGE & DISPOSAL

Pesticide Storage: Keep in original container. Store away from direct sunlight, feed, or foodstuffs. Keep container tightly sealed when not in use. Keep from freezing.

Pesticide and Container Handling If empty: Nonrefillable container. Do not reuse or refill this container. Place in trash or Offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

Certis USA, L.L.C. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

The following claims and product information may or may not be presented on the product's label and labeling: $\frac{2}{2}$

- The _____ Promise Satisfaction guaranteed or your money back!
- If you apply _____ brand _____ as directed and you are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address below. We will refund the purchase price you paid.

ວ່ວງ ວ່ວງ ອີ່ງ

000000

ີ ອ ວິວີ ອ ວິ ສີ ອ

در د و در د

- Notice: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use pot in accordance with directions.
- Questions or Comments? Call ______ or visit our website at ______.
- 3 Garden Products in One!