

70051-52

09/24/2007

1/5

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Biopesticides and Pollution Prevention Division (7511P)  
1200 Pennsylvania Avenue NW  
Washington, D.C. 20460

EPA Reg. Number:

70051-52

Date of Issuance:

9/24/07

Term of Issuance:

Unconditional

Name of Pesticide Product:

Thuricide HPC

NOTICE OF PESTICIDE:

Registration  
 Reregistration (under FIFRA, as amended)

Name and Address of Registrant (Include ZIP Code):

Certis USA  
9145 Guilford Road  
Suite 175  
Columbia, MD 21046

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document, EPA has reregistered the product listed above. This action is taken under the authority of section 4(g)(2)(C) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain registration of your product.

A copy of the approved label for the product is enclosed for your records.

Signature of Approving Official:

*W. Michael McDavit*

Date:

9/24/07

W. Michael McDavit, Associate Director, Biopesticides and Pollution Prevention Division (7511P)

EPA Form 8570-6

CONCURRENCES

SYMBOL	7511P	7511P					
SURNAME	Reynolds	Reynolds					
DATE	9/26/07	9/26/07					

EPA Form 1320-1A (1/90)

Printed on Recycled Paper

OFFICIAL FILE COPY

# Thuricide HPC

2/5

## BIOLOGICAL INSECTICIDE

### HIGH POTENCY AQUEOUS CONCENTRATE

For control of insect pests in vegetables, fruits, nuts, field crops, and ornamentals.  
For control of insect pests in shade trees grown in nurseries and greenhouses.

#### ACTIVE INGREDIENT:

*Bacillus thuringiensis* subspecies *kurstaki* strain SA-12 solids, spores and  
Lepidopteran active toxins. (at least 6 million viable spores per milligram)\* .....15.0%

OTHER INGREDIENTS: .....85.0%

TOTAL .....100.0%

\* The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

Net Contents: 53 Gallons (200L)

EPA Reg. No. 70051-52

EPA Est. No. 70051-CA-001

Lot No.:

ACCEPTED

SEP 24 2007

KEEP OUT OF REACH OF CHILDREN  
CAUTION

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 70051-52

Manufactured by  
Certis USA, L.L.C.  
9145 Guilford Road  
Suite 175  
Columbia, MD 21046

CERTIS

#### Precautionary Statements

##### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing.

Wash thoroughly with soap and water after handling.

#### FIRST AID

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.  
Hot Line Number: 1-800-255-3924.

#### ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This product must not be applied aerially within 1/4 mile of any habitats of endangered or threatened lepidoptera. No manual application can be made within 300 feet of any threatened or endangered lepidoptera.

#### Personal Protective Equipment:

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, and P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Controls Statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendation:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product through any type of irrigation system. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Thorough coverage is essential when using Thuricide HPC Concentrate.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water wear:

- coveralls
- waterproof gloves
- shoes plus socks

**GENERAL**

Apply at first sign of infestation when worms are small and repeat at weekly intervals as needed to maintain control. Thuricide HPC attacks the larval gut and must be eaten by the insect to be effective. Thorough underleaf coverage is, therefore, essential for best results. After eating Thuricide HPC, worms stop feeding and die within 2-3 days. Since the active ingredient, *Bacillus thuringiensis*, is exempt from tolerance requirements, Thuricide HPC may be applied up to the day of harvest.

**MIXING**

Always shake or stir Thuricide HPC thoroughly before use. Pour application rate amount of Thuricide HPC onto the surface of water in a nearly filled spray tank. Maintain agitation. Use the higher rates against very heavy worm infestations or where it is difficult to get good coverage due to rank foliage. Do not allow dilute sprays to stand in the tank for more than 12 hours. Combinations with chemical insecticides, fungicides, spreader-stickers, and stickers in the tank are generally not deleterious to Thuricide HPC if the mix is used promptly.

**GROUND APPLICATION**

Use application rate amount of Thuricide HPC in a minimum of 5 to 50 gallons of water, depending on the type of crop and requirements of state regulations. Add a sticker, adhesive or deposit builder where heavy dew or rain is encountered.

**GROUND APPLICATION, TREES**

Use application rate amount of Thuricide HPC in 100 gallons of water in high pressure, high gallonage, hydraulic sprayers. Wet foliage thoroughly with spray but not to the point of excessive runoff. For mist blowers use application rate amount in 10 gallons of water. To prevent washoff by rain add a sticker to the finished spray.

**AERIAL APPLICATION**

Use application rate amount of Thuricide HPC in 5 to 20 gallons of water. A sticker should be added under conditions of heavy dew or rain. Apply early morning or evening when the air is calm.

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determines the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Crop	Pest	Quarts per acre
ALFALFA	Alfalfa Caterpillar	1/4
ALMONDS <sup>1</sup>	Redhumped Caterpillar	1 - 2
ALMONDS <sup>1</sup> , CHERRIES <sup>1</sup> Apply when caterpillars are actively feeding (2nd - 4th instar).	Tent Caterpillar	1 - 2
APPLES <sup>1</sup> , PEARS <sup>1</sup> Apply when eggs or newly hatched larvae first appear. Continue applying as a part of the normal cover spray program until pest is adequately controlled.	Redbanded Leafroller Tufted Apple Budmoth Variegated Leafroller Tent Caterpillar Fruit Tree Leafroller Gypsy Moth <sup>5</sup>	1 - 2 1 - 2 1 - 2 1 - 2 1 - 2 1 - 2
BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, KALE, MUSTARD GREENS, TURNIP GREENS	Diamondback Moth Imported Cabbage Worm Green Cloverworm Cabbage Looper	1/2 - 1 1/2 - 1 1/2 1/2 - 1 1/2 1 - 2
BEANS, BEETS, CARROTS, CELERY, CHARD, CHINESE CABBAGE (Choy sum), COS LETTUCE, ENDIVE, ESCAROLE, GARLIC, KOHLRABI, LENTILS, LETTUCE, ONIONS, PARSLEY, RADISHES, SPINACH, SQUASH	Cabbage Looper	1 - 2
CUCUMBERS, MELONS Begin treatment when worms are small to avoid damage and scarring to the crop.	Cabbage Looper	1 - 2
EGGPLANT, PEPPERS	Tomato Hornworm Tomato Fruitworm <sup>2</sup> Cabbage Looper	1/2 - 1 1/2 1 - 2 1 - 2
CURRENTS, BLUEBERRIES, CANE BERRIES (BLACKBERRIES, DEWBERRIES, RASPBERRIES) Apply by ground equipment only. Begin treatment as soon as possible after hatching and before larvae are protected by leaf folds.	Omnivorous Leafroller	1 - 2

Crop	Pest	Quarts per acre
COTTON	Tobacco Budworm	1 - 2
	Cotton Bollworm	1 - 2

For the suppression of light to moderate infestations, apply at first sign of egg laying or newly hatched worms (1st and 2nd instar larvae). Reapply at 3-5 day intervals to sustain worm suppression or until adequate reduction of worm population is achieved. Use THURICIDE HPC under a pest management program that includes close scouting. If heavy worm populations occur, it may be necessary to apply an insecticide with rapid knockdown and good larvicidal activity as a replacement. Apply from 5-15 gallons total spray per acre with ground equipment and from 2-5 gallons total spray per acre with aerial equipment.

**DIRECTIONS FOR USE OF TANK-MIX COMBINATIONS OF THURICIDE HPC AND CHLORDIMEFORM (FUNDAL<sup>3</sup> OR GALECRON<sup>4</sup> OR 4EC)\***

Crop	Insecticide	Pest	Rate per acre
COTTON	Thuricide HPC + chlordimeform SP* or chlordimeform 4EC*	Tobacco Budworm	1/2 - 1 pt. +
		Cotton Bollworm	1/8 - 1/4 lb. or 1/4 - 1/2 pt.

Use Thuricide HPC plus chlordimeform for a combined ovicide/larvicide effect. Use the low rates for suppression of light infestations, and the high rates for suppression of moderate infestations. Apply at first sign of egg laying or newly hatched worms. Reapply at 3-5 day intervals to sustain suppression, or until adequate worm reduction is achieved. Apply from 2-5 gallons total spray per acre with aerial equipment. Follow all restrictions, safety precautions and directions contained on the FUNDAL or GALECRON labels.

\*Not registered for use in California

Crop	Pest	Quarts per acre
COTTON	Cabbage Looper	1 - 2
PEANUTS	Cabbage Looper	1/2 - 1 1/2
	Velvetbean Caterpillar	1/2 - 1 1/2
	Podworm <sup>2</sup>	1 - 2
POTATOES	Cabbage Looper	1 - 2
SOYBEANS	Soybean Looper	1 - 1 1/2
	Cabbage Looper	1 - 1 1/2
	Green Cloverworm	1/2 - 1 1/2
	Velvetbean Caterpillar	1/2 - 1 1/2
	Podworm <sup>2</sup>	1 - 2
SUGARBEETS	Cabbage Looper	1 - 2
	Omnivorous Leafroller	1 - 2
TOBACCO	Cabbage Looper	1 - 1 1/2
	Tobacco Budworm	1 - 1 1/2
	Tobacco Hornworm	1 - 1 1/2
TOMATOES	Tomato Hornworm	1/2 - 1 1/2
	Cabbage Looper	1/2 - 1 1/2
	Tomato Fruitworm <sup>2</sup>	1 - 2
WATERMELONS (Florida)	Rindworm Complex	1/2 - 1 1/2

Crop	Pest	Quarts per acre
GRAPES	Grape Leaf Folder	1 - 2
		Start treating as soon as possible after hatching and before larvae are protected by leaf folds. Apply by ground equipment in up to 200 gallons total spray per acre to obtain thorough coverage of leaf surfaces.
GRAPES	Omnivorous Leafroller	1/2 - 1 1/2
MINT	Cabbage Looper	1 - 2
STRAWBERRIES	Cabbage Looper Roughskinned Cutworm	1 - 2
		Use 20 gallons water minimum per acre when using ground equipment and 5 gallons water minimum per acre by aircraft.
SUGAR MAPLE (greenhouse and nursery grown only)	Spring & Fall Cankerworm	1 - 2
	Fall Webworm	1 - 2
	Tent Caterpillar	1 - 2
	Gypsy Moth	1 - 2
PEACHES, NECTARINES	Redhumped Caterpillar	1 - 2
	Tent Caterpillar	1 - 2
	Omnivorous Leafroller	1 - 2
BANANAS (Hawaii)	Banana Skipper	1 - 2
		Use calibrated ground equipment with adequate water to apply to point of runoff.
ORANGES	Fruit Tree Leafroller Orange Dog	1 - 2
		1/2 - 1
ORANGES	Citrus Cutworm	1 - 2
		Use 50 gallons water minimum to 600 gallons maximum per acre when using ground equipment, and 10 gallons water minimum per acre by aircraft.
FILBERTS <sup>1</sup>	Tent Caterpillar* Filbert Leafroller**	1 - 2
		1 - 2
* Apply when caterpillars are actively feeding (2nd-4th instar.)		
** Apply at 10% egg hatch.		
PECANS <sup>1</sup>	Redhumped Caterpillar Fall Webworm Tent Caterpillar	1 - 2
		1 - 2
		1 - 2
WALNUTS <sup>1</sup>	Tent Caterpillar Fall Webworm Omnivorous Leafroller Redhumped Caterpillar	1 - 2
		1 - 2
		1 - 2
		1/4 - 1
Use 100 gallons water minimum per acre when using ground equipment, and 10 gallons water minimum per acre by aircraft.		

Crop	Pest	Quarts per acre
KIWI	Omnivorous Leafroller	1 - 2
	Fruittree Leafroller	1 - 2
	Cabbage Looper	1 - 2
*SHADE TREES, ORNAMENTALS *greenhouse and nursery grown shade trees only	Spring & Fall Cankerworm	1/2 - 1
	Fall Webworm	1/2 - 1
	Tent Caterpillar	1/2 - 1
	Redhumped Caterpillar	1/2 - 2
	Oakmoth Larvae	1 - 2
	Gypsy Moth <sup>5</sup>	1 - 2
STORED SOYBEANS, GRAINS	Indian Meal Moth	1 - 2
	Almond Moth	1 - 2

<sup>1</sup> To achieve adequate coverage of large trees or dense foliage, aerial application of not less than 15 gpa is preferred.

<sup>2</sup> Suppression only.

<sup>3</sup> FUNDAL 4EC (EPA Reg. No. 2139-100) Nor-Am Ag Prod., Inc. FUNDAL SP (EPA Reg. No. 2139-98) Nor-Am Ag Prod., Inc.

<sup>4</sup> GALECRON 4EC (EPA Reg. No. 100-551-AA) Ciba-Geigy Corp. GALECRON SP (EPA Reg. No. 100-554-AA) Ciba-Geigy Corp.

<sup>5</sup> Gypsy Moth: Aerial Application. THURICIDE HPC may also be applied by air to prevent excessive defoliation by Gypsy Moth. For best results, spray when leaf expansion reaches 40% - 50%, and respray 14 days later. Use two quarts of THURICIDE HPC in 1-4 gallons finished spray per acre. Incorporate in the spray mix an evaporation retardant, such as molasses, Cargill's Insecticide Base polyethylene glycol E-400, and a film forming sticker. To obtain thorough coverage use spray equipment suitable to deliver droplets with an average diameter of less than 300 microns.

To control and prevent Indian Meal Moth and Almond Moth infestations of stored grains and soybeans, prepare a spray mixture which includes one gallon of water for every 6 fluid ounces THURICIDE HPC. The spray mixture may be applied either by treating the top 4 inches of grain as it is being augured into storage (applying 0.6 pint of mixture per bushel in the grain stream), or by treating the surface of grain after it is in the bin. The table below can be used as a guide in determining the total amount of THURICIDE HPC needed according to the bin diameter or the number of bushels to be treated.

Bin Diameter (ft.)	Surface Area (sq. ft.)	Bushels (to 4 inch depth)	THURICIDE HPC Rate
8	50	13	6 fluid oz.
12	113	30	14 fluid oz.
16	201	53	24 fluid oz.
20	314	84	1 qt. 2 fluid oz.
24	452	120	1 qt. 20 fluid oz.
28	615	163	2 qts. 8 fluid oz.
32	804	214	2 qts. 28 fluid oz.

• To insure thorough coverage when making applications to the grain surface after it is in the bin, apply spray mixture in three applications. Mix the grain with a scoop or rake to a depth of 4 inches after each application.

• Stored grain may be treated anytime, but for best results, treat

grain at the time it is placed into storage or shortly thereafter, or in the early spring prior to egg laying. Full season control is normally experienced.

• Retreat only if reinfestation occurs.

• For the protection of bagged grain, apply spray mixture to entire grain mass, and mix thoroughly prior to bagging. THURICIDE HPC at 1 1/2 qts. per 10 gallons of water will treat approximately 100 bushels.

• Treated grain may be used at any time after treatment.

**Greenhouse Application**

THURICIDE HPC may also be used in the greenhouse and on flowers at 1/2 - 1 qt. per 100 gallons of water for control of listed insects on crops on this package label.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal.

**Pesticide Storage:** Store in a cool place. Activity may be impaired by storage at temperatures above 90°F. Protect from freezing.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**WARRANTY**

Certis USA, L.L.C. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OF MERCHANTABILITY IS MADE.