



Please read instructions on reverse before completing form.

Form Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States  
Environmental Protection Agency  
Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

### Application for Pesticide - Section I

1. Company/Product Number 70051-80	2. EPA Product Manager Phil Hutton	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Condor WP	PM#	
5. Name and Address of Applicant (Include ZIP Code) Certis USA, L.L.C. 9145 Guilford Road, Suite 175 Columbia, MD 21046 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

### Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)  
New establishment number on label.

### Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Metal Plastic Glass Paper Other (Specify) _____		
<i>certification must be submitted</i>		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/>	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

### Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name Christine A. Dively		Title Director of Regulatory Affairs		Telephone No. (include Area Code) 301-483-3806	
<b>Certification</b> I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature <i>Christine A. Dively</i>		3. Title Director of Regulatory Affairs			
4. Typed Name Christine A. Dively		5. Date April 9, 2003			

# CERTIS

Certis USA  
9145 Guilford Road  
Suite 175  
Columbia, MD 21046

(301) 604-7340

FAX (301) 604-7015  
www.certisusa.com

276

April 14, 2003

Mr. Alan Reynolds  
Biopesticides & Pollution Prevention Division (7511C)  
Office of Pesticide Programs  
1921 Jefferson Davis Hwy.  
Crystal Mall 2 Building  
Arlington, VA 22202

RE: New Establishment Number and Updated First Aid Statements

Dear Mr. Reynolds:

Certis USA L.L.C. (EPA Company No. 70051) is respectfully submitting the enclosed EPA Forms 8570-1. The purpose for these notifications is to change the establishment number on the following labels:


Raven Oil Flowable, EPA Reg. #70051-82 ✓  
Lepinox G, EPA Reg. #70051-88 ✓  
Cutlass Wettable Powder, EPA Reg. #70051-79 ✓  
Condor XL, EPA Reg. #70051-85 ✓  
Condor WP, EPA Reg. #70051-80 ✓  
Condor Bioinsecticide, EPA Reg. #70051-78 ✓  
Crymax WP, EPA Reg. #70051-90

These products were all formerly owned by Ecogen, Inc. and it has just recently come to our attention that the establishment number is incorrect. Therefore, we have corrected the establishment number on all of the enclosed labels. In addition, the following amendments are included to both update the First Aid Statement in accordance with PR Notice 2001-1, and change the establishment number.

Lepinox WDG, EPA Reg. #70051-89  
Crymax Bioinsecticide, EPA Reg. #70051-86

Enclosed you will find:

- 1) Two copies of the Application for Pesticide Notification (EPA Form 8570-1) for each product submitted.
- 2) Two copies of each label for notification.
- 3) Five copies of each label for amendment.



# Condor<sup>®</sup>WP

## WETTABLE POWDER BIOINSECTICIDE

CONDOR<sup>®</sup> wettable powder bioinsecticide is a biological insecticide for the control of lepidopteran pests.

**Active Ingredient:**

*Bacillus thuringiensis* subspecies *kurstaki* strain EG2348 solids, spores and Lepidopteran active toxins .....43.0%

**Inert Ingredients:** .....57.0%

**Total** .....100.0%

The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

**Net Contents:**

5 U.S. Pound Bag

EPA Reg. No. 70051-80

EPA Est. No. 62171-MS-001

### NOTIFICATION

Date Reviewed: 6-13-03  
Reviewed By: [Signature]

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

<b>FIRST AID</b>
<b>If on skin or clothing:</b> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.
<b>If in eyes:</b> Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after 5 minutes, then continue rinsing eye.
<b>If inhaled:</b> Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
<b>If swallowed:</b> Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.
Call Poison control center or doctor for treatment advice.

Manufactured by  
Certis USA  
9145 Guilford Road  
Suite 175  
Columbia, MD 21046

## CERTIS

Hot Line Number: 1-800-255-3924

### PRECAUTIONARY STATEMENTS

#### CAUTION

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled. Avoid breathing dust or spray mist. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

#### Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

<b>User Safety Recommendations</b>
User should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Storage:** Store in a cool, dry place inaccessible to children.

**Pesticide Disposal:** Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

<b>AGRICULTURAL USE REQUIREMENTS</b>
Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this section only apply to uses of this product that are covered by the Worker Protection Standard.
Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.
PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Do not enter treated areas without protective clothing until sprays have dried.**

**Preharvest Interval:** CONDOR WP may be applied to the crops listed in the APPLICATION RATE TABLE at any time, up to and on the day of harvest.

**Mode of Action:** After consuming a lethal dose of CONDOR WP, larvae will cease to feed, but may remain alive on foliage for several days before disappearing. Immediately after ingestion of CONDOR WP, larvae begin to move slowly, become discolored, shrivel and blacken prior to death.

**MIXING INSTRUCTIONS**

CONDOR WP may be applied with conventional ground, aerial or hand held application equipment with quantities of water sufficient to provide thorough coverage of infested plants. Do not apply this product through any type of irrigation system. To obtain a suitable mixture with water, add enough water to allow maximum agitation. With agitator running, slowly add in the CONDOR WP. Continue agitation. Add remainder of water and other spray materials and agitate until mixed. Maintain suspension with loading and spraying. Do not mix more CONDOR WP than can be used in a 12-hour period. Rinse and flush spray equipment thoroughly following each use. Do not contaminate water when disposing of equipment washwaters.

In order to make proper decisions on application rates to be used, follow the recommendations in the APPLICATION RATE TABLE and these guidelines:

**APPLICATION GUIDELINES**

Pest category	Pest Pressure (number of larvae/plant)			
	Low <sup>1</sup> (<0.3)	Moderate <sup>2</sup> (0.3-1.0)	High <sup>3</sup> (1.0-5.0)	Extreme <sup>4</sup> (>5.0)
<b>Pounds of Product to be Applied per Acre</b>				
Category 1	1.5	2.0	2.0	2.0
Category 2	1.0	1.5	2.0	2.0
Category 3	.5	1.0	1.5	2.0

<sup>1</sup> Recommended spray interval of 7 - 10 days.

<sup>2</sup> Recommended spray interval of 6 - 8 days.

<sup>3</sup> Recommended spray interval of 4 - 6 days.

<sup>4</sup> Recommended spray interval of 3 - 5 days.

**Category 1 Pests include:** artichoke plume moth, navel orangeworm, oriental fruit moth, tomato fruitworm (also called bollworm and corn earworm), tufted apple budmoth.

**Category 2 Pests include:** Amorbia, armyworms, cabbage looper, citrus cutworm, diamondback moth, leafrollers, melonworm, peach twig borer, pickleworm, soybean looper, tomato pinworm, tobacco budworm and tortrix moth.

**Category 3 Pests include:** all caterpillar pests shown in the APPLICATION RATE TABLE, except those shown in Categories 1 and 2.

For crops such as Fruits, Nuts and Vines, applications are often timed to stage of development and recommendations from local Extension personnel should always be followed.

**APPLICATION INSTRUCTIONS**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

CONDOR WP is a selective insecticide for use against the lepidopteran larvae listed in the APPLICATION RATE TABLE. Larvae must consume deposits of CONDOR WP to be affected. Always follow these directions:

- Make applications when larvae are still small (early instars) and actively feeding on foliage or other plant parts.
- Make applications before noticeable foliar damage occurs.
- Thorough spray coverage is essential for good insect control. For ground applications, directed drop nozzles should be used for certain vegetable crops.
- When insect infestations are heavy, use the higher label rates, shorten the spray interval, and/or use larger total spray volume to improve spray coverage (see APPLICATION GUIDELINES for selection of rates and intervals).
- Applications should be repeated at an interval sufficient to maintain control, depending upon plant growth, insect pressure and weather conditions after spraying. (Refer to APPLICATION GUIDELINES).
- For ground applications, use a minimum spray volume of 20 gallons per acre. For aerial applications, use a minimum spray volume of 5 gallons per acre.
- For orchard applications, a spray volume of 100 gallons per acre and treatment of each row is recommended.
- Do not use screens smaller than 50 mesh.
- Local conditions may affect the use of CONDOR WP. Consult your State Agricultural Extension Specialist for specific recommendations related to local crop protection problems.
- Spray water/spray tank solutions should not exceed pH 8.0. If necessary, buffer water to near neutral pH.

**HAND-HELD EQUIPMENT**

When using hand-held equipment, mix 3 teaspoons per gallon of water or 1 1/2 pounds per 100 gallons of spray solution. Spray to wet, but not to runoff.

**TANK-MIX**

Combinations of CONDOR WP with commonly used insecticides, fungicides, or other spray tank adjuvants are generally not deleterious to performance. It is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities prior to mixing in spray tank. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rate should be exceeded. Application must be made in accordance with the more restrictive of label limitations and precautions.

- For improved durability of spray deposits, an approved spreader/sticker is strongly recommended for hard-to wet crops such as cole crops.

**APPLICATION RATE TABLE**

**I. VEGETABLES & COLE CROPS  
(Fresh and Processed)**

Crop such as:	Insect Pest
Artichokes	Leeks
Arugala	Lentils
Asparagus	Lettuce (Head
Beans	Leaf, Romaine)
Beets	Malanga
Bok Choy	Melons (Cantaloupe,
Broccoli	Crenshaw, Honeydew,
Brussels sprouts	Muskmelon,
Cabbage	Watermelon, etc.)
Cardoni	Napa
Carrots	Okra
Cauliflower	Onions
Celeriac	Parsley
Celery	Parsnips
Chick peas	Peas
Chicory	Peppers
Chinese cabbage	Potatoes
Collards	Pumpkins
Cucumber	Radishes
Cucurbits	Rutabaga
Dry bulb onions	Salsify
Eggplants	Shallots
Escarole	Soybean foliage
Endive	Spinach
Garlic	Squash
Green onions	Sugar Beets
Greens (Beets,	Sweet potatoes
China, Dandelion,	Swiss Chard
Mustard, Turnip)	Tomatoes
Horseradish	Turnips
Kale	Watercress
Kohlrabi	

Rate/Acre: 0.5 - 2.0 pounds

**II. HERBS & SPICES**

Crop such as:	Insect Pest
Basil	Armyworms
Chives	Diamondback moth
Cilantro	European corn corer
Dill	Green cloverworm
Oregano	Imported cabbageworm
Peppermint	Loopers
Thyme	Saltmarsh caterpillar

Rate/Acre: 0.5 - 2.0 pounds

**III. PASTURE & HAY CROPS**

Crop such as:	Insect Pest
Alfalfa (hay & seed)	Alfalfa caterpillar
Pasture (grasses & hay)	Armyworms*
Silage	European skipper
	Loopers*
	Webworm

Rate/Acre: 0.5 - 2.0 pounds

\* Product should be applied when early instar larvae first appear. If infestation persists, make a second application 7-10 days later. Combination of CONDOR with contact insecticide is recommended for control of 4th and 5th instar larvae.

**IV. FRUIT, NUT AND VINE CROPS**

Crop such as:	Insect Pest	Rate/Acre
Pome and Stone Fruit Trees:	Cankerworm (Spring & Fall)	0.5 - 2.0 pounds
Apples	Eastern tent caterpillar	
Apricots	Fall webworm	
Cherries	Fruitree leafroller	
Nectarines	Gypsy moth	
Peaches	Navel orangeworm	
Pears	Omnivorous leafroller	
Plums	Oriental fruit moth	
Prunes	Pandemis leafroller	
Quince	Peach twig borer	
	Redbanded leafroller	
	Redhumped caterpillar	
	Tortrix moth (Orange and Garden)	
	Tufted apple budmoth	
	Variegated leafroller	
	Walnut caterpillar	
Nut Trees:	Citrus cutworm	0.5 - 2.0 pounds
Almonds	Filbert leafroller	
Chestnuts	Filbert webworm	
Filberts	Navel orangeworm	
Pecans	Oblique banded leafroller	
Walnuts	Peach twig borer	
	Roughskinned cutworm	
Citrus	Amorbia	0.5 - 2.0 pounds
	Citrus cutworm	
	Fruitree leafroller	
	Orangedog	
Small Fruit and Berries:	Acherna sphinx moth	0.5 - 2.0 pounds
Blackberries	Armyworms	
Blueberries	Blueberry leafroller	
Cranberries	Fruitree leafroller	
Currants	Grape berry moth	
Raspberries	Gypsy moth	
Strawberries	Loopers	
	Oblique banded leafroller	
	Tobacco budworm	
Grapes	Grape berry moth	0.5 - 2.0 pounds
	Cherry fruitworm	
	Grape leafroller	
	Grapeleaf skeletonizer	
	Green fruitworm	
	Omnivorous leafroller	
	Orange tortrix	
	Saltmarsh caterpillar	
Tropical and Other Fruit:	Amorbia	0.5 - 2.0 pounds
Avocados	Loopers	
	Orange tortrix	
	Omnivorous leafroller	
	Omnivorous looper	
	Spanworm	
Bananas	Banana skipper	0.5 - 2.0 pounds.
Kiwi	Omnivorous leafroller	0.5 - 2.0 pounds
Persimmons	Citrus cutworm	0.5 - 2.0 pounds
Pomegranate	Fall webworm	
	Filbert webworm	
	Omnivorous leafroller	
	Redhumped caterpillar	
	Tent caterpillar	
Pineapple	Gummosis-Batrachedra cornosae	0.5 - 2.0 pounds
	Thecla-Thecla basiliides	
Tropical fruits	Hornworms	0.5 - 2.0 pounds
	Leafrollers	
	Loopers	
	Omnivorous leafroller	

V. FIELD CROPS

Crop such as:	Insect Pest	Rate/Acre
Canola/Rape Seed Evening Primrose	Armyworms Diamondback moth Imported cabbageworm Loopers	0.5 - 2.0 pounds
Corn (Field, Sweet, Popcorn)	Armyworms European corn borer Southwestern corn borer	0.5 - 2.0 pounds
Cotton*	Beet armyworm Bollworm Cabbage looper Cotton leaf perforator Saltmarsh caterpillar Tobacco budworm	0.5 - 2.0 pounds
Hops	Armyworms Loopers Oblique banded leafroller Omnivorous leaf-tier Spotted cutworm	0.5 - 2.0 pounds
Joboba	Looper ( <i>Anacamptodes spp.</i> )	0.5 - 2.0 pounds
Peanuts	Green cloverworm Loopers Podworms Velvetbean caterpillar	0.5 - 2.0 pounds
Rice	Armyworms Green cloverworm Loopers Saltmarsh caterpillar Velvetbean caterpillar	0.5 - 2.0 pounds
Safflower	Armyworms Loopers Saltmarsh caterpillar	0.5 - 2.0 pounds
Small Grains (Barley, Oats, Rye, Wheat, etc.)	Armyworms Loopers	0.5 - 2.0 pounds
Sorghum	European corn borer Saltmarsh caterpillar Velvetbean caterpillar	0.5 - 2.0 pounds
Soybeans	Green cloverworm Soybean looper Velvetbean caterpillar	0.5 - 2.0 pounds
Sunflowers	Banded sunflower moth Beet armyworm Headmoth Loopers Sunflower moth	0.5 - 2.0 pounds
Tobacco	Tobacco budworm Tobacco hornworm Loopers	0.5 - 2.0 pounds

\* Use CONDOR to control light to moderate populations of newly hatched worms in integrated pest management programs. Repeat treatments at four to five day intervals or as long as necessary until results are acceptable. For control of cotton bollworm and tobacco budworm, ovicides or synthetic pyrethroids can be combined with CONDOR in accordance with the more restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

VI. COMMERCIAL FLOWERS & ORNAMENTAL PLANTS

Crop such as:	Insect Pest	
Bedding plants	Armyworms	Loopers
Flowers	Azalea moth	Oleander moth
Greenhouse	Diamondback moth	Omnivorous leafroller
Ornamentals	Ello moth (hornworm)	Omnivorous looper
Vegetables	lo moth	Tobacco budworm

Rates/Acre 0.5 - 2.0 pounds

VII. FOREST, SHADE TREE & NURSERY STOCK

Crop such as:	Insect Pest	
Forest	Bagworm	Pine butterfly
Shade trees	Blackheaded budworm	Redhumped caterpillar
Nursery trees	Browntail moth	Saddleback caterpillar
	California oakworm	Saddle prominent caterpillar
	Cankerworm (Spring and Fall)	Spruce budworm
	Douglas fir tussock moth	Tent caterpillar
	Elm spanworm	Tortix
	Fall webworm	Western tussock moth
	Fruitree leafroller	
	Greenstriped mapleworm	
	Gypsy moth	
	Jack pine budworm	
	Mimosa webworm	

Rate/Acre 0.5 - 2.0 pounds

VIII. TURF

Crop such as:	Insect Pest	
Turf	Armyworms Sod webworm	Tropical sod webworm

Rate/Acre 0.5 - 2.0 pounds

WARRANTY

Certis USA, L.L.C. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.