

70051-90

10f8



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

March 23, 2010

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Christine A. Dively
Director of Regulatory Affairs
CERTIS USA, LLC
9145 Guilford Road, Suite 175
Columbia, MD 21046

RE: Product Name: Crymax WP
EPA Reg. No: 70051- 90
Application for Notification dated November 13, 2009, to add container reuse statement to the Storage and Disposal section as permitted by PR Notices 2007-4 and 2008-1.

Dear Ms. Dively,

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under Pesticides Registration Notice (PRN) 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PRN 98-10 and it has been determined that the action(s) requested falls within the scope of PRN 98-10. Our records have been duly noted, and the letter submitted with this application has been stamped "Notification, received and accepted" and will be placed accordingly in our records.

Questions concerning this action should be directed to Mary Paden (703) 308-0411 or email at paden.mary@epa.gov.

Sincerely,

Sheryl K. Reilly

Sheryl K. Reilly, Ph.D., Chief
Microbial Pesticides Branch
Biopesticides and Pollution Prevention
Division (7511P)

CONCURRENCES							
SYMBOL	▶ 7511P						
SURNAME	▶ Paden						
DATE	▶ 3-18-10						

2098

Please read instructions on reverse before completing form.

Form Approved

B No. 2070-0060

Print Form



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 70051-90	2. EPA Product Manager Microbial Pesticides Branch	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Crymax WP	PM# 92	
5. Name and Address of Applicant (Include ZIP Code) Certis U.S.A., L.L.C. 9145 Guilford Road, Suite 175 Columbia, MD 21788 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: NOTIFICATION accepted <input checked="" type="checkbox"/> EPA Reg. No. _____ Date Reviewed: 03-18-10 Product Name _____ Reviewed By: m J Pader	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container 5 Pound Bags	5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)

Name Christine Dively	Title Director Reg. Affairs	Telephone No. (Include Area Code) 301-483-3806
--------------------------	--------------------------------	---

Certification

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received (Stamped)

2. Signature <i>Christine A. Dively</i>	3. Title Director Reg. Affairs	6. Date Application Received (Stamped)
4. Typed Name Christine Dively	5. Date November 13, 2009	

4088

Crymax[®]

WETTABLE POWDER BIOINSECTICIDE

CRYMAX[®] wettable powder bioinsecticide is a biological insecticide for the control of lepidopteran pests.

Active Ingredient:

Bacillus thuringiensis subspecies *kurstaki* strain EG7841 solids, spores and Lepidopteran active toxins*40.00%

Other Ingredients:60.00%

Total100.00%

*The percent active ingredient does not indicate product performance and potency measurements are not federally standardized.

Net Contents:

5 U.S. Pound Bag

EPA Reg. No. 70051-90

EPA Est. No. 62171-MS-001

Lot No.:

NOTIFICATION *Accepted*

Date Reviewed: 03-18-10

Reviewed By: *m. J. Pader*

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

Call a Poison control center or doctor for further treatment advice.

Hot Line Number: 1-800-255-3924

Manufactured by

Certis USA, L.L.C.

9145 Guilford Road

Suite 175

Columbia, MD 21046

CERTIS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled. Avoid breathing dust or spray mist. Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Shoes plus socks
- Waterproof gloves

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Mixers/loaders and applicators must wear a dust/mist filtering respirator meeting NIOSH standards of at least N-95, R-95, or P-95. Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters.

This product must not be applied aerially within 1/4 mile of any habitats of endangered or threatened lepidoptera. No manual application can be made within 300 feet of any threatened or endangered lepidoptera.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

5048

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this section only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks.

Preharvest Interval: CRYMAX® may be applied to the crops listed in the APPLICATION RATE TABLE at any time, up to and on the day of harvest.

Mode of Action: After consuming a lethal dose of CRYMAX®, larvae will cease to feed, but may remain alive on foliage for several days before disappearing. Immediately after ingestion of CRYMAX® larvae begin to move slowly, become discolored, shrivel and blacken prior to death.

MIXING INSTRUCTIONS

CRYMAX® may be applied with conventional ground, aerial or hand held application equipment with quantities of water sufficient to provide thorough coverage of infested plants. To obtain a suitable mixture with water, add enough water to allow maximum agitation. With agitator running, slowly add in the CRYMAX®. Continue agitation. Add remainder of water and other spray materials and agitate until mixed. Maintain suspension while loading and spraying. Do not mix more CRYMAX® than can be used in a 24-hour period. Rinse and flush spray equipment thoroughly following each use. Do not contaminate water when disposing of equipment washwaters.

In order to make proper decisions on application rates to be used, follow the recommendations in the APPLICATION RATE TABLE.

APPLICATION INSTRUCTIONS

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

CRYMAX® is a biological insecticide for use against the lepidopteran larvae listed in the APPLICATION RATE TABLE. Larvae must consume deposits of CRYMAX® to be affected. Always follow these directions:

- Make applications when larvae are still small (early instars) and actively feeding on foliage or other plant parts.
- Make applications before noticeable foliar damage occurs.
- Thorough spray coverage is essential for good insect control. For ground applications, directed drop nozzles should be used for certain vegetable crops.
- When insect infestations are heavy, use the higher label rates, shorten the spray interval, and/or use larger total spray volume to improve spray coverage.

- Applications should be repeated at an interval sufficient to maintain control, depending upon plant growth, insect pressure and weather conditions after spraying.
- For crops such as Fruits, Nuts, and Vines, applications are often timed to stage of development and recommendations from local Extension personnel should always be followed.
- Local conditions may affect the use of CRYMAX®. Consult your State Agricultural Extension Specialist for specific recommendations related to local crop protection problems.
- Spray water/spray tank solutions should not exceed pH 8.0. If necessary, buffer water to near neutral pH.

HAND HELD EQUIPMENT

When using hand held equipment, mix 3 teaspoons per gallon of water or 2 pounds per 100 gallons of spray solution. Spray to wet, but not to runoff.

TANK MIX

Combinations of CRYMAX® with commonly used insecticides, fungicides, or other spray tank adjuvants are generally not deleterious to performance. It is advisable to test physical compatibility by mixing all components in a small container in proportionate quantities prior to mixing in spray tank. This product cannot be mixed with any product containing a label prohibition against such mixing. No label dosage rate should be exceeded. Application must be made in accordance with the more restrictive of label limitation and precautions.

For improved durability of spray deposits, a spreader/sticker approved for use on growing crops may be used for hard-to-wet crops such as cole crops.

CHEMIGATION

Apply this product only through center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move sprinkler systems. **Do not apply this product through any other type of irrigation system.** Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

If you have questions about calibration, contact your State Extension Service Specialist, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

CHEMIGATION SYSTEM CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be

600f8

discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

The active ingredient in CRYMAX® will settle in the tank and injection lines; adequate agitation must be provided before and during the injection period. Use only in systems that apply uniformly and have appropriate check valves. When application is complete, thoroughly flush the injection system and sprinkler lines.

MIXING RECOMMENDATIONS FOR CHEMIGATION:

Follow general MIXING INSTRUCTIONS and keep the ratio at three parts water to one part CRYMAX®. Also, provide mild uniform agitation throughout the solution but do not agitate excessively.

SPRAY VOLUME

For chemigation, use irrigation levels of 0.15 to 0.5 inches of water per acre. Up to 1 inch of irrigation water may be used, but efficacy may be reduced. The product should be applied continuously for the duration of the water application.

APPLICATION RATE TABLE

I. Vegetable & Cole CROPS		
Crop such as:		Insect Pest
Artichokes	Lentils	Alfalfa looper
Arugala	Lettuce (Head,	Armyworm
Asparagus	Leaf, Romaine)	Artichoke plume moth
Beans	Malanga	Beet armyworm
Beets	Melons	Cabbage budworm
Bok Choy	(Cantaloupe,	Cabbage looper
Broccoli	Crenshaw,	Cabbage Webworm
Brussels sprouts	Honeydew,	Celery leaf-tier
Cabbage	Muskmelon,	Corn earworm
Cardoni	Watermelon,	Cross-striped
Carrots	etc.)	cabbageworm
Cauliflower	Napa	Diamondback moth
Celeriac	Okra	European corn borer
Celery	Onions	Green cloverworm
Chick peas	Parsley	Imported cabbageworm
Chicory	Parsnips	Melonworm
Chinese cabbage	Peas	Omnivorous leafroller
Collards	Peppers	Pickleworm
Cucumber	Potatoes	Rindworm complex
Cucurbits	Pumpkins	Saltmarsh caterpillar
Dry bulb onions	Radishes	Southern armyworm
Eggplant	Rutabaga	Soybean looper
Escarole	Salsify	Tobacco budworm
Endive	Shallots	Tomato fruitworm
Garlic	Soybean foliage	Tomato hornworm
Green onions	Spinach	Tomato pinworm
Greens (Beets,	Squash	Velvetbean caterpillar
China,	Sugar beets	Yellowstriped armyworm
Dandelion,	Sweet potatoes	
Mustard,	Swiss chard	
Turnip)	Tomatoes	
Horseradish	Turnips	
Kale	Watercress	
Kohlrabi		
Rate/Acre: 0.5 - 2.0 pounds		
II. HERBS & SPICES		
Crop such as:		Insect Pest
Basil		Alfalfa looper
Chives		Armyworm
Cilantro		Diamondback moth
Dill		European corn borer
Oregano		Green cloverworm
Peppermint		Imported hornworm
Thyme		Loopers
		Saltmarsh caterpillar
Rate/Acre: 0.5 - 2.0 pounds		
III. PASTURE & HAY CROPS		
Crop such as:		Insect Pest
Alfalfa (hay & seed)		Alfalfa caterpillar
Pasture (grasses & hay)		Armyworm
Silage		Beet armyworm
		European skipper
		Loopers
		Webworm
		Yellowstriped armyworm
Rate/Acre: 0.5 - 2.0 pounds		

7098

IV. FRUIT, NUT & VINE CROPS

Crop such as:	Insect Pest	
Pome and Stone	Cankerworm (Spring & Fall)	Pandemis leafroller
Fruit Trees:	Cherry fruitworm	Peach twig borer
Apples	Eastern tent caterpillar	Redbanded leafroller
Apricots	Fall webworm	Redhumped caterpillar
Cherries	Fruittree leafroller	Tortrix moth
Nectarines	Green fruitworm	(Orange and Garden)
Peaches	Gypsy moth	Tufted apple budmoth
Pears	Navel orangeworm	Variegated leafroller
Plums	Obliquebanded leafroller	Walnut caterpillar
Prunes	Omnivorous leafroller	Western tent caterpillar
Quince	Oriental fruit moth	
Nut Trees:	Citrus cutworm	Omnivorous leafroller
Almonds	Filbert leafroller	Pecan nut casebearer
Chestnuts	Filbert webworm	Peach twig borer
Filberts	Fruittree leafroller	Redhumped caterpillar
Pecans	Hickory shuckworm	Roughskinned cutworm
Pistachios	Navel orangeworm	Western tent caterpillar
Walnuts	Obliquebanded leafroller	
Citrus:	Amorbia	Omnivorous leafroller
	Citrus cutworm	Orangedog
	Fruittree leafroller	
Small Fruit and Berries:	Achema sphinx moth	Omnivorous looper
Blackberries	Armyworms	Tobacco budworm
Blueberries	Blackheaded fireworm	
Boysenberries	Blueberry leafroller	
Cranberries	Cranberry girdler	
Currants	Fruittree leafroller	
Longanberries	Grape berry moth	
Raspberries	Gypsy moth	
Strawberries	Loopers	
	Obliquebanded leafroller	
Grapes:	Grape berry moth	Omnivorous leafroller
	Cherry fruitworm	Orange tortrix
	Grape leafroller	Saltmarsh caterpillar
	Grapeleaf skeletonizer	Yellowstriped armyworm
	Green fruitworm	
Tropical and Other Fruit:	Amorbia	Omnivorous leafroller
Avocados	Loopers	Omnivorous looper
	Orange tortrix	Spanworm
Bananas	Banana skipper	
Kiwi	Omnivorous leafroller	
Persimmons	Citrus cutworm	Omnivorous leafroller
Pomegranate	Fall webworm	Redhumped caterpillar
	Filbert webworm	Tent caterpillar
Pineapple	Gummosos-Batrachedra comosae	
	Thecla-Thecla basioides	
Tropical fruits	Hornworms	Loopers
	Leafrollers	Omnivorous leafroller

Rate/Acre: 0.5 - 2.0 pounds

V. FIELD CROPS

Crop such as:	Insect Pest	
Canola/ Rape Seed	Armyworm	
Evening Primrose	Diamondback moth	
Meadow foam	Imported cabbageworm	
	Loopers	
Corn (Field, Sweet, Popcorn, Seed)	Armyworm	
	European corn borer	
	Southwestern corn borer	

V. FIELD CROPS, continued

Crop such as:	Insect Pest	
Cotton*	Beet armyworm	Saltmarsh caterpillar
	Bollworm	Soybean looper
	Cabbage looper	Tobacco budworm
	Cotton leaf perforator	Yellowstriped armyworm
Hops	Armyworm	Omnivorous leaftier
	Loopers	Spotted cutworm
	Obliquebanded leafroller	
Jojoba	Looper (<i>Anacamptodes sp.</i>)	
Peanuts	Green cloverworm	Podworm
	Loopers	Velvetbean caterpillar
Rice	Armyworm	Saltmarsh caterpillar
	Green cloverworm	Velvetbean caterpillar
	Loopers	
Safflower	Armyworm	Saltmarsh caterpillar
	Loopers	
Small Grains (Barley, Oats, Rye, Wheat, etc.)	Armyworm	
	Loopers	
Sorghum	European corn borer	Saltmarsh caterpillar
	Headworm	Velvetbean caterpillar
Soybeans	Green cloverworm	Soybean looper
	Podworm	Velvetbean caterpillar
Sunflowers	Banded sunflower moth	Loopers
	Beet armyworm	Sunflower moth
	Headmoth	
Tobacco	Tobacco budworm	
	Tobacco hornworm	
	Loopers	

Rate/Acre: 0.5 - 2.0 pounds

* Use CRYMAX® at 0.25 lb/acre to control light to moderate populations of newly hatched tobacco budworm and bollworm in integrated pest management programs. Repeat treatments at four to five day intervals or as long as necessary until results are acceptable. Ovicides or synthetic pyrethroids can be combined with CRYMAX® in accordance with the more restrictive of label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

VI. COMMERCIAL FLOWERS & ORNAMENTAL PLANTS

Crop such as:	Insect Pest	
Bedding plants	Armyworm	Oleander moth
Flowers	Azalea moth	Omnivorous leafroller
(Greenhouse and Field)	Beet armyworm	Omnivorous looper
Greenhouse	Diamondback moth	Tobacco budworm
Ornamentals	Ello moth (hornworm)	
Greenhouse	Florida fern caterpillar	
Vegetables	lo moth	
Container Stock	Loopers	

Rate/Acre: 0.5 - 2.0 pounds

8098

VII. FOREST, SHADE TREE & NURSERY STOCK

Crop such as:	Insect Pest	
Forest	Bagworm	Pine butterfly
Shade trees	Blackheaded budworm	Redhumped caterpillar
Nursery trees	Browntail moth	Saddleback caterpillar
	California oakworm	Saddle prominent caterpillar
	Douglas fir tussock moth	caterpillar
	Elm spanworm	Spring and Fall cankerworm
	Fall webworm	caterpillar
	Fruittree leafroller	Spruce budworm
	Greenstriped mapleworm	Tent caterpillar
	Gypsy moth	Tortix
	Jack pine budworm	Western tussock moth
	Mimosa webworm	

Rate/Acre: 0.5 - 2.0 pounds

VIII. TURF

Crop such as:	Insect Pest	
Turf	Armyworm	
	Sod webworm	
	Tropical sod webworm	

Rate/Acre: 0.5 - 2.0 pounds

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place inaccessible to children.


Pesticide Disposal: Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Certis USA, L.L.C. warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the insect problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF THE FITNESS OR MERCHANTABILITY IS MADE.

CRYMAX® is a trademark of Certis USA, L.L.C.
U.S. Patent No. 5441884, 5650308, 5776449, 5843744
Copyright © 2001 Certis USA



082907
101209