



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

70051-121

Date of Issuance:

9/6/2017

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

SoilGard WDG

Name and Address of Registrant (include ZIP Code):

Certis USA LLC
 9145 Guilford Road, Suite 175
 Columbia, MD 21046

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.

Signature of Approving Official:

Jeannine Kausch, Product Manager 92
 Microbial Pesticides Branch
 Biopesticides and Pollution
 Prevention Division (7511P)
 Office of Pesticide Programs

Date:

9/6/2017

2. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 09/06/2017

If you have any questions, please contact Nicola Steinmetz by phone at (703) 347-8567 or via email at steinmetz.nicola@epa.gov.

Sincerely,



Jeannine Kausch, Product Manager 92
Microbial Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

SoilGard® WDG

MICROBIAL FUNGICIDE

{Alternate Brand Names: None}

ACTIVE INGREDIENT:

Gliocladium virens strain GL-2112%*

OTHER INGREDIENTS:88%

TOTAL:100%

*Contains $\geq 1 \times 10^7$ CFU/g of product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

MANUFACTURED BY:

Certis USA LLC
9145 Guilford Road, Suite 175
Columbia, MD 21046


EPA Reg. No. 70051-121
EPA Est. No. 70051-CA-001



Master Label Consisting of Sublabels

A. Commercial Use

B. Homeowner Use



Sublabel A: Commercial Use

SoilGard® WDG

MICROBIAL FUNGICIDE

{Alternate Brand Names: None}



FOR ORGANIC PRODUCTION



[placeholder pending OMRI approval]

This bag will treat up to _____ acres when used at the highest application rate.
{Note: Number of acres will be filled as applicable to size of retail packaging}

For the Control of "Damping-Off" and Root Rot Pathogens of [Ornamental and [Food Crop Plants] [and] [Tobacco] Grown in Greenhouses, Nurseries, and Interiorscapes and Outdoors on Agricultural and Ornamental Crops] {or} [Vegetable [and] [Tobacco] Plants Grown Indoors and Outdoors]

ACTIVE INGREDIENT:

Gliocladium virens strain GL-21 12%*

OTHER INGREDIENTS: 88%

TOTAL: 100%

*Contains $\geq 1 \times 10^7$ CFU/g of product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See [side][back][right][left][other] panel for [additional] [[precautionary] [and] [first aid] statements] [[use] directions [for use]]

[Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.]

MANUFACTURED BY:

Certis USA LLC
9145 Guilford Road, Suite 175
Columbia, MD 21046



EPA Reg. No. 70051-121

EPA Est. No. 70051-CA-001

Lot Number/Container Batch Code:
If unopened and not refrigerated: Not for sale or use after [insert date 6 months after production]

Item Code:

Package Code:

Net Weight:

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled. Avoid breathing dust or spray mist. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Mixer/loaders and applicators must wear:

- NIOSH-approved particulate respirator with any N, R, or P filter with NIOSH approval number prefix TC-84A (if tank mixing with any oil-based adjuvants, spreaders, or spreader/stickers, use only R or P filters); or
- NIOSH-approved powered air purifying respirator with an HE filter with NIOSH approval number prefix TC-21C.
- Repeated exposure to high concentrations of microbial proteins can cause allergic sensitization.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, wear: coveralls, chemical-resistant gloves (made of any waterproof material), and shoes plus socks.

EXCEPTION: If the product is soil injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses including Interiorscapes, and other non-agricultural uses, do not enter treated areas without protective clothing until soil-incorporation and potting is complete.

GENERAL INFORMATION

Gliocladium virens, the active component in SOILGARD® WDG, is a soil fungus that is antagonistic to plant pathogenic fungi such as [*Pythium*], [*Rhizoctonia*], [*Sclerotinia* spp.], [*Phytophthora capsici*], [*Sclerotium*], [and] [*Fusarium*] [root] [and] [crown] rots]], thereby aiding in control of these "damping off" and root rot pathogens. SOILGARD®

WDG is a granular formulation containing a high concentration of *G. virens* spores suitable for incorporation into soil or soilless media. SOILGARD® WDG acts as a preventative and will protect non-infected plants. It may not, however, cure already diseased plants.

PRECAUTIONS

Where possible, allow the treated soil or media to incubate for 1 day prior to planting to achieve best results. Do not use other soil fungicides with SOILGARD® WDG at time of incorporation. Later addition of fungicides is acceptable but usually unnecessary. Best results are achieved if the contents of an open package are used within 30 days. Be careful to seal out moisture from the unused material by closing the inner bag tightly.

SOILGARD® WDG can be mixed with water for application as a soil drench, as a root dip, or through irrigation systems. The dry granules can also be mixed directly into soil or growing media. Use lower rates in sandy soils, warm soils (60°F or higher temperature at 4-inch depth), and as a preventative against low to moderate disease pressure. Use higher rates in heavy soils (high clay and/or organic matter), low temperatures (less than 60°F at 4-inch depth), as a preventative against heavy disease pressure, or as a curative for low to moderate disease pressure early in the growing season. The soil or growing medium should be moist but not wet at the time of application; the application water may be of sufficient volume to provide the necessary moisture unless the soil is extremely dry or difficult to wet. Do not apply SOILGARD® WDG to very dry, compacted, frozen, or water-logged soils, and do not over-water or allow soil to become saturated for extended periods (>3 hours) after application.

COMPATIBILITY WITH PESTICIDES, FERTILIZERS, AND ADJUVANTS

Do not apply chemical fungicides to soil or plant media at the same time as SOILGARD® WDG. In most cases, additional soil-applied fungicides should not be required to control seedling diseases after proper application of SOILGARD® WDG. If additional fungicide applications are desired, wait a minimum of 2 weeks after applying SOILGARD® WDG. The effectiveness of SOILGARD® WDG may be decreased if soil fungicides are applied to the same area before it has become fully established in the root zone.

SOILGARD® WDG can be applied after soil fumigation, solarization, or steam sterilization to prevent or delay reinfestation by pathogenic fungi. Do not apply SOILGARD® WDG to fumigated soil until fumigants have dissipated to levels safe for planting the crop. Do not apply SOILGARD® WDG to steam-sterilized soil or other planting media until it has cooled to a temperature of 100°F or lower.

Do not combine SOILGARD® WDG with other pesticides, surfactants, or fertilizers unless prior experience has shown the combination to be physically compatible, effective, and non-injurious under the conditions of use. SOILGARD® WDG has not been fully evaluated for compatibility with all adjuvants or surfactants. If a mixture with adjuvants or surfactants is planned, first conduct a compatibility test (“jar test”) by mixing propor-

tional quantities of SOILGARD® WDG and these products in a small volume of water.

FOR INCORPORATION INTO GROWING MEDIA [for control of [Pythium], [Rhizoctonia], [Sclerotinia spp.], [Phytophthora capsici], [Sclerotium], [and] [[Fusarium] [root] [and] [crown] [rots]]]

(Ornamentals, Tobacco, and Food Crop Plants Grown Indoors and Outdoors)

For incorporation into potting medium in greenhouse flats, plug trays, or pots:

- Mix [4-6 [ounces] [oz]] {or} [¼-¾ [pound] [lb]]* of SOILGARD® WDG per cubic yard of potting soil or other plant-growing medium. Granules must be distributed uniformly throughout the medium.
- If the potting medium has been heat sterilized, allow it to cool to 100°F or lower before adding SOILGARD® WDG.
- Allow at least 24 hours after incorporation of SOILGARD® WDG before planting seeds, cuttings, or other non-rooted plant material into the treated medium. Plants with established roots can be transplanted into treated soil immediately after mixing with SOILGARD® WDG.
- Do not use SOILGARD® WDG in media totally devoid of organic matter (such as sand). *G. virens* requires organic matter to grow and work. If the only available organic substrate is the seed, bulb, or other plant material, *G. virens* may cause injury to it.

***NOTE:** The rates in the range provided may require the use of more than one package.

For incorporation into seedling and planting beds (indoors and outdoors):

- Thoroughly till the soil in the planting bed prior to addition of SOILGARD® WDG.
- Incorporate SOILGARD® WDG granules uniformly within the plant root zone at the rate of [4-6 [ounces][oz]] {or} [¼-¾ [pound][lb]]* per cubic yard of soil. The table below can be used to determine how much SOILGARD® WDG to use depending on planting or rooting depth and area to be treated.
- SOILGARD® WDG is unlikely to completely eliminate disease from soil already severely infested with pathogenic fungi. For best results, use SOILGARD® WDG in the greenhouse at the time of planting, so that it has already colonized the root ball of the transplant.

Planting depth (inches)	Ounces* per 100 square feet	Square feet per ounce	Pounds* per 100 square feet	Square feet per pound
2	2-4	27-40.5	0.125-0.25	432-648
4	5-7	13.5-20.25	0.3-0.4	216-324
6	7-11	9-13.5	0.4-0.7	144-216
8	10-15	6.75-10.125	0.6-0.9	108-162
10	12-19	5.375-8.125	0.75-1.2	86-130
12	15-22	4.5-6.75	0.9-1.4	72-108

***NOTE:** The rates in the range provided may require the use of more than one package.

FOR DRENCH OR SPRAY APPLICATION TO SOIL [for control of [*Pythium*], [*Rhizoctonia*], [*Sclerotinia spp.*], [*Phytophthora capsici*], [*Sclerotium*], [and] [*Fusarium*] [root] [and] [crown] [rots]]]

Mix SOILGARD® WDG in [a minimum of 2 gallons of] water [per pound of product] and apply as a drench or coarse spray to soil or other plant growth media. Stir or agitate thoroughly and maintain agitation during application to keep SOILGARD® WDG suspended uniformly. The total amount of water required will be determined by the area to be covered. Apply SOILGARD® WDG immediately after mixing with water; do not allow the mixture to stand overnight or for prolonged periods. The finished mix can be sprayed using low-pressure watering nozzles on a spray boom or spray wand, drenched with a watering can, injected into the soil with a water wheel or shank injector, or applied with other high-volume delivery devices. If applying with spray equipment, use flood jet or wide orifice flat fan nozzles with screens and swirl plates removed. If desired, SOILGARD® WDG can be mixed with water and passed through a 50 mesh screen for spraying through standard orifice tee jet or flat fan nozzles.

Refer to the table below for specific rates and use instructions for different applications.

Application rates and additional instructions for drench or spray application of SOILGARD® WDG to soil or other plant media.

CROP or USE SITE		RATE*	ADDITIONAL INSTRUCTIONS
Transplanted crops (such as fruiting and leafy vegetables, melons, strawberries, tobacco, field grown ornamental plants, and fruit/nut trees and vines)	Before transplanting	1/8 - 1/2 pound per 100 gallons of water	Apply 4 fl. oz. of finished drench per plant in flats (unit trays) or pots within 1 week before transplanting into the field (during the hardening-off phase). For densely seeded trays or flats, apply 100 gallons of finished drench per 800 square feet. SOILGARD® WDG can also be applied in furrow immediately before transplanting as described for seed furrows in the next section.
	At transplanting	1/8 - 1/2 pound per 100 gallons of water	Apply 4 – 8 fl. oz. of finished drench (depending on the size of the root ball) in each transplant hole using a water wheel/shank injector, watering can, or low pressure sprayer with flood/drench nozzles.
	After transplanting	1/2 - 2 pounds per acre	Repeat (maintenance) applications can be made as needed through directed spray or drench at the base of each plant, or through the irrigation system (see “For Application Through Irrigation Systems” below).

{ } Symbolizes explanatory text to the reviewer
 [] Indicates optional text

CROP or USE SITE	RATE*	ADDITIONAL INSTRUCTIONS
Seed furrows or planting beds (such as potatoes and other tubers, bulbs, tobacco, and direct-seeded crops)	½ - 2 pounds per acre	Apply in 50 – 100 gallons of water per acre as banded drench in-furrow at or immediately before planting, and before covering seed or seed pieces with soil. Apply a band of sufficient width to cover the seed furrow and initial rooting zone (usually 4-8 inches, centered on the furrow). Drench can also be applied as a seed line treatment using shank or other injection equipment in sufficient water to treat both the seed furrow and initial rooting zone.
Ornamental plant and flower beds (indoor and outdoor)	½ - 1 ounce per 100 gallons of water	Thoroughly till the soil in the bed prior to application. Before planting, apply the finished drench at 100 gallons per 800 square feet (if 4" deep or less) or 100 gallons per 400 square feet (more than 4" deep). Do not water the treated area for 12 hours after application. Allow at least 24 hours between drench application and planting of seeds or non-rooted cuttings. Plants with established roots can be transplanted immediately into treated beds.
Turfgrass	½ - 1 ounce per 1,000 square feet	Compacted soil in high-traffic areas should be aerated before applying SOILGARD® WDG. For best results, turf should be lightly pre-irrigated immediately before application to wet the foliage. Broadcast as a coarse spray in 1 – 2 gallons of water per 1,000 square feet. Irrigate the treated area with ¼ – ½ inch of water via overhead sprinklers immediately after application to move the product into the soil profile and maintain moisture. Repeat application every 1 – 3 weeks as necessary to maintain disease control.
Greenhouse flats, plug trays, or pots (prior to planting)	½ - 1 ounce per 100 gallons of water	Apply the finished drench at 50 – 100 gallons per 800 square feet (if 4" deep or less) or 100 gallons per 400 square feet (more than 4" deep). Do not water the treated area for 12 hours after application. Allow at least 24 hours between drench application and planting of seeds or non-rooted cuttings, or transplanting seedlings without true leaves. Plants with true leaves and established roots can be transplanted immediately into treated pots.
Potted or container plants (including tobacco) with established roots (greenhouse, nurseries, interior-scapes)	½ - 2 ounces per 100 gallons of water	Apply the finished drench at the base of the plant at a rate of 4 fl. oz. per pot for shallow pots (up to 4 inches deep) or plants up to 6 inches tall. For deeper pots and/or plants over 6 inches tall, apply 8 to 16 fl. oz. per pot. May be repeated every 1 – 4 weeks as needed.

{ } Symbolizes explanatory text to the reviewer

[] Indicates optional text

***NOTE:** The rates in the range provided may require the use of more than one package.

FOR APPLICATION THROUGH IRRIGATION SYSTEMS [for control of [*Pythium*], [*Rhizoctonia*], [*Sclerotinia spp.*], [*Phytophthora capsici*], [*Sclerotium*], [and] [*Fusarium*] [root] [and] [crown] [rots]]]

SOILGARD® WDG can be applied only to the base of the plant through ground-directed surface or subsurface drip (trickle) or micro-irrigation, or through flood, furrow, or border irrigation.

Mix SOILGARD® WDG into a large volume (50 gallons or more) of water under continuous agitation and inject through a filter or screen (50 mesh) into the irrigation lines. [Alternatively, first suspend the product in a smaller volume of water (a minimum of 2 gallons per pound of SOILGARD® WDG) with thorough mixing, and then pour this suspension through a filter or screen (50 mesh) into the final volume of water for injection into the irrigation system.] Apply SOILGARD® WDG immediately after mixing with water; do not allow the mixture to stand overnight or for prolonged periods. When using large pressurized systems, inject SOILGARD® WDG after the system has been fully charged with water and over sufficient time (typically 30 – 60 minutes, depending on the system) to allow uniform distribution of the product. After application, flush the system with water for an additional 10 – 20 minutes to avoid fouling of irrigation lines due to fungal growth.

Apply SOILGARD® WDG through irrigation systems up to 1 week before or immediately after planting or transplanting. Apply at ½ - 2* pounds per acre (0.2 – 1* ounces per 1,000 square feet). Refer to the Chemigation Instructions below for additional use instructions for application through irrigation systems. Do not apply SOILGARD® WDG through any irrigation system unless allowed in the Chemigation Instructions. Application can be repeated every 1 – 4 weeks if necessary due to disease pressure.

***NOTE:** The rates in the range provided may require the use of more than one package.

FOR ROOT DIP APPLICATION OR ROOT COATING [for control of [*Pythium*], [*Rhizoctonia*], [*Sclerotinia spp.*], [*Phytophthora capsici*], [*Sclerotium*], [and] [*Fusarium*] [root] [and] [crown] [rots]]]

SOILGARD® WDG can be applied as a dip or coating for bare-root plantings of crops (such as horseradish, strawberries or caneberries, tobacco, ornamental shrubs, fruit trees) or to rooted cuttings of poinsettias and other ornamental plants. Dip or immerse roots in a suspension containing [4 – 8 [ounces][oz]] {or} [¼ - ½ [pound][lb]]* of SOILGARD® WDG per gallon of water immediately before planting. Roots or planting sets can also be coated with SOILGARD® WDG by shaking them in a sealed plastic bag containing the suspension of SOILGARD® WDG in water, or by moistening them with water and shaking in a sealed plastic bag containing dry granules. Do not apply SOILGARD® WDG directly to fresh (non-rooted) cuttings.

***NOTE:** The rates in the range provided may require the use of more than one package.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: SOILGARD® WDG is packaged in an airtight, foil-lined, resealable bag designed to protect the fungus from exposure to moisture and oxygen, which can impact storage life. Store in a cool, dry place or under refrigeration. In an unopened bag, SOILGARD® WDG will retain its biological activity for up to one year at refrigerated temperature (39°F or 4°C) and up to 6 months at ambient temperature (77°F or 25°C). Long-term exposure to air and higher temperatures (>77°F) may accelerate loss of bio-activity. Exposure to temperatures above 100°F for extended periods can be detrimental to SOILGARD® WDG. An open bag of SOILGARD® WDG can be kept for up to 30 days if the bag is tightly resealed. Squeeze excess air from the bag before resealing.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of onsite or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available or dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CHEMIGATION INSTRUCTIONS

Precautions:

Apply this product through ground-directed, pressurized irrigation systems such as surface or subsurface drip (trickle) or micro-irrigation (including micro-jets, micro-sprayers, spaghetti tubes, or individual tubes), or through flood, furrow, or border irrigation. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Public water system chemigation:

“Public water system” means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

1. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
5. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials

that are compatible with pesticides and capable of being fitted with a system interlock.

6. Do not apply when wind speed favors drift beyond the area intended for treatment.
7. Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and injector system and flush with clean water before use. Failure to provide a clean tank, free of scale or residues, may reduce effectiveness of this product.
8. Dilute the product in water following the label mixing directions. It may be pre-mixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip. Maintain continuous application in sufficient water to apply the specified rate evenly to the entire treated area.

Drip (trickle) and micro-irrigation chemigation:

1. The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Dilute the product in water following the label mixing directions. It may be pre-mixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff or excessive drip from treated rows, beds, or pots. Maintain continuous application in sufficient water to apply the specified rate evenly to the entire treated area.



Flood, furrow, or border chemigation:


1. Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity such as a drop structure or weir box to decrease potential of water source contamination from the backflow if water flow stops.
2. Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:
 - a. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
 - b. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
 - c. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
 - d. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
 - e. The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
 - f. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
3. Dilute the product in water following the label mixing directions. It may be pre-mixed in a supply tank with water, fertilizer, or other appropriate tank-mixed agricultural chemicals. Agitation is necessary. Apply to moderately moist soils. Use volumes that thoroughly wet the soil but that do not cause significant runoff. Maintain continuous application in sufficient water to apply the specified rate evenly to the entire treated area.

WARRANTY

Certis USA LLC warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the crop, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.

{Optional statements}

- Optional Graphics
 - Graphic depictions of diseases may appear with or without a descriptor listing the name of the disease.
 - Front Panel Bursts
 -  FOR ORGANIC PRODUCTION
 -  {placeholder pending OMRI approval}
 - This bag will treat up to _____ acres when used at the highest application rate. {Note: Number of acres will be filled as applicable to size of retail packaging}
 - For the Control of "Damping-Off" and Root Rot Pathogens of [Ornamental and [Food Crop Plants] [and] [Tobacco] Grown in Greenhouses, Nurseries, and Interiorscapes and Outdoors on Agricultural and Ornamental Crops] {or} [Vegetable [and] [Tobacco] Plants Grown Indoors and Outdoors]
- Referral Statements
 - See [side][back][right][left][other] panel for [additional] [[precautionary] [and] [first aid] statements] [[use] directions [for use]]
 - See [attached] [booklet] [book] for [additional] [[precautionary] [and] [first aid] statements] [[use] directions [for use]]
 - Peel [back] [book] [here] [to open] [▶] [→] {may be accompanied by an arrow or other indicator}
 - Peel here [for [additional] [more] [complete] label instructions]. {may be accompanied by an arrow or other graphic indicating location of peel back corner}

- Hotline Number Text {may be used in place of or in addition to the current “Hot Line Number 1-800-255-3924”}
 - In the event of a medical or chemical emergency, [] contact North America 1-800-255-3924 or Worldwide Intl. +01-813-248-0585.
 - In case of emergency [endangering health or the environment] [involving this product], call [INFOTRAC] [other hotline] at [phone number].
- PRN 97-4 Access Numbers/Website References
 - Visit us at: [company website address]
 - [Product] Questions[?] [company phone number]
 - Comments[?]: [company phone number and/or website address]
- Other Miscellaneous Items
 - Barcodes/SKUs
 - Label revision numbers
 - Allow the insertion of “this product” in place of the product name everywhere it appears in the directions for use.
 - Allow the use of equivalent units of measures (e.g., “1½ “ in place of “1.5”) and appropriate equivalent abbreviations (e.g., “2-4 oz per 1,000 sq ft” or “2-4 oz/1,000 ft²” or similar in place of “2-4 ounces per 1,000 square feet”) throughout the directions for use.

{End Commercial Use Directions}

Sublabel B: Homeowner Use

SoilGard® WDG

MICROBIAL FUNGICIDE



FOR ORGANIC GARDENING



[Placeholder pending OMRI approval]

This bag will treat up to _____ [x inch pots] {and/or} [square feet] when used at the highest application rate. {Note: Number of pots and/or square feet will be filled as applicable to size of retail packaging}

For the Control of "Damping-Off" and Root Rot Pathogens of Ornamental and Food Plants {and/or} [Ornamental Flowering Tobacco] Grown in Homes, Home Greenhouses, and Outdoors in Home Gardens

ACTIVE INGREDIENT:

Gliocladium virens strain GL-21 12%*

OTHER INGREDIENTS: 88%

TOTAL: 100%

*Contains $\geq 1 \times 10^7$ CFU/g of product.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See [side][back][right][left][other] panel for [additional] [[precautionary] [and] [first aid] statements] [[use] directions [for use]]

[Refer to inside of label booklet for additional precautionary information and Directions for Use including First Aid and Storage and Disposal.]

MANUFACTURED BY:

Certis USA LLC
9145 Guilford Road, Suite 175
Columbia, MD 21046



EPA Reg. No. 70051-121
EPA Est. No. 70051-CA-001
Lot Number/Container Batch Code:
If unopened and not refrigerated: Not for sale or use after [insert date 6 months after production]
Item Code:
Package Code:
Net Weight:

FIRST AID

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Hot Line Number: 1-800-255-3924.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if inhaled. Avoid breathing dust or spray mist. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

For Incorporation Into Growing Media [for control of [*Pythium*], [*Rhizoctonia*], [*Sclerotinia spp.*], [*Phytophthora capsici*], [*Sclerotium*], [and] [*Fusarium*] [root] [and] [crown] [rots]]]

(Ornamentals, Ornamental Flowering Tobacco, and Food Plants Grown Indoors and Outdoors).

{Directions for 0.07 oz. package} Thoroughly blend SOILGARD® WDG with growing media. One 0.07 oz. packet (2 grams) will treat eight 8-inch pots. To apply SOILGARD® WDG, fill a clean 8-inch pot 3/4 full with potting soil. Spread the contents of the SOILGARD® WDG packet onto the soil and mix thoroughly with a small spade. It is important that SOILGARD® WDG be uniformly mixed into the media to provide the best performance possible. Transplants or plants for repotting can then be placed into the pot immediately. Wait at least 24 hours before planting seeds, bulbs or cuttings without roots. Moisten the media with water and keep moist (but not wet) for the next three days. The protective fungus will spread throughout the root system of the plant protecting it from disease organisms.

{Directions for 0.54 oz. package} Thoroughly blend SOILGARD® WDG with growing media. One 0.54 oz. packet (15 grams) will treat 4 cubic feet of media, enough to fill 72

6-inch pots. To apply SOILGARD® WDG, empty a 20 lb. bag of potting soil into a clean 5-gallon bucket. Spread SOILGARD® WDG onto the soil in the bucket and mix thoroughly with a spade or shovel. Alternatively, a corner can be cut from the potting soil bag and SOILGARD® WDG can be incorporated by shaking, stirring, or kneading the bag. It is important that SOILGARD® WDG be uniformly mixed into the media to provide the best performance possible. Soil can then be placed in pots. Moisten the media with water and keep moist (but not wet) for the next three days. Transplants or plants for repotting can be placed into pots immediately, but wait at least 24 hours to plant seeds, bulbs or non-rooted cuttings. The protective fungus will spread throughout the root system of the plant protecting it from disease organisms.

For Incorporation into Garden Rows [for control of [Pythium], [Rhizoctonia], [Sclerotinia spp.], [Phytophthora capsici], [Sclerotium], [and] [[Fusarium] [root] [and] [crown] [rots]]]

SOILGARD® WDG can be used to control soil-borne pathogens [including *Rhizoctonia*, *Pythium*, *Sclerotinia*, *Phytophthora*, *Sclerotium*, and *Fusarium*] that might attack plants grown outside in the home garden. Disease can be prevented by applying SOILGARD® WDG in the garden when plants are seeded or transplants are moved into the outside home garden. Disease prevention will exist for up to two months after application, depending on the disease pressure present. Application rates: apply and incorporate SOILGARD® WDG at the rate of 2 – 4* [ounces] [oz] per 100 [square feet] [sq ft] [ft²] (200 ft. x 6 inch row). Plants with established roots can be transplanted into treated soil immediately but allow at least 24 hours after incorporation before planting seeds, non-rooted cuttings, or bulbs.

***NOTE:** The rates in the range provided may require the use of more than one package.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store unopened product in a dry, cool place or in refrigerated conditions and away from foodstuffs. Seal out moisture from unused product in an opened package by closing the bag tightly after squeezing out excess air. Keep unused product in an opened package cool and away from foodstuffs in the original package and use within 30 days after opening.

Container Handling

Nonrefillable container. Do not reuse or refill this container.



If empty: Place in trash or offer for recycling if available.


If partly filled: Call your local solid waste agency [or 1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY

Certis USA LLC warrants that the material contained herein conforms to the description on the label and is reasonably fit for the purposes referred to in the directions for use. Timing and method of application, weather, watering practices, nature of soil, the disease problem, condition of the plant, incompatibility with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. To the extent consistent with applicable law, buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herein. NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY IS MADE.

{Optional statements}

- Optional Graphics
 - Graphic depictions of diseases may appear with or without a descriptor listing the name of the disease.
- Front Panel Bursts
 -  FOR ORGANIC GARDENING
 -  {placeholder pending OMRI approval}
 - This bag will treat up to _____ [x inch pots] {and/or} [square feet] when used at the highest application rate. {Note: Number of pots and/or square feet will be filled as applicable to size of retail packaging}
 - For the Control of "Damping-Off" and Root Rot Pathogens of Ornamental and Food Plants {and/or} [Ornamental Flowering Tobacco] Grown in Homes, Home Greenhouses, and Outdoors in Home Gardens
- Referral Statements
 - See [side][back][right][left][other] panel for [additional] [[precautionary] [and] [first aid] statements] [[use] directions [for use]]
 - See [attached] [booklet] [book] for [additional] [[precautionary] [and] [first aid] statements] [[use] directions [for use]]
 - Peel [back] [book] [here] [to open] [▶] [→] {may be accompanied by an arrow or other indicator}
 - Peel here [for [additional] [more] [complete] label instructions]. {may be accompanied by an arrow or other graphic indicating location of peel back corner}
- Hotline Number Text {may be used in place of or in addition to the current "Hot Line Number 1-800-255-3924"}

- In the event of a medical or chemical emergency, [] contact North America 1-800-255-3924 or Worldwide Intl. +01-813-248-0585.
- In case of emergency [endangering health or the environment] [involving this product], call [INFOTRAC] [other hotline] at [phone number].
- PRN 97-4 Access Numbers/Website References
 - Visit us at: [company website address]
 - [Product] Questions[?] [company phone number]
 - Comments[?]: [company phone number and/or website address]
- Other Miscellaneous Items
 - Barcodes/SKUs
 - Label revision numbers
 - Allow the insertion of “this product” in place of the product name everywhere it appears in the directions for use.
 - Allow the use of equivalent units of measures (e.g., “1½ “ in place of “1.5”) and appropriate equivalent abbreviations (e.g., “2-4 oz per 1,000 sq ft” or “2-4 oz/1,000 ft²” or similar in place of “2-4 ounces per 1,000 square feet”) throughout the directions for use.

{End Homeowner Use Directions}