



070060-00008-051399

Printed: 09:38:57 Thursday, 24 Jun, 1999 # 11 / 3411

Systems Integration Group, Inc.

PM 91

70060-8

5/13/99

1 of 16



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Biopesticides and Pollution Prevention Division (7501W)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
70060-8

Date of Issuance:
5/13/99

Term of Issuance:
Unconditional

Name of Pesticide Product:
M-98-A05 Crop Protectant

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

Engelhard Corporation
101 Wood Avenue
P.O. Box 770
Iselin, NJ 08830-0770

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others

This product is unconditionally registered in accordance with FIFRA sec. 3(c)(5) subject to the comments listed below:

1. Make the following label changes:
 - a. Add the phrase "EPA Registration No. 70060-8 to your label before you release the product for shipment
 - b. Add the appropriate Establishment Number to your label before you release the product for shipment
2. Submit five copies of the final printed labeling before you release this product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec.6(e). Your release for shipment of the product constitutes acceptance of these conditions.

Unconditional registration does not eliminate the need for continual reassessment of a pesticide. If EPA determines, at any time, that additional data are required to maintain in effect, an existing registration, the Agency will require submission of such data under Section 3(c)(2)(B) of FIFRA.

Signature of the Approving Official

2 of 16

M-98-A05 CROP PROTECTANT

M-98-A05 Crop Protectant Forms a Barrier Film, Which Acts as a Broad Spectrum Agricultural Crop Protectant for Controlling Damage from Various Insect, Mite, and Disease Pests, Growth Enhancer, and Protectant Against Sunburn and Heat stress

ACTIVE INGREDIENT:
 Kaolin:..... 95.0%
 Other INGREDIENTS:..... 5.0%
 Total: 100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION


STATEMENT OF PRACTICAL TREATMENT (FIRST AID)

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

Manufactured by:
ENGELHARD CORPORATION
 101 Wood Avenue
 Iselin, NJ 08830
 Emergency phone number: 1-800-424-9300

Net Weight 50 lbs.

EPA Reg. No. 70060-
 EPA Est. No. 70060-GA-1



PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. May cause irritation to the respiratory system. Avoid breathing dust or mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT

Applicators, pickers, and other handlers must wear:
Long sleeved shirt, long pants, socks and shoes, and a dust/mist-filtering respirator (MSHA/NIOSH approval number prefix TC-21C), or a NIOSH approved respirator with any N. R. P. or HE filter.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

AGRICULTURAL USE REQUIREMENTS

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, in forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. **The restricted-entry interval (REI) is 4 hours from the time of application. Do not allow worker entry into treated areas during the restricted-entry interval (REI).** Personal protection equipment required for early entry workers are long-sleeved shirts, long pants, and shoes plus socks.

I. GENERAL INFORMATION

M-98-A05 protectant forms a mineral-based particle film intended for protection of agricultural crops. When M-98-A05 is applied to plants, a dry white film results. The M-98-A05 barrier will help reduce the damage to plants and crops caused by certain insects, mites, disease, and environmental stresses such as solar effects. This product may be used as the primary component in a pest management system that utilizes supplementary compatible pesticides.

Plant Response Precautions:

Lack of adhesion or film formation on foliage or fruit may indicate spray incompatibility with the plant surface or previously applied materials.

Ensure that plant leaf and fruit surfaces are dry before applying or poor adhesion or film formation will occur.

Do not spray where a visible white film is undesirable or cannot be washed off.

a. Mix Instructions:

For Agitating Sprayer Tanks

For Concentrate Application - Use 50 lbs. of M-98-A05 per 100 gallons of water.

For Dilute Application - Use 50 lbs. of M-98-A05 per 200 gallons of water.

The following mixing sequence must be followed:

1. Slowly add M-98-A05 powder into the water in a recirculating sprayer tank. Add directly into the mix basket. If there is no mix basket, add powder very slowly to the recirculating water. Mix thoroughly.
2. Add Engelhard M-03 or Engelhard D-22 Spreader/Sticker per the use instructions on their labels.

If Organic growing practices are used, use Engelhard M-03 or O-23 Spreader/Sticker per use instructions on the label. (Check with your organic material regulating body to determine if Engelhard M-03 or O-23 spreader/sticker is approved).

3. Add tank mix pesticides, if any.
4. Continue agitation until all of the material is sprayed from the tank.
5. At the end of the application, spray until empty and flush system and nozzles with fresh water.

Properly dispose of rinse water.

For Non-agitating Sprayer Tanks, Such as Backpack Sprayers

The following mixing sequence must be followed:

1. Use M-98-A05 powder at a rate of ½ lb. of M-98-A05 powder per one gallon of water.
2. While stirring or swirling, slowly add M-98-A05 powder into ¼ to ½ of the water that will be used in the batch.
3. Add Engelhard M-03 or Engelhard D-22 Spreader/Sticker per the use instructions on their labels.

If Organic growing practices are used, use Engelhard M-03 or O-23 Spreader/Sticker spreader per use instructions on the label. (Check with your organic material regulating body to determine if Engelhard M-03 or O-23 Spreader/Sticker is approved).

4. Mix thoroughly by shaking the closed container vigorously for 30 seconds.
5. Add tank mix pesticides, if any.
6. Add the remainder of the batch water and shake for an additional 30 seconds.
7. Shake the sprayer occasionally during application
8. At the end of the application, spray until empty and flush system and nozzles with fresh water. Properly dispose of rinse water.

b. Compatibility:

M-98-A05 is not generally effected by most other insecticides, miticides, and fungicides. However, the user should test tank mixes before use. When mixing with other products, make up a small batch and observe slurry and film characteristics. Curdling, precipitation, lack of film formation, or changes in viscosity are signs of incompatibility. Do not tank mix with sulfur, copper, or bourdeaux mixture fungicides. **Always add tank mix pesticides after the M-98-A05 powder and spreader/sticker have been added.** Use of anti-foaming agents is not recommended.

Use of Engelhard-supplied spreader/stickers is essential. Use of other spreader/stickers may cause poor film formation, inadequate adhesion, settling, clogging, equipment damage or may make M-98-A05 difficult to remove from fruit surfaces. Only Engelhard M-03, D-22, and O-23 spreader/stickers are approved for use with M-98-A05.

Do not use additional spreader/stickers. Products that specify other spreader/stickers cannot be tank mixed with M-98-A05.

c. General Application Guidelines (see, also, specific crop use instructions):

Coverage: Use sufficient spray volume to obtain thorough coverage. For optimal performance, applications must coat all portions of plant that are to be protected, including both sides of the leaves. Applications can be made with any commercial air blast or high-pressure sprayer that provides enough air turbulence to coat both sides of the leaves, bark, and fruit. Tractor speed must not exceed 3 mph.

Plant Color Change: All plant surfaces must turn a hazy white color after drying. Additional treatments will turn the plant surfaces a deeper white. This is normal, and is indicative of appropriate film formation. Generally, two or more coatings are required for complete coverage and for establishment of the foundation of the protective barrier.

Foliage Dryness: Apply only to dry foliage. Do not apply to wet foliage or poor coverage and adhesion will result.

Droplet Size: Nozzles that produce a fine spray are recommended when M-98-A05 is used under normal temperature and humidity conditions. Under very hot and dry conditions, nozzles that produce a coarser spray and larger volumes may be necessary. Otherwise, the product may dry before it contacts the foliage and will result in poor film formation.

Spray Methods: Air blast or high-pressure sprayers are required. Observe specific crop label instructions for directions regarding spray volume. Calibrate spray equipment per equipment manufacturer to deliver the required volume. The flow rate of this product is similar to water. Use strainers in the spray system and behind each nozzle per normal practice.

Reapplication: Reapplication is generally required every 7-14 days when using M-98-A05 with Engelhard M-03 or O-23 Spreader/Sticker and every 10 to 17 days when using M-98-A05 with Engelhard D-22 Spreader/Sticker. Intervals can be widened later in the season when new growth diminishes.

When the dry foliage has lost its white appearance, reapplication is necessary. Heavy rainfall, new growth, and wind erosion will affect film quality. Reapply to re-establish coverage after heavy rain as soon as the foliage is dry.

Overhead Irrigation Considerations: Overhead irrigation will wash off the film and reduced efficacy will result. Overhead irrigation is not preferred, but, if used, should be used just prior to the next application to prevent premature wash-off. Do not apply M-98-A05 through sprinkler irrigation systems or by air.

d. General Insect and Mite Control:

M-98-A05 is a protectant/repellent that forms a film barrier to insects and mites. For optimum performance, begin application before a particular insect or mite outbreak occurs; thorough coverage must be in place before the pest outbreak. Thorough, uniform, and consistent coverage is essential for effective control.

e. Growth Enhancer, Sunburn and Heatstress Protectant:

When applied at recommended rates and frequencies, benefits such as increased plant vigor and improved yields may occur on many fruit and nut trees. Under high ambient temperatures, M-98-A05 reduces canopy temperature and, therefore, can help to reduce heat and water stress. When M-98-A05 is used, many pome and stone fruit varieties have shown improved fruit color, smoothness, and size with less russet, sunburn, and cracking.

For sunburn reduction, apply before conditions leading to sunburn occur. Apply concentrate spray at least every 7-14 days, to ensure complete coverage of fruit surfaces. Under windy conditions, particle film may be rubbed off by leaf movement making reapplication necessary.

If initiating sprays for sunburn reduction where there have been no prior sprays, thorough coverage of all fruit surfaces must be in place just prior to sunburn-causing conditions. To achieve optimal coverage, it is necessary to start the spray program with either two concentrate applications 5-7 days apart, or, one application with twice the concentrate use rate of per acre (for example, for nominal 12' trees, 100 lbs. per acre in 200 gallons of water).

f. Packing and Processing:

Washing is required unless films weather off before harvest. Most residues wash off with packing line brushing and forced water sprays. An approved washing detergent may be used in the packing line and/or wash tank. A pre-harvest washing trial is recommended to determine if a washing detergent is necessary. Waxing further improves fruit appearance.

g. Organic Growing Considerations:

1. Washing is usually necessary to remove film. Consult your organic materials regulating body for approved fruit cleaning products. Plain water brushing and/or hand wiping can remove film from firm smooth-skinned fruit such as apples, pears, and nectarines.
2. Tank mixes with pyrethrins, potassium salts of fatty acids, and rotenone are generally compatible but curdling, precipitation, uneven film formation, or changes in viscosity are signs of incompatibility. Test mix compatibility before use. If these conditions occur, do not tank mix M-98-A05.
3. Do not tank mix or over spray with sulfur, copper, or bourdeaux mixture fungicides.

4. Use Engelhard M-03 or O-23 Spreader/Sticker only. Check with your organic material regulating body to determine if M-03 or O-23 Spreader/Sticker is approved.

II. TREE FRUIT AND NUT CROPS

Apply sufficient spray volume to obtain thorough coverage.

The plant surface must turn a hazy white color after drying. Additional treatments will turn the plant surface a deeper white. This is normal, and is indicative of appropriate film formation. Generally, two or more coatings are required for complete coverage and for establishment of the foundation of the protective barrier.

Maintain film coverage to maximize potential solar protectant benefits. If rainy weather reduces coverage on leaves and fruit, reapply to reestablish coverage.

Estimate* rates based on the table below:

Concentrate Application

<u>Approx. Tree Height (ft)</u>	<u>Gallons of Mix per Acre</u>	<u>Lbs. of M-98-A05 per Acre</u>
6'	50	25
12'	100	50
18'	150	75
24'	200	100

Dilute Application**

<u>Approx. Tree Height (ft)</u>	<u>Gallons of Mix per Acre</u>	<u>Lbs. of M-98-A05 per Acre</u>
6'	100	25
12'	200	50
18'	300	75
24'	400	100

*Estimate rates are based on tree height only. Tree spacing, pruning practices, time of season, and cultivar type affect spray rates. High density or low-density foliage will require correspondingly higher or lower rates. A visual inspection of film deposition and completeness of coverage is crucial for fine-tuning spray amounts.

**Apply to near-drip. Do not apply to excessive drip.

Continue applications up to 10 days prior to harvest to maximize M-98-A05 horticultural benefits.

Fungicides, insecticides, miticides, and bactericides may be tank mixed or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

For heat stress relief for tree fruit and nut crops

Apply at 7 to 14 day intervals starting when temperatures are anticipated to reach 90 degrees F or when heat stress is known to occur. Do not spray during bloom. Resume treatments within three days of first petal fall. Application intervals may be widened to 14 to 21 days as new growth slows or when dry periods occur, being sure to maintain good film coverage to maximize potential benefits of M-98-A05.

Washing is recommended; most residues wash off with brushing and forced water sprays. An approved fruit cleaning detergent may be used in the packing line and/or wash tank. A pre-harvest washing trial is recommended to determine if a detergent is necessary. Waxing further improves fruit appearance.

Pears

For pear psylla control:

- Dilute applications are recommended. Adjust volume per tree height to insure thorough coverage without excessive drip.
- Apply every 7 –14 days from green cluster bud through popcorn. A spray at popcorn is recommended to cover the bloom period. Resume M-98-A05 application within three days of first petal fall, following a 10 to 14 day spray schedule under normal non-rainy conditions. Reapplication following heavy rain may be necessary.
- Application intervals may be widened to 14 to 21 days as new growth slows or when dry periods occur.

For suppression of twospotted spider mite, European red mite, pear rust mite, mealy bug, thrips, leafhoppers, green fruit worm, codling moth, California pear slug, Lygus bugs, obliquebanded leafroller, pandemus leafroller, tarnished plant bug, stink bugs, and fabraea leaf spot:

Spray per recommendations for psylla control or spray before expected infestation and continue at 7 to 14 day intervals. Do not widen spray intervals beyond 14 days to achieve season-long suppression. Do not spray during bloom. Resume treatments within three days of first petal fall. Supplemental pest control methods may be needed for full control; especially for codling moth.

Continue applications to maximize M-98-A05 horticultural benefits.

Fungicides, insecticides, miticides, and bactericides may be tank mixed, if compatible, or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no case apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

Apples

For control of over-wintering obliquebanded leafroller

- Apply two sprays 7 days apart from delayed dormant to pink.

For control of leafhoppers

- Apply every 7 to 14 days starting within three days of first petal fall.

For suppression of European red mite, apple rust mite, two-spotted mite, codling moth, plum curculio, Leafminers, apple sucker, Lygus bugs, pandemus leafroller, tarnished plant bug, stink bugs, apple maggot, thrips, green fruitworms, and aphids:

- Apply at 7 to 14 day intervals starting at delayed dormant. Do not spray during bloom. Resume treatments within three days of first petal fall applying at 7 to 14 day intervals. Application intervals may be widened to 14 to 21 days as new growth slows or when dry periods occur. Be sure to maintain a good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control; especially for codling moth.

M-98-A05 may improve fruit quality if applications resume at initial petal fall and are continued up until 10 days prior to harvest. When applied at recommended rates and frequencies, benefits such as increased plant vigor and improved yields may occur in certain apple cultivars. Under high ambient temperatures, M-98-A05 reduces canopy temperature and, therefore, can help to reduce heat and water stress. Many apple cultivars have shown improved fruit color, smoothness, and size with less russet, sunburn, and cracking when M-98-A05 is used.

For sunburn reduction, apply before conditions leading to sunburn occur. Apply concentrate spray at least every 7-14 days, or as needed to ensure complete coverage of fruit surfaces. Under windy conditions, particle film may be rubbed off by leaf movement making reapplication necessary.

If initiating sprays for sunburn reduction where there have been no prior sprays, thorough coverage of all fruit surfaces must be in place just prior to sunburn-causing conditions. To achieve optimal coverage, it is necessary to start the spray program with either two

concentrate applications 5-7 days apart, or, one application with twice the concentrate use rate of per acre (for example, for nominal 12' trees, 100 lbs. per acre in 200 gallons of water).

Fungicides, insecticides, miticides, and bactericides may be tank mixed, if compatible, or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

Stone Fruit – Apricots, cherries, nectarines, peaches, plums, and prunes

For control of Japanese beetle

- Apply one week prior to the expected infestation period and follow with three to four additional applications at 5 to 7 day intervals.

For suppression of European red mite, twospotted spider mite, peach silver mite, thrips, stink bugs, oriental fruit moth, tarnished plant bug, European earwig, June beetle, peach twig borer, naval orange worm, and plum curculio

- Apply at 7 to 14 day intervals starting at bud swell and ending at pink. Do not spray during bloom. Resume treatments within three days of first petal fall. Application timing may be widened when new growth slows or when dry period occur. Be sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

Continue applications to maximize M-98-A05 horticultural benefits.

Fungicides, insecticides, miticides, and bactericides may be tank mixed, if compatible, or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

Special Washing Considerations for Stone Fruit: Washing is required; especially for fuzzy peaches. Most residues wash off with brushing and forced water sprays. An approved fruit cleaning detergent may be used in the packing line and/or wash tank. Prior to brushing, a pre-soak in approved fruit cleaning detergent is usually needed for fuzzy peaches. A pre-harvest washing trial is recommended to determine if a detergent is necessary. Waxing further improves fruit appearance.

Citrus Fruits – Lemons, Limes, Grapefruit, and Oranges

For suppression of thrips and mites

- Apply at 7 to 14 day intervals starting within three days of first petal fall. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

Continue applications to maximize M-98-A05 horticultural benefits.

Fungicides, insecticides, miticides, and bactericides may be tank mixed, if compatible, or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

Nut Trees – Walnuts, Pecans, and Almonds

For suppression of mites

- Apply at 7 to 14 day intervals starting after nut set. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

Continue applications to maximize M-98-A05 horticultural benefits.

Fungicides, insecticides, miticides, and bactericides may be tank mixed or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

III. SMALL FRUIT CROPS

Apply sufficient spray volume to obtain thorough coverage.

Apply only to fruits to be used for processing.

Application of plain water via normal sprayer prior to harvest may help to reduce M-98-A05 residues. This may be particularly helpful for wine grapes.

Continue applications to maximize M-98-A05 horticultural benefits.

Fungicides, insecticides, miticides, and bactericides may be tank mixed, if compatible, or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

Blackberries, Raspberries, Dewberries, Boysenberries, Loganberries, and Blueberries

For suppression of blackberry psyllid, European raspberry aphid, Japanese beetle, leafhoppers, thrips, twospotted spider mite, and European red mite

- Apply 12.5 to 50 lbs. per acre at 7 to 14 day intervals starting after fruit set. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

Wine Grapes

For suppression of European red mite, two-spotted spider mite, omnivorous leafroller, grape leafroller, grape leaf skeletonizer, leafhoppers, Japanese beetle, June beetle, and thrips

- Apply at 12.5 to 37.5 lbs. per acre at 7 to 14 day intervals starting after fruit set. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

IV. FIELD AND VEGETABLE CROPS

Apply sufficient spray volume to obtain thorough coverage.

Fungicides, insecticides, miticides, and bactericides may be tank mixed, if compatible, or over-sprayed according to their product label use instructions. If tank mixing pesticides, follow the pre-harvest interval of those pesticides but in no cases apply tank mixes within 10 days prior to harvest. If using M-98-A05 alone, reapply as needed up to harvest.

For difficult to wash or unwashed crops, especially in low rainfall areas, it is recommended to stop spraying 30 days prior to harvest or when fruit are small. For crops where the produce is not directly sprayed by M-98-A05 (e.g. potato, shelled beans, and radishes), M-98-A05 may be applied up to the day of harvest. Cotton may be sprayed until 10 days before harvest.

Washing is recommended; most residues wash off with brushing and forced water sprays; an approved detergent may be used in the packing line and/or wash tank. A pre-harvest washing trial is recommended to determine if a detergent is necessary.

For heat stress relief and sunburn protection of field and vegetable crops

- Apply at 7 to 14 day intervals starting when heat stress or sunburn is known to occur. Application timing may be widened to 14 to 21 days as new growth slows or when dry periods occur, being sure to maintain good film coverage to maximize potential benefits of M-98-A05.

Beans

For suppression of alfalfa looper, aphids, Fall armyworm, flea beetle, Mexican bean beetle, European red mites, Japanese beetle, leafhoppers, Lygus bug, stink bugs, tarnished plant bug, three cornered alfalfa hopper, thrips, twospotted spider mite, velvetbean caterpillar

- Apply 6.25 to 12.5 lbs. per acre at 7 to 14 day intervals starting after fruit set. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

**Collards, Garden beet, Sugar beet, Horseradish, Radishes
Rutabagas, and Turnips**

For suppression of European red mites, flea beetle, harlequin bug, leafhoppers, tarnished plant bug, twospotted spider mite, imported cabbage worm

- Apply 6.25 to 12.5 lbs. per acre at 7 to 14 day intervals starting after fruit set. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

Cotton

For suppression of cotton fleahopper, European red mites, flea beetles, tarnished plant bug, thrips, twospotted spider mite, whiteflies

- Apply 6.25 to 25.0 lbs. per acre at 7 to 14 day intervals starting after fruit set. Application timing may be widened during dry periods to once every 14 to 21 days, being sure to maintain good film coverage to maximize potential benefits of M-98-A05. Supplemental pest control methods may be needed for full control.

Potato, Tomato, Eggplant, Pepper

For suppression of European red mites, Colorado potato beetle, flea beetles, lace bugs, leafhoppers, stink bugs, tarnished plant bug, tomato fruit worm, tomato pinworm, twospotted spider mite

- Apply 6.25 to 25 lbs. per acre at 7 to 14 day intervals starting one week before infestation is expected. Supplemental pest control methods may be needed for full control.

Onions

For suppression of onion thrips

- Apply 6.25 to 12.5 lbs. per acre at 7 to 14 day intervals starting one week before infestation is expected. Supplemental pest control methods may be needed for full control.

Cucurbits

For suppression of cucumber beetles

Apply 6.25 to 25 lbs. per acre at 7 to 14 day intervals starting one week before infestation is expected. Supplemental pest control methods may be needed for full control.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in a dry, sheltered location. Product is slippery when wet. In case of spill or leak, avoid breathing dust, clean up and dispose of wastes in compliance with applicable Federal, State, and local regulations.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to applicable Federal, State, and local procedures.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of the smoke.

CONDITIONS OF SALE, WARRANTY STATEMENT, DISCLAIMER

THE MANUFACTURER WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND THAT, SUBJECT TO THE INHERENT RISKS DISCUSSED BELOW, THIS PRODUCT IS REASONABLY FIT FOR THE PURPOSES SET FORTH IN THIS LABEL WHEN USED UNDER NORMAL CONDITIONS AND STRICTLY IN ACCORDANCE WITH THE DIRECTION, WARNINGS, AND OTHER STATEMENTS ON THIS LABEL. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS

INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER. ALL SUCH RISKS ARE ASSUMED BY BUYER/USER. THE DIRECTIONS, WARNINGS AND OTHER STATEMENTS ON THIS LABEL ARE BASED UPON REASONABLE EXPERTS' EVALUATIONS OF REASONABLE TESTS OF EFFECTIVENESS, AND UPON REPORTS OF FIELD EXPERIENCE. TESTS HAVE NOT BEEN MADE UNDER ALL CONDITIONS.

THE MANUFACTURER NEITHER INTENDS, NOR DOES IT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, EXPRESSED OR IMPLIED, AND IT EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

THIS WARRANTY DOES NOT EXTEND TO, AND THE BUYER SHALL BE SOLELY RESPONSIBLE FOR, ANY AND ALL LOSSES OR DAMAGE WHICH MAY RESULT FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL, DIRECTIONS, WARNINGS OR CAUTIONS.

BUYER'S EXCLUSIVE REMEDY AND MANUFACTURER'S OR SELLER'S EXCLUSIVE LIABILITY FOR ANY AND ALL CLAIMS, LOSSES, DAMAGES OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY, TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF, OR REFUND OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INDIRECT OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

ENGELHARD OFFERS THIS PRODUCT FOR SALE AT ALL TIMES SUBJECT TO THE FOREGOING.

M-98-A05 Crop Protectant
An HPF™ Particle Film Technology Product

© 1999 Engelhard Corporation
HPF is a trademark of Engelhard Corporation

M970A05Label5991