

PM 13

70126-1

5315817
161 5311196
181 1/5

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
70126-1

Date of Issuance:
MAR 20 1997

NOTICE OF PESTICIDE:
 Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance:
Conditional

Name of Pesticide Product:
Organic Resources
Multipurpose
Insecticide

Name and Address of Registrant (include ZIP Code):

Organic Resources, Inc.
8023 Vantage Drive, Suite 600
San Antonio, TX 78230

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A), provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label change:
 - a. Add the phrase "EPA Registration Number 70126-1" to the label.
 - b. Revise the rate application for fruit, nut, vine and speciality crops from 4 to 12 lbs per acre to 2 to 6 lbs per acre as it appears in the reference product labeling.

CONCURRENCES

SYMBOL	Signature of Approving Official:	Date:			
SURNAME					
DATE					

2/5

- c. Since this product can be used in food/feed areas of food handling establishments, delete the paragraph beginning with "DO NOT USE IN: edible product areas of food/feed handling establishments....".
- d. Since your product contains three active ingredients, you must identify Piperonyl Butoxide in the statement "Contains Natural Pyrethrin and Diatomaceous Earth".
- e. Delete the word "Natural" from the statement "Contains Natural Pyrethrin and Diatomaceous Earth" since statements or words relative to composition directly or indirectly implying safety of a product or pesticide is not acceptable.
- f. Based on the results of the product chemistry review, revise the percentage of silicon dioxide in the ingredients statement to be 82.9% and the total inert ingredients to be 15.9%.

3. Submit two copies of the revised final printed label for the record. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager 13
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

BEST AVAILABLE COPY

3/5

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Harmful if absorbed through the skin. Avoid contact with eyes, skin or clothing. Use adequate ventilation and avoid breathing dust.

Personal Protective Equipment:

Applicators and other handlers must wear long-sleeved shirt and long pants, waterproof gloves, shoes and socks and a dust-rated filtering respirator (NIOSH/NIOSH) approved number prefix: TX-21C. Follow manufacturer's instructions for cleaning and maintaining Personal Protective Equipment. If no instructions for washables are given, use detergent and hot water. Keep and wash Personal Protective Equipment separately from other laundry.

User Safety Recommendations

Use should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove Personal Protective Equipment immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothes.

Environmental Hazards:

This product is toxic to fish. Do not apply directly to any body of water. Do not apply when weather conditions favor drift from the treatment area. Do not contaminate water by disposing of equipment wash waters.

STORAGE AND DISPOSAL

Storage: Store in original container in a cool dry place, out of sunlight. Disposal: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse container. If possible, flush container with water and use the water in application. Wrap empty flushed container and discard in trash.

WARRANTY STATEMENT

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and conditions of application equipment will vary, seller makes no claims other than those on this label.

ORGANIC RESOURCES, Inc.
A Delaware Corporation
8023 Vantage Drive, Suite 600
San Antonio, TX 78290

EPA Reg. No. _____ EPA Est. No. _____ - TX-001

Batch No. _____ Lot _____
ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 20 1997

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

70126-1

MASTER LABEL
ORGANIC RESOURCES
MULTIPURPOSE INSECTICIDE
CONTAINS NATURAL PYRETHRIN
AND DIATOMACEOUS EARTH

FOR USE INDOORS AND OUTDOORS INCLUDING BUT NOT LIMITED TO: HOMES, ANIMAL QUARTERS, ON PETS, GRASSES, LAWNS & TURF (INCLUDING GOLF COURSES), GARDENS, VEGETABLE & GROWING CROPS (OUTDOORS & GREENHOUSES), ORNAMENTALS, SPECIALTY CROPS, FIELD GRAIN CROPS, FRUIT, NUT, VINE AND ORIENTAL VEGETABLE CROPS.

ACTIVE INGREDIENTS:

Pyrethrin	0.2%
Piperonyl Butoxide	1.0%
Silicon Dioxide (from Diatomaceous Earth)	98.0%
INERT INGREDIENTS	10.8%
Total	100.0%

*Equivalent to 0.8% (butylcarbamoyl(5-propyl)pyrethrin) ether and 0.2% related compounds.

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger.

If inhaled: Remove person to fresh air. Apply artificial respiration if indicated.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If in eyes: Flush eyes with plenty of water. If irritation persists, call a physician immediately.

SEE BACK PANEL FOR ADDITIONAL DIRECTIONS

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, are: Coveralls, waterproof gloves, shoes and socks. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 22 hours.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Keep unprotected persons, children and pets out of the treated area until sprays have dried.

GENERAL INFORMATION: Carefully follow the directions for each type of use. This material may be used on edible crops growing outdoors or in greenhouses, up to and including the day of harvest. Use rates will vary with insect populations and amount of foliage.

USE RATES FOR VEGETABLES, FIELD CROPS, ORNAMENTALS AND MISCELLANEOUS CROPS: Use 2 to 6 pounds of product per acre depending upon the insect population and amount of foliage. The material can be used either:

AS A DUST, the best results are obtained when the plants are wet or dew still remains on the leaf surface. It is suggested that the plants are misted prior to application to obtain the maximum benefit and use the least amount of product for control. Reapply only as necessary.

AS A WETTABLE POWDER, apply in sufficient water to obtain maximum coverage. Dilution of the material should be at the maximum rate of one half pound in one half gallon of water or at a minimum of one pound to 25 gallons of water (4 Teas. per gallon). It is best to keep the material agitated during application. Apply the material uniformly to both the upper and lower leaf surfaces. Reapply only as necessary. Rates will vary with insect populations and amount of foliage, but adequate control can be obtained at 2 to 6 pounds product per acre.

VEGETABLE CROPS:

FOR THE CONTROL OF INSECTS SUCH AS: Aphis, Aphids, Mexican Bean Beetle, 12 Spotted Cucumber Beetle, Army Worms, Beet Webworms, Blister Beetles, Cabbage Looper, Cabbage Worms, Caterpillars, Cutworms, Colorado Potato Beetle, Corn Earworm, Cucumber Beetle, Diamond Back Moth Larva, European Corn Borer, Fire Ants, Flea Beetle, Garden Fleahoppers, Harlequin Bug, Imported Cabbageworm, Japanese Beetles, Leafhoppers, Leaf Miners, Leaflice, Loopers, Lygus Bugs, Mites, Oblique-banded Leafhoppers, Plant Bugs, Stink Bugs, Squash Vine Borer, Thrips, Vegetable Weevils, Webworms and White Flies.

GROWING CROPS (Outdoors and Greenhouses):

Root and Tuber Vegetables: including but not limited to Artichoke, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke,

5/16

GROWING CROPS (Outdoors and Greenhouses):

Root and Tuber Vegetables: Including but not limited to Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichoke, Jerusalem Artichoke, Beets, Sugar Beets, Edible Burdock, Carrots, Cassava (Bitter & Sweet), Celery, Celery Root, Chervil (Turnip Root), Chicory, Chufa, Dasheen, Ginger, Ginseng, Horseradish, Laren, Parsley (Turnip Rooted), Parsnip, Potato, Radish, Japanese Radish, Rutabaga, Salsify, Black Salsify, Sweet Potato, Tanler, Tarrow Root, Turnip, Turnip, Yam, Bean Yam.

Leaves of Root and Tuber Vegetables: Including but not limited to Beet, Sugar Beet, Edible Burdock, Carrot, Cassava (Bitter & Sweet), Celery, Chervil, Chicory, Dasheen, Parsnip, Radish, Japanese Radish, Rutabaga, Black Salsify, Sweet Potato, Tanler, Turnip and Yam (True).

Bulb Vegetables: Including but not limited to Garlic, Leek, Onion (Bulb & Green), Shallot.

Leafy Vegetables: Including but not limited to Amaranth, Leafy Amaranth, Chinese Spinach, Tampala, Arrugula, Celery, Celtuce, Chervil, Cilantro, Corn Salad, Chrysanthemum (edible leaves), Chrysanthemum Garland, Cress (Garden), Upland Cress (Yellow Rocket Winter Cress), Dandelion, Dock, Endive, Fennel, Lettuce (Head and Leafy), Orach, Parsley, Pursians (Garden and Winter), Rhubarb, Spinach, Fine Spinach (Malabar Ceylon), Spinach (New Zealand), Swiss Chard.

Brassica (Cole) Leafy Vegetables: Including but not limited to Broccoli, Chinese Broccoli, Broccoli Raab, Brussel Sprouts, Cabbage, Chinese Cabbage (Bok Choy, Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Collards, Kale, Kohlrabi, Mustard Greens, Rape Greens.

Legume Vegetables (Succulent or Dried): Including but not limited to Adzuki Beans, Field Beans, French Beans, Kidney Beans, Lima Beans, Mch Beans, Mung Beans, Navy Beans, Pinto Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black Eyed Peas, Catjang, Chinese Longbeans, Cowpeas, Crowder Peas, Southern Peas, Yard-Longbeans, Broad Beans (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jack Beans (Sword Beans), Lablab (Hyacinth Beans), Lentils, Peas (Garden Peas, Field Peas, Sugar Peas), Pigeon Peas, Soybeans. Foliage of Legume Vegetables: Including but not limited to plant parts of any Legume vegetable included in the legume vegetable group that will be used as animal food including any variety of Beans, Field Peas, Soybeans.

Fruiting Vegetables: Including but not limited to Eggplant, Ground Cherry, Okra, Pepinos, Pepper (Bell Pepper, Chili Pepper, Cooking Pepper, Pimentos, Sweet Peppers), Tomatillo, Tomatoes.

Cucurbit Vegetables: Including but not limited to Balsam Pear (Bitter Melon), Chinese Waxgourd, Citron Melon, Cucumber, Gherkin, Edible Gourds, Melons Including Hybrids (Cantaloupe, Casaba, Crenshaw, Honeydew Melons, Honey Balls, Mango Melon, Muskmelon, Persian Melon), Pumpkin, Squash (Summer & Winter), Watermelon including hybrids.

Oriental Vegetables: Including but not limited to Japanese Artichoke, Chinese Broccoli (Gai Lon), Chinese Cabbage (Bok Choy Napa), Chinese Mustard Cabbage (Gai Choy), Cilantro, Dasheen, Ginger, Ginseng, Chinese Longbeans, Mung Beans, Citron Melon, Balsam Pear (Bitter Melon), Japanese Radish (Daikon), Chinese Spinach, Chinese Waxgourd.

FIELD GRAIN CROPS

For the control of insects such as Ants, Aphids, Armyworms, Chinch Bugs, Boll Weevil, Bollworm, Budworm, Caterpillars, Corn Earworms, Fire Ants, Fleabeetles, Fleahoppers, Flies, Hornworms, Looper, Lygus Bugs, Midge, Mites, Pear Cystitis, Pink Boll Worms, Thrips, White Flies.

Cereal Grains: Including but not limited to Barley, Buckwheat, Corn, Millet (Proso & Pearl), Oats, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticale, Wheat, Wild Rice.

Grass Forage, Fodder and Hay: Including but not limited to Bermudagrass, Bluegrass, Bromegrass, Fescue and any type grass, Gramineae Family (Green or Cured), Barley, Buckwheat, Corn, Millet (Proso & Pearl), Oats, Popcorn, Rice, Rye, Sorghum (Milo), Teosinte, Triticale, Wheat, Wild Rice, that will be fed to or grazed by livestock, all pasture, range grasses and grasses grown for hay or silage.

Non-Grass Animal Feeds: Including but not limited to Alfalfa, Velvet Bean, Clover, Kudzu, Leapedeza, Lupine, Sainfoin, Trefoil Vetch, Crown Vetch, Milk Vetch.

ORNAMENTALS:

For the control of insects such as: Ants, Aphids, Armyworms, Chinch Bugs, Caterpillars, Fire Ants, Flea Beetles, Fleahoppers, Flies, Fruit Flies, Japanese Beetles, Leafhoppers, Leafminers, Leafrollers, Loopers, Lygus Bugs, Mealy Bugs, Mites, Plant Bugs, Thrips, White Flies.

Ornamentals: Such as African Violets, Aster, Azalea, Begonia, Calceolaria, Calendula, Calla, Camellia, Carnation, Cineraria, Chrysanthemum, Cypress, Daffodil, Dahlia, Dogwood, Elm, Eucalyptus, Fern, Ficus, Geranium, Gladiolus, Gypsophila, Holly, Juniper, Lily, Marigold, Oak, Palm, Peony, Petunia, Philodendron, Pine, Roses, Snapdragon, Sweetpeas, Tulips, Viburnum, Wandering Jew, Yew, Zinnia.

HERBS, SPICES:

For the control of insects such as: Ants, Aphids, Fire Ants, Loopers, Mites, Plant Bugs, Thrips, White Flies.

Herbs and Spices: Including but not limited to Anise, Balm, Basil, Burnet, Borage, Camomile, Caraway, Catnip, Chives, Clary, Coriander, Costmary, Cumin, Curry Leaf, Dill, Fennel (Italian & Sweet), Fenugreek, Horehound, Hyssop, Marigold, Marjoram (Sweet & Wild), Mint, Nasturtium, Oregano, Pennyroyal, Rosemary, Rue, Sage, Savory (Winter & Summer), Sweet Bay (Bay Leaf), Tansy, Tarragon, Thyme, Wintergreen, Woodruff, Wormwood.

FRUIT, NUT, VINE AND SPECIALTY CROPS:

Use Rate: Use 4 to 12 pounds per acre depending upon the insect population and amount of foliage. As a DUST, the best results are obtained when the plants are wet or dew still remains on the leaf surface. It is suggested that the plants are misted prior to application to obtain the maximum benefits and use the least amount of product for control. Reapply only as necessary. As a WETTABLE POWDER, apply in sufficient water to obtain maximum coverage. Dilution of the material should be at the maximum rate of one half pound in one half gallon of water or at a minimum of one pound to 25 gallons of water (4 tbsp. per gallon). It is best to keep the material agitated during application. Apply the material uniformly to both the upper and lower leaf surfaces. Reapply only as necessary. Rates will vary with insect populations and amount of foliage, but adequate control can be obtained at 4 to 12 pounds product per acre. Reapply only as necessary.

FOR THE CONTROL OF SUCH INSECTS AS Ants, Armyworms, Aphids, Blueberry Maggots, Cabbage Loopers, Caterpillars, Cherry Fruit Flies, Cutworms, Fire Ants, Fireworms, Fruit Flies, Fuller Rose Beetle, Gooseberry Fruit Worms, Imported Cabbage Worms, Japanese Beetles, Leafhoppers, Mites, Peach Borers, Pecan Weevil, Red-necked Borers, Raspberry Fruit Worms, Rose Chafers, Stink Bugs, Strawberry Leaf Rollers, Thrips, Weevils, Webworms, White

Flies.

CITRUS FRUITS: Including but not limited to Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (Sweet & Sour), Pummelo, Satsuma Mandarin.

POME FRUITS: Including but not limited to Apple, Crabapple, Loquat, Pear, Oriental Pear, Quince.

STONE FRUITS: Including but not limited to Apricot, Cherry (Sweet & Sour), Nectarine, Peaches, Plums, Prunes, Chickasaw Plums, Damaon Plum, Japanese Plum.

SMALL FRUITS AND BERRIES: Including but not limited to Blackberry, Blueberry, Boysenberry, Cranberry, Currants, Dewberry, Elderberry, Gooseberry, Grapes, Huckleberry, Loganberry, Ochia Berry, Raspberry (Black & Red), Strawberry, Youngberry.

SUBTROPICAL FRUITS: Including but not limited to Avocado, Banana, Carob, Barbados Cherry, Cherimoya, Dates, Feijoa, Figs (Adriatic, Calimyna, Kadota, Black Mission, California Brown Turkey, Brunswick), Guava, Kiwifruit, Lychee, Mango, Papaya, Passion Fruit, Persimmon, Pineapple, Pomegranate.

TREE NUTS: Including but not limited to Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (Hazelnut), Hickory Nut, Japanese Horsechestnut, Macadamia Nut (Bushnut), Pecan, Pistachio, Walnut (Black & English).

SPECIALTY CROPS: Such as Artichoke, Chayote, Asparagus, Coffee, Cotton, Hops, Jojoba, Ornamental Turf Grass, Sesame, Sunflower (Leaves & Seed), Tea, Tobacco.

On harvested tomatoes and fruit including grapes: Fruit Flies, Vinegar Flies. Apply with a hand or power sprayer or other suitable means on tomatoes and fruits in baskets, on trucks or in plants, raw stock stacked in the yard. Apply liberally at the rate of 5 oz. per 100 sq. ft. of area.

As a grain protectant: To protect against grain storage insects, dilute one pound in 5 gallons of water and apply directly to the grain at the rate of 4 to 5 gallons per 1000 bushels as the grain is carried along a belt or as it enters the auger or elevator. Monthly inspection should be made if the top two or three inches are found to be reinfested. Retreat at the rate of 1 to 2 gallons of dilution per 1000 bushels of stored product.

In mushroom production and processing: Mushroom Flies, Fungus Gnats. Apply dilution at a rate of 5 oz. per 1000 cu. ft. of space. Direct spray 8 to 10 feet away from beds and slightly upwards over beds. Keep circulating fans running but turn off refrigerators during application. In processing plants, use at the rate of 5 oz. per 2000 cu. ft. of space.

Mosquito Control: May be used on croplands as an adulticide to protect agricultural workers during harvesting. Use 2 to 12 ounces per acre when applied by fixed wing aircraft, helicopters or truck mounted equipment. Do not apply in a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

COMMERCIAL INDOOR AND OUTDOOR USE:

For use in and around homes, office buildings, restaurants, motels, warehouses, theaters, schools, hotels and food handling establishments. This product is intended for application with hand or power duster, sprayer or other suitable means, to hiding and running areas, and places where pests are found. Remove or cover food prior to use. All food preparation surfaces and utensils should be covered during treatment, or thoroughly washed before use.

BEST AVAILABLE COPY

SP/

CONTROLS: Cockroaches, Ants, Fire Ants, Earwigs, Silverfish, Crickets, Millipedes, Box Elder Bugs, Scorpions and Other Similar Crawling Insects: Coat with a layer of dust, or spray as a wettable powder (mix 6 Tbsp. per gallon of water) in areas where these pests are found or may hide such as cracks and crevices, behind and beneath stoves, refrigerators, sinks, cabinets, garbage cans, garbage trucks, in and around sewer pipes, drains, window frames, in attics, basements, driveways, sidewalks, patios around houses and buildings. Apply directly to insects when possible.

GRASSES, LAWNS & TURF (INCLUDING GOLF COURSES): Controls Armyworms, Caterpillars, Chinch Bugs, Cutworms, Fleas, Ants, Fire Ants, Ticks.

Use Rate: Use 16 oz. (1 lb.) per 1000 sq. ft. of area. Apply either as a DUST or as a WETTABLE POWDER. Use hose-end sprayer for lawns, add a small amount of water to the jar and add the amount needed for the measured area, stir to make a slurry, then add the remaining water to the top of the jar. Empty entire contents of the jar on the premeasured area.

FIRE ANTS:

Mound Drench or Water Method: Use 0.6 oz. (4 Tbsp.) to one gallon of water. For best results, saturate the perimeter of the mound first working toward the center of the mound in a circular motion (you may want the mixture to puddle). An ant mound 12-14 inches in diameter requires approximately one gallon of the mix.

Hand Sprayer: Use 0.6 oz. (4 Tbsp.) to one gallon of water. It is best to keep the material agitated during application. Apply the contents to a premeasured area.

Hose-end Sprayer: Use 16 oz. (1 lb.) per 1000 sq. ft. of area. Use a hose-end sprayer for lawns, add a small amount of water to the jar and add the amount needed for the measured area, stir to make a slurry, then add the remaining water to the top of the jar. Empty the entire contents of the jar on the premeasured area.

As a Dust: For best results, dust the perimeter of the mound first. With a stick, disturb the mound, then dust mound in a circular motion working toward the center until the entire mound is thoroughly dusted.

HOUSE FLIES: Apply to surface of window frames, around light fixtures, on porches, in garages and other similar places where these insects light or congregate.

BEDBUGS: Treat liberally baseboards, moldings, cracks, bedstead, and both sides of mattress.

FLEAS: Thoroughly treat floor, bedding in and around pet's sleeping quarters. Treat surrounding cracks and crevices, baseboards, furniture, carpeting and wherever fleas are suspected. Dust rugs and carpets with thin coat of dust. Rake (not sweep) nap back and forth, causing dust to enter rug or carpet. Leave overnight. When cleaning do not make special effort to remove dust in rug or carpet. It will kill live fleas, repel others and help deodorize carpet or rug.

ROACH CONTROL IN PUBLIC SEWERS: A power blower using 1 pound of product will coat the dry part of the sewer line and manhole for a maximum of 300 ft. each direction. It will also enter the sewers and dust may be seen emerging from the vent pipes on the houses on each side of the street.

FOOD AND FEED HANDLING ESTABLISHMENTS: Places other than private residence in which food is held, processed or served. **Food and Feed Areas:** Includes areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage and enclosed processing systems (mills, dairies, edible oils,

syrupe). (Serving areas when food is exposed and facility is in operation are also considered a food area.) Application limited to spot or crack and crevice treatments only.

Crack and Crevice Treatments Only: Apply in small amounts directly into cracks and crevices using equipment capable of directing insecticide in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases where cockroaches, ants, silverfish and crickets hide. Spot treatment may also be made to areas where any of the above pests hide or inhabit such as baseboards, storage areas, closets, around water pipes, doors, windows, behind and under refrigerators, cabinets, sinks, stoves and other equipment. Care should be taken to avoid depositing the product onto exposed food contact surfaces or introducing the material into the air.

Spot Treatment: No individual spot will exceed two (2) square feet. Limit spot treatments to wall or floor surfaces where pests have been seen or where pests are suspected of hiding.

DO NOT USE IN: Edible product areas of food/feed handling establishments, restaurants or other areas where food/feed are commercially prepared or processed. Do not use in serving areas while food is exposed or facility is in operation. Serving areas are areas where prepared foods may be prepared or held. In the home, cover all food handling surfaces and cover or remove all food and cooking utensils, or wash thoroughly after treatment.

NON-FOOD AREAS: Including garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

DOGS, CATS AND PETS:

CONTROLS: Fleas, Flies, Lice and Ticks. As a dust, lightly work the material into the hair of the animal, down to the skin. Pay strict attention to places where the animal cannot reach, such as the scalp. Dust the ears of the animal with care. Dust the animal's bedding and areas where the animal sleeps, as well as cracks and crevices adjacent to them. Repeat as necessary. Individual ticks observed on the animal should be dusted thoroughly.

As a spray, mix 4 Tbsp. per gallon of water. Spray animal thoroughly.

For wet application, mix 4 Tbsp. per gallon of water. Dip animals in solution paying attention to make sure animals are completely covered with material.

ANIMAL QUARTERS (including and not limited to Barns, Dairies, Milkrooms, Pet Kennels, Poultry Houses, Rums and Stables).

CONTROLS: Ants, Gnats, Fire Ants, Fleas, Flies, Lice, Mites, Mosquitoes, Ticks.

USE RATE: Use at the rate of 2 oz. per 400 sq. ft. of surface area. Apply with a hand or power duster or sprayer or other suitable means to the hiding places where pests are found. Repeat the treatment as necessary to maintain control.

SEE FRONT PANEL FOR ADDITIONAL INFORMATION AND DIRECTIONS.