

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



United States
Environmental Protection
Agency

Office of Pesticide Programs

FILE COPY

December 19, 2007

Donna Leventhal
KIK International Inc.
% Delta Analytical Corp.
7910 Woodmont Ave. Suite 1000
Bethesda, MD 20814

Subject: **KIK International Inc. / Pure Bright Germicidal Ultra Bleach**
EPA Registration Number: 70271-13
Application Dated: December 11, 2007
Receipt Date: December 12, 2007

Dear Ms. Leventhal:

This acknowledges receipt of your notification, Submitted under the provision of PR Notice 98-10, FIFRA Section 3(c) 9.

Proposed Notification

- To add non-pesticidal claims

General Comment

Based on a review of the material submitted, the following comment applies:

This notification is accepted and a copy has been inserted in your file for future reference.

Should you have any questions concerning this letter, please contact Wanda Henson at (703) 308-6345.

Sincerely,

A handwritten signature in black ink, appearing to read "Wanda Henson".

Wanda Henson
Product Reviewer (32)
Regulatory Management Branch II
Antimicrobials Division (7510P)



United States
Environmental Protection Agency
Washington, DC 20460

☐ Registration
☐ Amendment
☒ Other

OPP Identifier Number

NOTIF**Application for Pesticide - Section I**1. Company/Product Number **70271-13**2. EPA Product Manager
Emily Mitchell3. Proposed Classification
☐ None ☐ Restricted4. Company/Product (Name) **KIK International Inc. /
Pure Bright Germicidal Ultra Bleach**PM# **32**5. Name and Address of Applicant (Include ZIP Code)
**KIK International Inc.
c/o Delta Analytical Corp.
7910 Woodmont Ave. Suite 1000
Bethesda MD 20814**
☐ Check if this is a new address6. **Expedited Review.** In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to:

EPA Reg. No. _____

Product Name _____

Section - II☐ Amendment - Explain below
☐ Resubmission in response to Agency letter dated _____
☒ Notification - Explain below.☐ Final printed labels in response to Agency letter dated _____
☐ "Me Too" Application.
☐ Other - explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification under PR Notice 98-10 to add non-pesticidal claims

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III**1. Material this Product will be Packaged in:**Child-Resistant Packaging
☐ Yes*
☐ NoUnit Packaging
☐ Yes
☐ NoWater Soluble Packaging
☐ Yes
☐ No2. Type of Container
☐ Metal
☐ Plastic
☐ Glass
☐ Paper
☐ Other (Specify) _____*** Certification must
be submitted.**If "Yes,"
Unit Package wgt. No. per containerIf "Yes,"
Unit Package wgt. No. per container3. Location of Net Contents Information
☐ Label ☐ Container

4. Size(s) of Retail Container

5. Location of Label Direction
☐ On Label
☐ On Labeling accompanying product

6. Manner In Which Label Is Affixed to Product

☐ Lithograph
☐ Paper glued
☐ Stenciled☐ Other _____**Section - IV****1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)**Name
Donna LeventhalTitle
**Agent for
KIK International Inc.**Telephone No. (Include Area Code)
301-680-7971**Certification**

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any kind of knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature

3. Title

Agent for KIK International Inc.

4. Typed Name

Donna Leventhal

5. Date

December 11, 20076. Date Application
Received**(Stamped)**



December 11, 2007

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Avenue, N.W.
Washington, DC 20460

Attn: Emily Mitchell, PM 32

RE: Notification to add non-pesticidal claims
Product: Pure Bright Germicidal Ultra Bleach
EPA Reg. No.: 70271-13
Company: KIK International, Inc.

Dear Ms. Mitchell:

On behalf of KIK International, Inc., I am submitting a notification to add non-pesticidal claims to the label for Pure Bright Germicidal Ultra Bleach, EPA Reg. No. 70271-13. The following non-pesticidal claims have been added:

- "Around the House" to be added to the already approved claim "Disinfects, Sanitizes and Deodorizes" (per New York request)
- Value Size 2 [or 3] Pack
- Do not Break Open - To Be Sold As A Two [or Three] Pack
- Buy 3 Get 1 Free
- Compatible for Use in High Efficiency Washing Machines [or HE or High Efficiency Compatible]

The following documents are attached in support of this notification.

- EPA form 8570-1
- Two copies of page 2 of 24 with the new text noted above highlighted
- One complete copy of the label

Please note that no other changes have been made to this document.

4 8 28

If you have any questions regarding this submission, please contact me at 301-680-7971 or dleventhal@delta-ac.com.

Sincerely,



Donna Leventhal
Agent for KIK International, Inc.

Enclosures

h:\clients\kik\ultra\epa04a.notif.cvr.wpd

PURE BRIGHT™ GERMICIDAL ULTRA BLEACH

Active Ingredient:
Sodium Hypochlorite 6.0%
Other Ingredients: 94.0%
TOTAL: 100.0%

KEEP OUT OF REACH OF CHILDREN

DANGER [: Corrosive]

FIRST AID: Call poison control center or doctor for treatment advice. Have the product container or label with you when calling a Poison Control Center or doctor, or going for treatment. **If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. **If on skin or clothing:** Take off contaminated clothing. Rinse skin with plenty of water for 15-20 minutes. **If swallowed:** Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

See side [or back] panel for additional precautionary labeling. [additional precautions]

[Brackets above indicate optional language]

Manufactured by: KIK International, Inc.
33 Macintosh Blvd. Concord Ontario L4K4L5
Santa Fe Springs CA 90670 EPA Est No. 70271-CA-2 ☐
Denver CO 80207 EPA Est No. 70271-CO-1 ☐
Auburndale FL 33823 EPA Est No. 70271-FL-1 ☐
Hampton GA 30228 EPA Est No. 70271-GA-1 ☐
Eagan MN 55121 EPA Est No. 70271-MN-1 ☐
Houston TX 77054 EPA Est No. 70271-TX-1 ☐
Salem VA 24153 EPA Est No. 70271-VA-1 ☐
Tacoma WA 98421 EPA Est No. 70271-WA-1 ☐
Concord ON CN L4K4K5 EPA Est No. 70271-CAN-1 ☐

EPA Reg. No. 70271- 13

[actual establishment shown in code above or below label printed on bottle]

NET 96 FL. OZ. (3 QT.) 2.84

5225

[Statements in brackets on the following pages are instructional and not part of labeling]

[Additional optional claims:]

25.0% MORE PRODUCT [COMPARED TO THE 48 FL. OZ. SIZE]

BACTERICIDAL

BACTERICIDE • FUNGICIDE • *VIRUCIDE

BRIGHTENS LAUNDRY

BUY 3 GET 1 FREE

CLEANS

COMMERCIAL/INSTITUTIONAL USE

COMPATIBLE FOR USE IN HIGH EFFICIENCY WASHING MACHINES [OR HE OR HIGH

EFFICIENCY COMPATIBLE

DISINFECTS AND DEODORIZES

DISINFECTS • SANITIZES • DEODORIZES

DISINFECTS, SANITIZES, DEODORIZES AROUND THE HOUSE

DO NOT BREAK OPEN. TO BE SOLD AS A TWO [OR THREE] PACK

ELIMINATES ODORS

FOR A CLEANER, FRESHER HOUSEHOLD & [AND] LAUNDRY

FORMULATED FOR FOODSERVICE APPLICATIONS

FUNGICIDAL

HIGHER CONCENTRATION FORMULA

JUST AS SAFE ON BLEACHABLE FABRICS AS BEFORE

KILLS GERMS

KILLS *HIV-1 ON HARD NON-POROUS ENVIRONMENTAL SURFACES

MILDEWICIDAL

REMOVES STAINS

SAFE ON BLEACHABLE FABRICS

SPLASH RESISTANT WITH AN EASY OPEN SEAL

VALUE SIZE 2 [OR 3] PACK

WHITENS

CONTAINS SODIUM HYPOCHLORITE THE ACTIVE INGREDIENT USED IN [brand name ® or TM]

This product can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms), eating establishments, pet kennels, veterinary premises, farms, dairies, and food processing plants.

Kitchen: Clean and disinfect kitchen sinks and countertops.

Bathroom: Deodorize and disinfect bathtubs, showers, floors, vinyl, and tile.

Toilet Bowls: Sanitize your toilet.

Exercise care in handling. Check to make sure bottle is always tightly capped. Product should be carried and stored in an upright position to avoid spillage.

Do not use this product full strength for cleaning surfaces. Always dilute strictly in accordance with label directions. Wear gloves when cleaning for prolonged periods.

Always mix this product strictly in accordance with label instructions. Do not use bleach at full strength for cleaning. Protect yourself by wearing rubber gloves.

Contains no phosphorous.

Keep upright. Keep bottles upright and tightly capped.

Do not drop.

ULTRA FORMULA: For regular loads, just 3/4 cup provides the same whitening and cleaning benefits as 1 cup of regular bleach. Smaller bottle is easier to handle, pour and store.

[The following headings may be placed above or in front of appropriate directions (e.g., Hospital Use in front of HIV directions, Household Use in front of directions for dishes or floors)]

FARM - DAIRY USE

RESTAURANT USE

FOODSERVICE APPLICATIONS

HOSPITAL USE

HOUSEHOLD USE

FOOD PROCESSING PLANT USE

[Additional optional claims; parenthetical terms need not be included:] *The phrase "this product" may be used interchangeably with the product name "Pure Bright Germicidal Ultra Bleach" throughout the label* [The claims on this page (except for HIV) are for use on product labels that include directions for use as a disinfectant on pre-cleaned, hard, non-porous surfaces at a use rate of at least 1 part product to 21.3 parts water (e.g., 3/4 cup (6 oz.) or more product per gallon of water) at a 5 minute contact time. For HIV see note at bottom of page.]

Kills [list any of the organisms below] [and] 99.9% of common household germs¹

Kills 99.9% germs¹

Pure Bright Germicidal Ultra Bleach kills germs¹

¹ Kills common household germs including: [list any of the organisms below]. or ¹Effective at killing [list any of the organisms below] [on hard, non-porous surfaces].

This product, when used as directed on hard, non-porous surfaces, is effective against the following:

FUNGICIDAL

- Trichophyton mentagrophytes (Athlete's foot fungus)
- Aspergillus niger (mildew)

BACTERICIDAL

- Escherichia coli 0157:H7 (E. coli)
- Pseudomonas aeruginosa (pseudomonas)
- Salmonella choleraesuis (salmonella)
- Shigella dysenteriae
- Staphylococcus aureus (staph)
- Streptococcus pyogenes (strep)

VIRUCIDAL*

- *Adenovirus type 2
- *Avian Influenza A virus
- *Canine Parvovirus
- *Cytomegalovirus
- *Feline Panleukopenia virus (Parvovirus)
- *Hepatitis A virus
- *Herpes simplex virus 1 (herpes)
- *Herpes simplex virus 2 (herpes)
- *Human Immunodeficiency Virus type 1 (HIV-1) [+]
- *Influenza A virus
- *Poliovirus type 1
- *Respiratory syncytial virus
- *Rhinovirus type 37 (viruses that cause colds and flu)
- *Rotavirus

[Or, optional: can be listed in the following format:]

Pure Bright Germicidal Ultra Bleach kills germs:

Kills common household germs including Staphylococcus aureus (staph), Streptococcus pyogenes (strep), Salmonella choleraesuis (salmonella), Pseudomonas aeruginosa (pseudomonas), *Influenza A virus, *Rhinovirus type 37 (viruses that cause colds and flu), Trichophyton mentagrophytes (Athlete's foot fungus), Aspergillus niger (mildew), *Rotavirus and Escherichia coli 0157:H7 (E. coli). Also kills *Adenovirus type 2, *Avian Flu virus, *Hepatitis A, *Cytomegalovirus, *Respiratory Syncytial Virus, *Poliovirus type 1, *Herpes Virus 1 (herpes), *Herpes Virus 2 (herpes), *Canine Parvovirus, *Feline Parvovirus, Shigella dysenteriae, *Human Immunodeficiency Virus type 1 (HIV-1) [+]

[+ If HIV-1 claim is used, HIV-1 directions must be included on the printed label.]

Kills viruses that cause colds and flu** **Influenza A virus, Rhinovirus type 37

Kills avian influenza A on pre-cleaned environmental surfaces

PRECAUTIONARY STATEMENTS**Hazards to Humans and Domestic Animals**

DANGER: Corrosive, may cause severe skin and eye irritation or chemical burns to broken skin. Causes eye damage. Wear safety glasses and rubber gloves when handling this product. Wash after handling. Avoid breathing vapors. Vacate poorly ventilated areas as soon as possible. Do not return until strong odors have dissipated.

[In accordance with PR notice 95-1, the Environmental Hazards statement is only required for containers 5 gallons and larger.]

Environmental Hazards

This product is toxic to fish and aquatic organisms. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or public waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

Physical and Chemical Hazards

Strong oxidizer. Flush drains before and after use. Do not use or mix with other household chemicals, such as toilet bowl cleaners, rust removers, acid or ammonia containing products. To do so will release hazardous gasses. Prolonged contact with metal may cause pitting or discoloration.

STORAGE AND DISPOSAL

Store in a cool, dry area away from direct sunlight and heat to avoid deterioration. In case of spill, flood area with large quantities of water. Product or rinsates that cannot be used should be diluted with water before disposal in a sanitary sewer. Do not re-use empty container and place in trash collection. Do not contaminate food or feed by storage, disposal or cleaning of equipment.

[The Directions for Use on the following pages are similar directions with different formats and quantities in order to provide flexibility to distributors to address their customers' and market needs on labeling for the marketplace.]

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

One Tbsp. (½ oz.) of this product in 1 gallon of water is equivalent to approximately 200 parts per million (ppm) available chlorine. Use a chlorine test kit to determine exact available chlorine concentration and adjust as necessary to obtain specified ppm.

LAUNDRY

Use to bleach white and colorfast Acrylics, Cotton, Nylon, Polyester, Rayon (test to be sure). Do not use on Acetate, Leather, Silk, Spandex or Wool, Mohair and non-fast colors. Sort laundry by color and fabric. Separate whites from colors, light colors from dark colors.

BLEACH TEST: Before using, mix two tablespoons of bleach with 1/4 cup of water in a glass, rubber, porcelain, or plastic container and test a small piece of fabric in a place that doesn't show. Test all colors, including trim. Let stand one minute, then blot dry. No color change means the article can be bleached safely:

LAUNDRY: Before adding clothes, mix ¾ cup of bleach with water in top-loading 16 gallon machines or mix ⅓ cup bleach with water in front-loading 8 gallon machines. For large top loading automatics or larger heavily soiled loads, use 1¼ cup. Add clothes. If sanitization is desired, soak clothes for 5 minutes. Wash and rinse with usual cycles. If clothes are in machine the addition of bleach can cause damage.

TOP LOAD AUTOMATIC..... ¾ CUP LARGE TOP LOADING AUTOMATIC .. 1¼ CUP
FRONT LOAD AUTOMATIC..... ⅓ CUP LARGE/HEAVY SOILED LOADS 1¼ CUP

REMOVE STAINS: Mix ¼ cup of bleach with a gallon of water. Soak stained area for 5 minutes to remove grass, ink, coffee, tea, scorch, fruit, etc. Rinse thoroughly.

[OR: the following format may be used:]

LAUNDRY USE:

1. Before adding clothes, mix ¾ cup of bleach with water in top-loading 16 gallon machines or mix ⅓ cup bleach with water in front-loading 8 gallon machines. For large top loading automatics or larger heavily soiled loads, use 1¼ cup.

2. Add clothes.

3. Wash and rinse with usual cycles.

Do not use on Acetate, Leather, Silk, Spandex or Wool, Mohair and non-fast colors.

REMOVE STAINS:

1. Mix ¼ cup of bleach with a gallon of water.

2. Soak stained area for 5 minutes to remove grass, ink, coffee, tea, scorch, fruit, etc.

3. Rinse thoroughly.

[OR: additional format:]

LAUNDRY USE:

1. Sort your laundry by color and fabric type.

2. Add ¾ cup of bleach and laundry detergent to wash water. For extra large machines, add 1¼ cups.

3. Add laundry.

Do not bleach fabrics such as wool, silk, mohair, leather, spandex or nonfast colored fabrics.

Helpful Hint: We recommend mixing ¾ cup of bleach with 1 quart of water and adding 5 minutes after wash load has started.

[For all directions, superscript¹ refers to the claims section listing organisms against which the product is effective.]

Kitchen: Clean and disinfect countertops and sinks [or Clean, disinfect, and eliminate mildew on countertops and sinks]:

1. Use 3/4 cup of Pure Bright Germicidal Ultra Bleach [or this product] per gallon of water.
2. Wash and rinse surfaces then apply disinfecting solution.
3. Let stand 5 minutes.
4. Rinse thoroughly and air dry.

Kitchen and Bathrooms: Disinfect, deodorize [and kill 99.9% of germs¹] on counters, in sinks, showers, bathtubs and on vinyl and tile floors.

1. Wash surfaces with water.
2. Disinfect surfaces with a solution of 3/4 cup of bleach per gallon of water.
3. Let stand 5 minutes then rinse with water and allow to dry.

Disinfecting and Deodorizing Bathrooms [or Disinfecting, Deodorizing, and Eliminating Mildew in Bathrooms]: Disinfect and deodorize bathtubs, showers, sinks, floors, vinyl, tile.

1. Prewash surfaces and rinse.
2. Spread a solution of 3/4 cup of Pure Bright Germicidal Ultra Bleach [or this product] per gallon of water [or Spread a solution of 1 1/2 cups of Pure Bright Germicidal Ultra Bleach [or this product] per 2 gallons of water.]
3. Let stand 5 minutes, then drain and air dry.

Toilet bowls: Sanitize and deodorize toilet

1. Prewash toilet and flush.
2. Pour 1 cup of Pure Bright Germicidal Ultra Bleach [or this product] - swab with brush, making sure to get under the rim.
3. Let stand 10 minutes and flush. DO NOT use with bowl cleaners or any other household chemicals.

or:

Toilet Bowls: Sanitize and deodorize your toilet.

1. Prewash and flush your toilet before pouring 1 cup of bleach into the bowl.
2. Using a brush, scrub the entire bowl making sure to get under the rim.
3. Let stand 10 minutes and then flush. DO NOT use with bowl cleaners or any other household chemicals.

Disinfects, sanitizes, [eliminates mildew] and deodorizes by killing most germs and their odors.

HAND SANITIZING: Thoroughly wash hands with soap or detergent and rinse. Mix approximately 1 tsp. of bleach per gallon of water to prepare a 50 ppm solution (use chlorine test kit to determine exact available chlorine). Submerge hands. Let air dry.

SANITIZING FOOD CONTACT SURFACES: Before using this product, remove or carefully protect food. Remove gross food particles from surface. Prewash surface with a good detergent and rinse thoroughly with potable water. Mix approximately 1 Tbsp. of bleach per gallon of water to prepare a 200 ppm available chlorine solution. Cover surface with bleach solution for at least 2 minutes. Air dry.

DISINFECTING KITCHEN, DISHES, SINKS: [1.] Use ¼ cup bleach mixed with a quart of water to soak cleaned dishes, teapot, cups, sinks, etc. for 5 minutes. [2.] Rinse with a solution of approximately 1 Tbsp. of bleach per gallon of water to prepare a 200 ppm solution. Do not use on silverware. Bleach solution can be used on porcelain, enamel, etc. surfaces after cleaning. [3.] Let air dry.

DISINFECTING [AND ELIMINATING MILDEW ON] WALLS, FLOORS, AND OTHER HARD INANIMATE SURFACES NOT IN DIRECT CONTACT WITH FOOD: Prewash surfaces and rinse. Mix ¾ cup bleach per gallon of water. Spray, rinse, or wipe surface with bleach solution and let stand for 5 minutes. Drain and air dry.

SANITIZING TOILET BOWL: Prewash toilet and flush. Pour 1 cup bleach into toilet bowl, scrub with a brush, making sure to get under rim, let stand 10 minutes, flush. Do not use with bowl cleaners or any other household chemicals.

DEODORIZING GARBAGE CANS: Wash and rinse. Use ¾ cup bleach for each gallon of water in can. Empty and let drain.

EGG SHELL SANITIZING: Thoroughly clean eggs. Mix approximately 1 Tbsp. (½ oz) of bleach per gallon of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130°F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable rinse. The solution should not be re-used to sanitize eggs.

Plastic Cutting Boards:

1. Wash with water and dish detergent.
2. Clean with a solution of 1 tablespoon of bleach per gallon of water.
3. Let stand 2 minutes then rinse with water and allow to dry.

Wooden Cutting Boards:

1. Wash with water and dish detergent.
2. Clean with a solution of 3 tablespoons of bleach per gallon of water.
3. Let stand 2 minutes then rinse with a solution of 1 tablespoon of bleach per gallon of water and allow to dry.

[optional headings: [Veterinary Clinics] [Animal Life Science Laboratory] [Zoos] [Pet Shops] [Kennels] [Breeding and Grooming Establishments] [Animal Housing Facilities] [Poultry houses] [Hatcheries]:]

Special Instructions for Inactivating Avian Influenza A

For cleaning and disinfecting hard, non-porous surfaces: equipment, utensils, instruments, cages, kennels, stables, and catteries. Remove all poultry [or animals] and feeds from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings, and manure from floors, walls, and surfaces of facilities occupied or transversed by poultry [or animals]. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use solution of $\frac{3}{4}$ cup (6 fl. oz.) of product per gallon of water and let stand for 5 minutes, drain and air dry. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, scrappers used in removing litter and manure. Ventilate buildings, coops, and other closed spaces. Do not house poultry [or animals] or employ equipment until treatment has been absorbed, set or dried. All treated feed/water bowls, racks, troughs, automatic feeders, fountains, and waterers must be rinsed with potable water before reuse.

BLEACH USE TIPS:

Disinfecting Your Pet's Food Bowls & Litter Boxes*:

1. Wash thoroughly with water and dish detergent.
 2. Disinfect bowls with 1 tablespoon of bleach per gallon of water.
 3. Let stand 2 minutes then rinse with water and allow to dry.
- *For litter boxes, repeat step one and use $\frac{3}{4}$ cup of bleach mixed with a gallon of water. Then repeat step 3.

Removing Mold & Mildew Stains from Outdoor Patios:

1. Protect plants from overspray.
2. Use 1 cup of bleach per gallon of water to clean unsightly stains using a brush.
3. Rinse entire area, including plants, with water for 5 minutes.

Kill Germs and Odors in Garbage Cans:

1. Rinse with soap and water.
2. Put a solution of $\frac{3}{4}$ cup of bleach per gallon of water in the garbage can.
3. Let stand 5 minutes then drain.

Disinfecting Children's Furniture and Toys:

1. Ensure all surfaces are colorfast.
2. To kill 99.9% of germs¹, use a solution of $\frac{3}{4}$ cup of bleach per gallon of water to clean children's surfaces.
3. Let stand for 5 minutes, then rinse and allow to dry.

One Tbsp. (½ ounce) of Pure Bright Germicidal Ultra Bleach [or this product] in 1 gallon of water is equivalent to approximately 200 parts per million (ppm) available chlorine. Use a chlorine test kit to determine exact available chlorine concentration.

	Amount Bleach	Amount Water	Instructions
FOR SANITIZING			
Work Surfaces	1 Tbsp.	1 Gallon	Pre-wash with detergent, rinse, cover surface with bleach solution for at least 2 minutes, drain, let air dry.
Dishes, Glassware, Utensils	1 Tbsp.	1 Gallon	After washing, soak for at least two minutes in bleach solution. Drain and let air dry.
Bathtubs, Showers	1 Tbsp.	1 Gallon	Wash, rinse, apply bleach solution for at least two minutes, drain, let air dry.
Refrigerators, Freezers	1 Tbsp.	1 Gallon	Wash, rinse, apply bleach solution for at least two minutes, drain, let air dry.
FOR DISINFECTING			
Floors & Walls	¾ Cup	1 Gallon	Pre-wash surfaces and rinse. Spray, rinse or wipe surface with bleach solution, let stand for 5 minutes. Drain and air dry.
Mops, Brushes, Brooms & Rugs	¾ Cup	1 Gallon	Wash with detergent, apply bleach solution, soak at least 10 minutes. Rinse well.
FOR MOLD AND MILDEW			
Bathrooms and Kitchens	¾ Cup	1 Gallon	Pre-wash with detergent, rinse, cover surface with bleach solution for at least 5 minutes, drain, let air dry. Food contact surfaces must be rinsed with potable water before use.
FOR DEODORIZING			
Garbage Cans	¾ Cup	1 Gallon	After washing and rinsing, brush inside with solution. Empty and let drain.
Drains	1 Cup	-----	Pour into drain. Flush with hot water.
FOR BLEACHING/WHITENING			
Wooden Surfaces	½ Cup	1 Gallon	Apply for 5 minutes, rinse.
FOR STAIN REMOVAL			
All Surfaces	1 Cup	1 Gallon	Add bleach to detergent solution, apply, rinse.
FOR AGRICULTURAL USES			
Food Egg	1 Tbsp.	1 Gallon	Wash, rinse, spray with maximum 130°F bleach solution until completely wet; let air dry before casing or breaking. Once solution is used it must not be reused for this purpose.
Fruits and Vegetables	1 Tbsp.	20 Gallons	Wash, rinse, soak in circulating bleach solution (25 ppm) for 2 min. Spray vegetables with bleach solution before packing; use potable water to rinse fruit before packaging. Rinse is to occur only immediately prior to packaging.

Reformatting of existing language in table

FOR SANITIZING - Mix 1 Tbsp bleach with 1 Gallon of water.

Work Surfaces: Pre-wash with detergent, rinse, cover surface with bleach solution for at least 2 minutes, drain, let air dry.

Dishes, Glassware, Utensils: After washing, soak for at least 2 minutes in bleach solution. Drain and let air dry.

Bathtubs, Showers: Wash, rinse, apply bleach solution for at least 2 minutes, drain let air dry.

Refrigerators, Freezers: Wash, rinse, apply bleach solution for at least 2 minutes, drain let air dry.

FOR DISINFECTING - Mix 3/4 Cup bleach with 1 Gallon of water.

Floors, Walls: Pre-wash surfaces and rinse. Spray, rinse or wipe surface with bleach solution, let stand for 5 minutes. Drain and air dry.

Mops, Brushes, Brooms, Rags: Wash with detergent, apply bleach solution, soak at least 10 minutes. Rinse well.

FOR MOLD AND MILDEW - Mix 3/4 Cup bleach with 1 Gallon of water.

Bathrooms and Kitchens: Pre-wash with detergent, rinse, cover surface with bleach solution for at least 5 minutes, drain, let air dry. Food contact surfaces must be rinsed with potable water before use.

FOR DEODORIZING - Mix 3/4 Cup bleach with 1 Gallon of water.

Garbage Cans: After washing and rinsing, brush inside with solution. Empty and let drain.

FOR DEODORIZING - Pour 1 Cup bleach in drain.

Drains: Pour into drain. Flush with hot water.

FOR BLEACHING/WHITENING - Mix 1/2 Cup with 1 Gallon of water.

Wooden Surfaces: Apply for 5 minutes, rinse.

FOR STAIN REMOVAL - Mix 1 Cup with 1 Gallon of water.

All surfaces: Add bleach to detergent solution, apply, rinse.

■**Laundering:** To bleach and sanitize white and colorfast cotton, linen, nylon, Dacron, Orlon, polyester, Dynel and rayon in washing machine: $\frac{3}{4}$ cup of this product per load for conventional washing machine; $\frac{1}{2}$ cup for front load automatic. Add to pre-soak, wash water or first rinse. If clothes are in machine, dilute product in 1 quart water before adding.

■**To Whiten Nylon and Other Synthetics** that have turned yellow or grey: 1 tablespoon of this product per gallon water. Soak clean fabric in solution for 15 to 20 minutes. Rinse well. Repeat if necessary.

■**To Remove Stains:** Berry, wine, coffee, tea, ink, grass, dye, medicine stains, scorch and mildew stains. Make solutions of 2 tablespoons of this product to each quart water. Immerse fabric for 5 to 10 minutes. Rinse well in clear water. Repeat if necessary.

■**Today's permanent Press Fabrics are Bleachable** and need this product to get out stains and help prevent dirt build up. Wash with regular laundry as directed: Top-load automatics – $\frac{3}{4}$ cup per load. Wringer-type washers – $\frac{3}{4}$ cup per load. Front-load automatics – $\frac{1}{2}$ cup per load. Use this product with any good laundry soap or detergent. If your washer has an automatic bleach dispenser, follow washer directions. If not, add this product to wash water before laundry is put in. If laundry is put in before wash water then dilute this product in quart of water and add after machine has started agitating and fabrics are thoroughly wet.

■**Disinfecting and Deodorizing Bathrooms:** To disinfect, deodorize, and eliminate mold and mildew from washable surfaces such as tubs, showers, countertops, sinks, ceramic tile and vinyl flooring, [1.] Spread a solution of $1\frac{1}{2}$ cups of this product per 2 gallons of water on clean surface. [2.] Let stand 5 minutes, then drain.

■**ToiletBowls:** To sanitize and deodorize pre-cleaned toilet bowls, use 1 cup of this product. [1.] Flush, pour in bleach – swab with brush, making sure to get under the rim, and [2.] let stand for 10 minutes. [3.] Flush. DO NOT use with bowl cleaners or any other household chemicals.

■**Sickroom Equipment:** Wash all surfaces thoroughly. Rinse, then spread a solution of $1\frac{1}{2}$ cups of this product per 2 gallons of water over all surfaces. Let stand 5 minutes, then drain.

■**Garbage cans:** Wash thoroughly with warm soapy solution. Rinse then spread a solution of $1\frac{1}{2}$ cups of this product per 2 gallons of water over all surfaces. Let stand 5 minutes, then drain.

Avoid prolonged contact with metal since corrosion or discoloration may occur. Do not use this product on chipped enamel.

■**Sanitizing Nonporous Food Contact Surfaces:** Prepare a sanitizing solution by thoroughly mixing 2 Tbsp. (1 oz.) of this product with $2\frac{1}{2}$ gallons of water to provide approximately 200 ppm available chlorine by weight. Clean all surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

■**To Sanitize Milking Equipment:** Prepare sanitizing solution as above immediately prior to use. All surfaces to be sanitized should be properly cleaned before application of chlorine solution. Milking utensils should be submerged in the solution for at least 2 minutes and allowed to drain. Do not rinse equipment with water after treatment.

If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Sanitizers used in automated systems may be used for general cleaning but may not be reused for sanitizing purposes.

■**Sanitizing Porous Food Contact Surfaces:** Prepare a solution of approximately 600 ppm by thoroughly mixing 6 Tbsp. (3 ozs.) of this product with $2\frac{1}{2}$ gallons of water. Clean surfaces in the normal manner. Rinse all surfaces thoroughly with the 600 ppm solution, maintaining contact with the sanitizer for at least 2 minutes. Prepare a 200 ppm sanitizing solution by thoroughly mixing 2 Tbsp. (1 oz.) of this product with $2\frac{1}{2}$ gallon of water. Prior to using equipment, rinse all surfaces with 200 ppm available chlorine solution. Do not rinse with water and do not soak equipment overnight.

PROPORTIONS FOR DILUTION OF THIS PRODUCT

200 ppm: 1 oz. (2 Tbsp) in $2\frac{1}{2}$ gallons of water

600 ppm: 3 oz. (6 Tbsp) in $2\frac{1}{2}$ gallons of water

(Use a chlorine test kit to determine exact available chlorine concentration and adjust dosage as necessary.)

RESTAURANTS, TAVERNS, SODA FOUNTAINS, DAIRIES, ETC.

DIRECTIONS FOR SANITIZING EATING AND DRINKING UTENSILS:

Prepare sanitizing solution immediately prior to use.

1. Scrape and pre-wash utensils and glass whenever possible.
2. Wash with good detergent or compatible cleaner.
3. Rinse with clean water.
4. Sanitize in solution of 1 oz to $2\frac{1}{2}$ gallons of water (200 ppm).
5. Immerse utensils at least 2 minutes or for contact time specified by governing sanitary code.
6. Do not reuse sanitizing solution.

15 2 28

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NOTE: This product degrades with age. Use a chlorine test kit and increase dosage, as necessary, to obtain the required level of available chlorine.

TABLE OF PROPORTIONS AVAILABLE CHLORINE

600 ppm – 12 oz. in 10 gal. water

200 ppm – 4 oz. in 10 gal. water

SWIMMING POOL WATER DISINFECTION

For a new pool or spring start-up, superchlorinate with 107 to 213 oz. of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Adjust and maintain pool water pH to between 7.2 to 7.6. Adjust and maintain the alkalinity of the pool to between 50 to 100 ppm.

To maintain the pool, add manually or by a feeder device 21.3 oz. of this product for each 10,000 gallons of water to yield an available chlorine residual between 0.6 to 1.0 ppm by weight. Stabilized pools should maintain a residual of 1.0 to 1.5 ppm available chlorine. Test the pH, available chlorine residual and alkalinity of the water frequently with appropriate test kits. Frequency of water treatment will depend upon temperature and number of swimmers.

Every 7 days, or as necessary, superchlorinate the pool with 107 to 213 oz. of product for each 10,000 gallons of water to yield 5 to 10 ppm available chlorine by weight. Check the level of available chlorine with a test kit. Reentry into treated pools is prohibited above levels of 4.0 ppm due to risk of bodily harm.

At the end of the swimming pool season or when water is to be drained from the pool, chlorine must be allowed to dissipate from treated pool water before discharge. Do not chlorinate the pool within 24 hours prior to discharge.

WINTERIZING POOLS - While water is still clear & clean, apply 6.4 oz. of product per 1000 gallons, while filter is running, to obtain a 3 ppm available chlorine residual, as determined by a suitable test kit. Cover pool, prepare heater, filter and heater components for winter by following manufacturers' instructions.

SPAS, HOT-TUBS, IMMERSION TANKS, ETC.

SPAS/HOT-TUBS - Apply 10.7 oz. of product per 1000 gallons of water to obtain a free available chlorine concentration of 5 ppm, as determined by a suitable chlorine test kit. Adjust and maintain pool water pH to between 7.2 and 7.8. Some oils, lotions, fragrances, cleaners, etc. may cause foaming or cloudy water as well as reduce the efficiency of the product.

To maintain the water, apply 10.7 oz. of product per 1000 gallons of water over the surface to maintain a chlorine concentration of 5 ppm.

After each use, shock treat with 17 oz. of this product per 500 gallons of water to control odor and algae. Reentry into treated spas is prohibited above levels of 5.0 ppm due to risk of bodily harm.

During extended periods of disuse, add 6.4 oz. of product daily per 1000 gallons of water to maintain a 3 ppm chlorine concentration.

HUBBARD AND IMMERSION TANKS - Add 10.7 oz. of this product per 200 gallons of water before patient use to obtain a chlorine residual of 25 ppm, as determined by a suitable test kit. Adjust and maintain the water pH to between 7.2 and 7.6. After each use, drain the tank. Add 10.7 oz. to a bucket of water and circulate this solution through the agitator of the tank for 15 minutes and then rinse out the solution. Clean tank thoroughly and dry with clean cloths. **[NOT FOR USE IN CALIFORNIA]**

HYDROTHERAPY TANKS - Add 2.1 oz. of this product per 1000 gallons of water to obtain a chlorine residual of 1 ppm, as determined by a suitable chlorine test kit. Pool should not be entered until the chlorine residual is below 3 ppm. Adjust and maintain the water pH to between 7.2 and 7.6. Operate pool filter continuously. Drain pool weekly, and clean before refilling.

SANITIZATION OF NONPOROUS FOOD CONTACT SURFACES

RINSE METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to insure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2.1 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 4.25 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment and do not soak equipment overnight.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

IMMERSION METHOD - A solution of 100 ppm available chlorine may be used in the sanitizing solution if a chlorine test kit is available. Solutions containing an initial concentration of 100 ppm available chlorine must be tested and adjusted periodically to ensure that the available chlorine does not drop below 50 ppm. Prepare a 100 ppm sanitizing solution by thoroughly mixing 2.1 oz. of this product with 10 gallons of water. If no test kit is available, prepare a sanitizing solution by thoroughly mixing 4.25 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight.

Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. If solution contains less than 50 ppm available chlorine, as determined by a suitable test kit, either discard the solution or add sufficient product to reestablish a 200 ppm residual. Do not rinse equipment with water after treatment.

Sanitizers used in automated systems may be used for general cleaning but may not be re-used for sanitizing purposes.

FLOW/PRESSURE METHOD - Disassemble equipment and thoroughly clean after use. Assemble equipment in operating position prior to use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 4.25 oz. product with 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold under

pressure for at least 2 minutes to ensure contact with all internal surfaces. Remove some cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

CLEAN-IN-PLACE METHOD - Thoroughly clean equipment after use. Prepare a volume of a 200 ppm available chlorine sanitizing solution equal to 110% of volume capacity of the equipment by mixing the product in a ratio of 4.25 oz. product with 10 gallons of water. Pump solution through the system until full flow is obtained at all extremities, the system is completely filled with the sanitizer and all air is removed from the system. Close drain valves and hold under pressure for at least 10 minutes to ensure contact with all internal surfaces. Remove some cleaning solution from drain valve and test with a chlorine test kit. Repeat entire cleaning/sanitizing process if effluent contains less than 50 ppm available chlorine.

SPRAY/FOG METHOD - Preclean all surfaces after use. Use a 200 ppm available chlorine solution to control bacteria, mold or fungi and a 600 ppm solution to control bacteriophage. Prepare a 200 ppm sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 4.25 oz. product with 10 gallons of water. Prepare a 600 ppm solution by thoroughly mixing the product in a ratio of 12.8 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces treated with 600 ppm solution with a 200 ppm solution.

SANITIZING OF POROUS FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 12.8 oz. of this product with 10 gallons of water to provide 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes and allow the sanitizer to drain. Prepare a 200 ppm sanitizing solution by thoroughly mixing 4.25 oz. of this product with 10 gallons of water and rinse all surfaces with this 200 ppm solution. Do not rinse with water and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 12.8 oz. of this product with 10 gallons of water to provide 600 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution, maintaining contact for at least 2 minutes and allow the sanitizer to drain. Following this, prepare a 200 ppm sanitizing solution by thoroughly mixing 4.25 oz. of this product with 10 gallons of water and rinse all surfaces with this 200 ppm solution. Do not rinse with water and do not soak equipment overnight.

SPRAY/FOG METHOD - Preclean all surfaces after use. Prepare a 600 ppm available chlorine sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 12.8 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours. Prior to using equipment, rinse all surfaces with a 200 ppm available chlorine solution. Prepare a 200 ppm sanitizing solution by thoroughly mixing 4.25 oz. of this product with 10 gallons of water.

SANITIZATION OF NONPOROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 4.25 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 4.25 oz. of this product with 10 gallons of water to provide approximately 200 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY/FOG METHOD - Preclean all surfaces after use. Prepare a 200 ppm. available chlorine sanitizing solution of sufficient size by thoroughly mixing the product in a ratio of 4.25 oz. product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Prior to using equipment, thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

[The following directions, "Disinfection of Nonporous Non-food Contact Surfaces," are not to be used on labels that include the claims on page 3 or the disinfection directions on pages 6, 7, 8, or 9.]

DISINFECTION OF NONPOROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a disinfecting solution by thoroughly mixing 12.8 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean equipment surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the disinfecting solution, maintaining contact with the solution for at least 10 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD - Prepare a disinfecting solution by thoroughly mixing, in an immersion tank, 12.8 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the disinfecting solution for at least 10 minutes and allow the disinfectant to drain. Do not rinse equipment with water after treatment.

SANITIZATION OF POROUS NON-FOOD CONTACT SURFACES

RINSE METHOD - Prepare a sanitizing solution by thoroughly mixing 12.8 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean surfaces in the normal manner. Prior to use, rinse all surfaces thoroughly with the sanitizing solution, maintaining contact with the sanitizer for at least 2 minutes. Do not rinse equipment with water after treatment and do not soak equipment overnight.

IMMERSION METHOD - Prepare a sanitizing solution by thoroughly mixing, in an immersion tank, 12.8 oz. of this product with 10 gallons of water to provide approximately 600 ppm available chlorine by weight. Clean equipment in the normal manner. Prior to use, immerse equipment in the sanitizing solution for at least 2 minutes and allow the sanitizer to drain. Do not rinse equipment with water after treatment.

SPRAY/FOG METHOD - After cleaning, sanitize non-food contact surfaces with 600 ppm available chlorine by thoroughly mixing the product in a ratio of 12.8 oz. of this product with 10 gallons of water. Use spray or fogging equipment which can resist hypochlorite solutions. Always empty and rinse spray/fog equipment with potable water after use. Prior to using equipment, thoroughly spray or fog all surfaces until wet, allowing excess sanitizer to drain. Vacate area for at least 2 hours.

SEWAGE & WASTEWATER EFFLUENT TREATMENT

The disinfection of sewage effluent must be evaluated by determining the total number of coliform bacteria and/or fecal coliform bacteria, as determined by the Most Probable Number (MPN) procedure, of the chlorinated effluent has been reduced to or below the maximum permitted by the controlling regulatory jurisdiction.

On the average, satisfactory disinfection of secondary wastewater effluent can be obtained when the chlorine residual is 0.5 ppm after 15 minutes contact. Although the chlorine residual is the critical factor in disinfection, the importance of correlating chlorine residual with bacterial kill must be emphasized. The MPN of the effluent, which is directly related to the water quality standards requirements, should be the final and primary standard and the chlorine residual should be considered an operating standard valid only to the extent verified by the coliform quality of the effluent.

The following are critical factors affecting wastewater disinfection.

1. **Mixing:** It is imperative that the product and the wastewater be instantaneously and completely flash mixed to assure reaction with every chemically active soluble and particulate component of the wastewater.
2. **Contacting:** Upon flash mixing, the flow through the system must be maintained.
3. **Dosage/Residual Control:** Successful disinfection is extremely dependant on response to fluctuating chlorine demand to maintain a predetermined, desirable chlorine level. Secondary effluent should contain 0.2 to 1.0 ppm chlorine residual after a 15 to 30 minute contact time. A reasonable average of residual chlorine is 0.5 ppm after 15 minutes contact time.

SEWAGE AND WASTEWATER TREATMENT

EFFLUENT SLIME CONTROL - Apply a 100 to 1000 ppm available chlorine solution at a location which will allow complete mixing. Prepare this solution by mixing 21.3 to 213 oz. of this product with 100 gallons of water. Once control is evident, apply a 15 ppm available chlorine solution. Prepare this solution by mixing 3.2 oz. of this product with 100 gallons of water.

FILTER BEDS - SLIME CONTROL: Remove filter from service, drain to a depth of 1 ft. above filter sand, and add 170 oz. of product per 20 sq/ft evenly over the surface. Wait 30 minutes before draining water to a level that is even with the top of the filter. Wait for 4 to 6 hours before completely draining and backwashing filter.

DISINFECTION OF DRINKING WATER (EMERGENCY/PUBLIC/INDIVIDUAL/SYSTEMS)

PUBLIC SYSTEMS - Mix a ratio of 2 oz. of this product to 100 gallons of water. Begin feeding this solution with a hypochlorinator until a free available chlorine residual of at least 0.2 ppm and no more than 0.6 ppm is attained throughout the distribution system. Check water frequently with a chlorine test kit. Bacteriological sampling must be conducted at a frequency no less than that prescribed by the National Interim Primary Drinking Water Regulations. Contact your local Health Department for further details.

INDIVIDUAL SYSTEMS: DUG WELLS - Upon completion of the casing (lining), wash the interior of the casing (lining) with a 100 ppm available chlorine solution using a stiff brush. This solution can be made by thoroughly mixing 2.1 oz. of this product into 10 gallons of water. After covering the well, pour the sanitizing solution into the well through both the pipesleeve opening and the pipeline. Wash the exterior of the pump cylinder also with the sanitizing solution. Start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours flush well until all traces of chlorine have been removed from the water. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS: DRILLED, DRIVEN & BORED WELLS - Run pump until water is as free from turbidity as possible. Pour a 100 ppm available chlorine sanitizing solution into the well. This solution can be made by thoroughly mixing 2.1 oz. of this product into 10 gallons of water. Add 5 to 10 gallons of clean, chlorinated water to the well in order to force the sanitizer into the rock formation. Wash the exterior of the pump cylinder with the sanitizer. Drop pipeline into the well, start pump and pump water until strong odor of chlorine in water is noted. Stop pump and wait at least 24 hours. After 24 hours, flush well until all traces of chlorine have been removed from the water. Deep wells with high water levels may necessitate the use of special methods for introduction of the sanitizer to the well. Consult your local Health Department for further details.

INDIVIDUAL WATER SYSTEMS: FLOWING ARTESIAN WELLS - Artesian Wells generally do not require disinfection. If analyses indicate persistent contamination, the well should be disinfected. Consult your local Health Department for further details.

EMERGENCY DISINFECTION - When boiling of water for 1 minute is not practical, water can be made potable by using this product. Prior to addition of the sanitizer, remove all suspended material by filtration or by allowing it to settle to the bottom. Decant the clarified, contaminated water to a clean container and add 3 drops of this product to 20 gallons of water. Allow the treated water to stand for 30 minutes. Properly treated water should have a slight chlorine odor, if not, repeat dosage and allow the water to stand an additional 15 minutes. The treated water can then be made palatable by pouring it between clean containers for several times. This process has not been demonstrated to inactivate Cryptosporidium cysts.

PUBLIC WATER SYSTEMS

RESERVOIRS: ALGAE CONTROL - Hypo chlorinate streams feeding the reservoir. Suitable feeding points should be selected on each stream at least 50 yards upstream from the points of entry into the reservoir.

MAINS - Thoroughly flush section to be sanitized by discharging from hydrants. Permit a water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

NEW TANKS, BASINS, ETC. - Remove all physical soil from surfaces. Place 43 oz. of this product for each 5 cubic feet of working capacity (500 ppm available chlorine). Fill to working capacity and allow to stand for at least 4 hours. Drain and flush with potable water and place in service.

NEW FILTER SAND - Apply 170 oz. of this product for each 150 to 200 cubic feet of sand. The action of the product dissolving as the water passes through the bed will aid in sanitizing the new sand.

NEW WELLS - Flush the casing with a 50 ppm available chlorine solution of water containing 10.7 oz. of this product for each 100 gallons of water. The solution should be pumped or fed by gravity into the well after thorough mixing with agitation. The well should stand for several hours or overnight under chlorination. It may then be pumped until a representative raw water sample is obtained. Bacterial examination of the water will indicate whether further treatment is necessary.

EXISTING EQUIPMENT - Remove equipment from service, thoroughly clean surfaces of all physical soil. Sanitize by placing 45 oz. of this product for each 5 cubic feet capacity (approximately 500 ppm available chlorine). Fill to working capacity and let stand at least 4 hours. Drain and place in service. If the previous treatment is not practical, surfaces may be sprayed with a solution containing 10.7 oz. of this product for each 5 gallons of water (approximately 1000 ppm available chlorine). After drying, flush with water and return to service.

EMERGENCY DISINFECTION AFTER FLOODS

WELLS - Thoroughly flush contaminated casing with a 500 ppm available chlorine solution. Prepare this solution by mixing 10.7 oz. of this product with 10 gallons of water. Backwash the well to increase yield and reduce turbidity, adding sufficient chlorinating solution to the backwash to produce a 10 ppm available chlorine residual, as determined by a chlorine test kit. After the turbidity has been reduced and the casing has been treated, add sufficient chlorinating solution to produce a 50 ppm available chlorine residual. Agitate the well water for several hours and take a representative water sample. Retreat well if water samples are biologically unacceptable.

RESERVOIRS - In case of contamination by overflowing streams, establish hypochlorinating stations upstream of the reservoir. Chlorinate the inlet water until the entire reservoir obtains a 0.2 ppm available chlorine residual, as determined by a suitable chlorine test kit. In case of contamination from surface drainage, apply sufficient product directly to the reservoir to obtain a 0.2 ppm available chlorine residual in all parts of the reservoir.

BASINS, TANKS, FLUMES, ETC. - Thoroughly clean all equipment, then apply 43 oz. of product per 5 cu. ft. of water to obtain 500 ppm available chlorine, as determined by a suitable test kit. After 24 hours, drain, flush, and return to service. If the previous method is not suitable, spray or flush the equipment with a solution containing 10.7 oz. of this product for each 5 gallons of water (1000 ppm available chlorine). Allow to stand for 2 to 4 hours, flush and return to service.

FILTERS - When the sand filter needs replacement, apply 170 oz. of this product for each 150 to 200 cubic feet of sand. When the filter is severely contaminated, additional product should be distributed over the surface at the rate of 100 oz. per 20 sq.ft. Water should stand at a depth of 1 foot above the surface of the filter bed for 4 to 24 hours. When filter beds can be backwashed of mud and silt, apply 170 oz. of this product per each 50 sq. ft., allowing the

water to stand at a depth of 1 foot above the filter sand. After 30 minutes, drain water to the level of the filter. After 4 to 6 hours, drain, and proceed with normal backwashing.

DISTRIBUTION SYSTEM - Flush repaired or replaced section with water. Establish a hypochlorinating station and apply sufficient product until a consistent available chlorine residual of at least 10 ppm remains after 24 hour retention time. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER FIRES

CROSS CONNECTIONS OF EMERGENCY CONNECTIONS - Hypochlorination or gravity feed equipment should be set up near the intake of the untreated water supply. Apply sufficient product to give a chlorine residual of at least 0.1 to 0.2 ppm at the point where the untreated supply enters the regular distribution system. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER DROUGHTS

SUPPLEMENTARY WATER SUPPLIES - Gravity or mechanical hypochlorite feeders should be set up on a supplementary line to dose the water to a minimum chlorine residual of 0.2 ppm after a 20 minute contact time. Use a chlorine test kit.

WATER SHIPPED IN BY TANKS, TANK CARS, TRUCKS, ETC. - Thoroughly clean all containers and equipment. Spray a 500 ppm available chlorine solution and rinse with potable water after 5 minutes. This solution is made by mixing 10.7 oz. of this product for each 10 gallons of water. During the filling of the containers, dose with sufficient amounts of this product to provide at least a 0.2 ppm chlorine residual. Use a chlorine test kit.

EMERGENCY DISINFECTION AFTER MAIN BREAKS

MAINS - Before assembly of the repaired section, flush out mud and soil. Permit a water flow of at least 2.5 feet per minute to continue under pressure while injecting this product by means of a hypochlorinator. Stop water flow when a chlorine residual of test of 50 ppm is obtained at the low pressure end of the new main section after a 24 hour retention time. When chlorination is completed, the system must be flushed free of all heavily chlorinated water.

COOLING TOWER/EVAPORATIVE CONDENSER WATER

SLUG FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 21.3 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

INTERMITTENT FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

Subsequent Dose: When microbial control is evident, add 21.3 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

CONTINUOUS FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 2.1 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

LAUNDRY SANITIZERS

Household Laundry Sanitizers

IN SOAKING SUDS - Thoroughly mix 4.25 oz. of this product to 10 gallons of wash water to provide 200 ppm available chlorine. Wait 5 minutes, then add soap or detergent. Immerse laundry for at least 11 minutes prior to starting the wash/rinse cycle.

IN WASHING SUDS - Thoroughly mix 4.25 oz. of this product to 10 gallons of wash water containing clothes to provide 200 ppm available chlorine. Wait 5 minutes, then add soap or detergent and start the wash/rinse cycle.

Commercial Laundry Sanitizers

Wet fabrics or clothes should be spun dry prior to sanitization. Thoroughly mix 4.25 oz. of this product with 10 gallons of water to yield 200 ppm available chlorine. Promptly after mixing the sanitizer, add the solution into the prewash prior to washing fabrics/clothes in the regular wash cycle with a good detergent. Test the level of available chlorine, if solution has been allowed to stand. Add more of this product if the available chlorine level has dropped below 200 ppm.

FARM PREMISES

Remove all animals, poultry, and feed from premises, vehicles, and enclosures. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities occupied or transversed by animals or poultry. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. To disinfect, saturate all surfaces with a solution of at least 1000 ppm available chlorine for a period of 10 minutes. A 1000 ppm solution can be made by thoroughly mixing 21.3 oz. of this product with 10 gallons of water. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals or poultry, as well as the cleaned forks, shovels, and scrapers used for removing litter and manure. Ventilate buildings, cars, boats, and other closed spaces. Do not house livestock or poultry or employ equipment until chlorine has been dissipated. All treated feed racks, mangers, troughs, automatic feeders, fountains, and waterers must be rinsed with potable water before reuse.

PULP AND PAPER MILL PROCESS WATER SYSTEMS

SLUG FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain from 5 to 10 ppm available chlorine. Repeat until control is achieved.

Subsequent Dose: When microbial control is evident, add 21.3 oz. of this product per 10,000 gallons of water in the system daily, or as needed to maintain control and keep the chlorine residual at 1 ppm. Badly fouled systems must be cleaned before treatment is begun.

INTERMITTENT FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown.

Subsequent Dose: When microbial control is evident, add 21.3 oz. of this product per 10,000 gallons of water in the system to obtain a 1 ppm residual. Apply half (or 1/3, 1/4, or 1/5) of this initial dose when half (or 1/3, 1/4, or 1/5) of the water in the system has been lost by blowdown. Badly fouled systems must be cleaned before treatment is begun.

CONTINUOUS FEED METHOD - Initial Dose: When system is noticeably fouled, apply 107 to 213 oz. of this product per 10,000 gallons of water in the system to obtain 5 to 10 ppm available chlorine.

Subsequent Dose: Maintain this treatment level by starting a continuous feed of 2.1 oz. of this product per 1,000 gallons of water lost by blowdown to maintain a 1 ppm residual. Badly fouled systems must be cleaned before treatment is begun.

AGRICULTURAL USES

POST-HARVEST PROTECTION - Potatoes can be sanitized after cleaning and prior to storage by spraying with a sanitizing solution at a level of 1 gallon of sanitizing solution per ton of potatoes. Thoroughly mix 2.1 oz. of this product to 2 gallons of water to obtain 500 ppm available chlorine.

Disinfect leafcutting bee cells and bee boards by immersion in a solution containing 1 ppm available chlorine for 3 minutes. Allow cells to drain for 2 minutes and dry for 4 to 5 hours or until no chlorine odor can be detected. This solution is made by thoroughly mixing 2 tsp. of this product to 100 gallons of water. The bee domicile is disinfected by spraying with a 0.1 ppm solution until all surfaces are thoroughly wet. Allow the domicile to dry until all chlorine odor has dissipated. [Not for use in California]

FOOD EGG SANITIZATION - Thoroughly clean all eggs. Thoroughly mix 4.25 oz. of this product with 10 gallons of warm water to produce a 200 ppm available chlorine solution. The sanitizer temperature should not exceed 130 degrees F. Spray the warm sanitizer so that the eggs are thoroughly wetted. Allow the eggs to thoroughly dry before casing or breaking. Do not apply a potable water rinse. The solution should not be re-used to sanitize eggs.

FRUIT & VEGETABLE WASHING - Thoroughly clean all fruits and vegetables in a wash tank. Thoroughly mix 10.7 oz. of this product in 200 gallons of water to make a sanitizing solution of 25 ppm available chlorine. After draining the tank, submerge fruit or vegetables for 2 minutes in a second wash tank containing the recirculating sanitizing solution. Spray rinse vegetables with the sanitizing solution prior to packaging. Rinse fruit with potable water only prior to packaging.

AQUACULTURAL USES

FISH PONDS - Remove fish from ponds prior to treatment. Thoroughly mix 213 oz. of this product to 10,000 gallons of water to obtain 10 ppm available chlorine. Add more product to the water if the available chlorine level is below 1 ppm after 5 minutes. Return fish to pond after the available chlorine level reaches zero.

FISH POND EQUIPMENT - Thoroughly clean all equipment prior to treatment. Thoroughly mix 4.25 oz. of this product to 10 gallons of water to obtain 200 ppm available chlorine. Porous equipment should soak for one hour.

MAINE LOBSTER PONDS - Remove lobsters, seaweed, etc. from ponds prior to treatment. Drain the pond. Thoroughly mix 12,800 oz. of this product to 10,000 gallons of water to obtain at least 600 ppm available chlorine. Apply so that all barrows, gates, rocks and dams are treated with product. Permit high tide to fill the pond and then close the gates. Allow water to stand for 2 to 3 days until the available chlorine level reaches zero. Open and allow 2 tidal cycles to flush the pond before returning lobsters to the pond. [NOT FOR USE IN CALIFORNIA]

CONDITIONING LIVE OYSTERS - Thoroughly mix 10.7 oz. of this product to 10,000 gallons of water at 50 to 70 degrees F to obtain 0.5 ppm available chlorine. Expose oysters to this solution for at least 15 minutes, monitoring the available chlorine level so that it does not fall below 0.05 ppm. Repeat entire process if the available chlorine level drops below 0.05 ppm or the temperature falls below 50 degrees F. [Not for use in California]

CONTROL OF SCAVENGERS IN FISH HATCHERY PONDS - Prepare a solution containing 200 ppm of available chlorine by mixing 4.25 oz. of product with 10 gallons of water. Pour into drained pond potholes. Repeat if necessary. Do not put desirable fish back into refilled ponds until chlorine residual has dropped to 0 ppm, as determined by a test kit.

SANITIZATION OF DIALYSIS MACHINES

Flush equipment thoroughly with water prior to using this product. Thoroughly mix 12.8 oz. of this product to 10 gallons of water to obtain at least 600 ppm available chlorine. Immediately use this product in the hemodialysate system allowing for a minimum contact time of 15 minutes at 20 degrees C. Drain system of the sanitizing solution and thoroughly rinse with water. Discard and DO NOT reuse the spent sanitizer. Rinsate must be monitored with a suitable test kit to ensure that no available chlorine remains in the system.

This product is recommended for decontaminating single and multipatient hemodialysate systems. This product has been shown to be an effective disinfectant (*virucide, fungicide, bactericide, pseudomonicide) when tested by AOAC and EPA test methods. This product may not totally eliminate all vegetative microorganisms in hemodialysate delivery systems due to their construction and/or assembly, but can be relied upon to reduce the number of microorganisms to acceptable levels when used as directed. This product should be used in a disinfectant program which includes bacteriological monitoring of the hemodialysate delivery system. This product is NOT recommended for use in hemodialysate or reverse osmosis (RO) membranes.

Consult the guidelines for hemodialysate systems which are available from the Hepatitis Laboratories, CDC, Phoenix, AR 85021.

ASPHALT OR WOOD ROOFS AND SIDINGS

To control fungus and mildew, first remove all physical soil by brushing and hosing with clean water, and apply a 5000 ppm available chlorine solution. Mix 10.7 oz. of this product per gallon of water and brush or spray roof or siding. After 30 minutes, rinse by hosing with clean water. [Not for Use in California]

BOAT BOTTOMS

To control slime on boat bottoms, sling a plastic tarp under boat, retaining enough water to cover the fouled bottom area, but not allowing water to enter enclosed area. This envelope should contain approximately 500 gallons of water for a 14 foot boat. Add 37.3 oz. of this product to this water to obtain a 35 ppm available chlorine concentration. Leave immersed for 8 to 12 hours. Repeat if necessary. Do not discharge the solution until the free chlorine level has dropped to 0 ppm, as determined by a swimming pool test kit. [Not for Use in California]

ARTIFICIAL SAND BEACHES

To sanitize the sand, spray a 500 ppm available chlorine solution containing 10.7 oz. of this product per 10 gal. of water at frequent intervals. Small areas can be sprinkled with a watering can. [Not for Use in California]

WAREWASHING

FOR SANITIZING TABLEWARE IN LOW TEMPERATURE DISHWASHING MACHINE -
Dispense this product into final rinse water at 100 ppm available chlorine. Do not allow concentration to fall below 50 ppm. Air dry. Dispenser should be set to deliver 6.5 cc of sanitizing solution per gallon of water to give 100 ppm of available chlorine. Only a qualified service representative should set or adjust dispenser on the machine.

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST *HIV (AIDS VIRUS) ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Kills *HIV-1 (AIDS virus) on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in healthcare settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of *Human Immunodeficiency Virus type 1 (HIV-1) (associated with AIDS).

Personal Protection: Disposable latex or vinyl gloves, gowns, masks, and/or eye coverings as appropriate must be worn during all cleaning and decontamination procedures of blood and other body fluids.

Cleaning Procedures: Blood and other body fluids must be thoroughly cleaned from surfaces and objects before applying this product.

Disinfectant Use and Contact Time: Effective against *HIV-1 (AIDS virus) on hard non-porous surfaces in the presence of a moderate amount of organic soil (e.g. 5% blood serum). Prepare disinfectant by mixing 21 oz. of this product per gallon of water to provide 10,000 ppm of available chlorine. Leave surfaces wet for 5 minutes. Drain and let air dry.

Disposal of Infectious Materials: Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to preclean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.