Master Label

Page 1 of 9

ADDITIONAL BRAND NAMES: ECOSIDE & ECOSENSE

Agroneem®

Sub-label A: For Agriculture, Horticulture & Greenhouses

Sub-label B: For Home Gardens

Sub-label C: For Golf Courses, Lawns, Playgrounds,

Parks, & Ornamentals

Emulsifiable Concentrate

AGRICULTURAL, HORTICULTURAL, GREENHOUSE, AND OUTDOOR DOMESTIC TERRESTRIAL PESTICIDE

ACTIVE INGREDIENT:

Azadirachtln

0.15%

NEEM EXTRACT:

15.00%

OTHER INGREDIENTS:

84.85%

TOTAL:

100.00%

(4.8 GRAMS/GALLON ACTIVE INGREDIENT)

KEEP OUT OF REACH OF CHILDREN

CAUTION


EPA REGISTRATION No. 70310-1 EPA ESTABLISHMENT No. 070310-IND-002

> Agro Logistic Systems, Inc. P.O. Box 5799 Diamond Bar, CA 91765 U.S.A.

TEL: (800) 425-3388 Website: www.agrologistic.com

Net contents: 8 Fl. Oz to 2 1/2 Gal.





Under the Federal Insecticide. Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin or if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or dothing. Avoid breathing spray. Remove contaminated dothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to inter-tidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

Batch No:

Manufacturing Date:

Expiry Date:

PRODUCT OF INDIA

Mfd. by:

Bahar Agrochem and Feeds PVt. Ltd Lote Parshuram, INDIA

MANUFACTURED FOR

Agro Logistic Systems, Inc. P.O. Box 5799 Diamond Bar, CA 91765 U.S.A.

TEL: (800) 425-3388 Website: www.agrologistic.com

EPA REGISTRATION No. 70310-1 EPA ESTABLISHMENT No. 070310-IND-002

Sub-label A: Agriculture, Horticulture, & Greenhouse use ADDITIONAL BRAND NAMES: ECOSIDE& ECOSENSE

Agroneem®

Emulsifiable Concentrate

ACTIVE INGREDIENT:

Azadirachtin

0.15%

NEEM EXTRACT:

15.00%

OTHER INGREDIENTS:

84.85%

TOTAL:

100.00%

(4.8 GRAMS/GALLON ACTIVE INGREDIENT)

Plant Origin
Antifeedant
Insect Repellent
Insecticide
Nematicide

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID		
ti swaliowesi	- Call a polition control center or doctor intensitately for treatment advice - Have person sip a glass of water if able to swallow On not induce vorniting unless told to do so by a poison control center or doctor. - Do not give anything by mouth to an unconscious person.	
If an skin or classing	Take off contaminated dicthing: Rinse skin immediately with plenty of water for 15-20 minutes Call a poison control center or doctor for breatment advice.	
lf inhaled	Hove person to fresh air, If person is not breathing call 911 or an ambutance, then give artificial respiration, preferably mouth to mouth if possible. Call a poison control center or doctor for further treatment advice.	
) ที่กฤห	- Hold eye open and rinse slowly and gesitly with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. - Call postun control center or doctor for treatment advice.	

Net contents: 8 Fl. Oz to 2 1/2 Gal.

poison control center or doctor, or going for treatment".

DIRECTIONS FOR USE

Page 2 of 9

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State and Tribe, consult the agency responsible for pesticide regulation.

Refer to the attached booklet for the listing of crops and pests treated, dilution directions and application rates.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, potification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personnel protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into breated areas during the

Do not enter or allow worker entry into breated areas during the restricted entry interval (REI) of 4 hours. PPE required for early entry into treated areas that is permitted under the Worker Protection Standard and that throwes contact with anything that has been treated such as plants, soil or water is:

Coveralls

Chemical resistant gloves, such as barrier laminates or butyl rubber or nitrile rubber or PVC.

Shoes and socks

NON- AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used on agricultural plants on farms, forests, nurseries, or greenhouses.

STORAGE AND DISPOSAL

On not contaminate water, food, or feed by storage and disposal, PESTICIDE STORAGE: Keep in original packaging in cool and dry place away from sunlight and foodstuffs. Do not refrigerate. For Commercial Agricultural Use:

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple Rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smake.

J.

DIRECTIONS FOR CROP APPLICATION

ATION Page 3 of 9

PREVENTATIVE APPLICATION: APPLICATION RATES ARE FOR ONE ACRE EXCEPT AS NOTED.

As a preventative application mix 32 oz. to 48 oz. of AGRONEEM in 50 - 100 gallons water or, sufficient amount of water to ensure total foliar coverage. Start applying when the plants are tail enough to be susceptible to infestation. Use the following:

- For Baby Vegetables, apply 10 days after planting and repeat again in 7- 10 days.
- . For Regular Vegetables, apply when the plant has grown two leaves or is 6 inches tall. Repeat application every 10 to15 days.
- For Fruit Trees, apply when the tree ends dormancy and continue to repeat applications every 10 15 days.

 (For fruit trees use 64 oz. AGRONEEM i.e. two quarts in 150 gallons of water)
- For Berries, apply when the plant ends dormancy and continue to repeat applications every 10 15 days. (For berries use 64 oz. AGRONEEM i.e. two quarts in 150 gallons of water).
- For Grasses and Forage Crops like Alfalfa, apply in spring, approximately 2 weeks after dormancy period is over, and repeat 10-15 days.

STANDARD APPLICATION: ALL APPLICATION RATES ARE FOR ONE (1) ACRE PER CROP

48 oz. AGRONEEM (1-1/2 quarts) mixed with 100 gallons water or sufficient amount of water to ensure total foliar coverage. When applying at early plant stage use dilution of 48 oz. AGRONEEM (1-1/2 quarts) with 50 gallons of water. When applying in high population crops like berries and fruit trees, use dilution of 64 oz. to 128 oz of AGRONEEM (2- 4 quarts), with 150 – 300 gallons of water.

Repeat application at 7 to 10 day intervals, at least once and if necessary additional times.

CROPS	INSECT PESTS/MITES
Top fruits such as Citrus, Lemons, Oranges, Grapefruits, Pummaio,	White Files, Mealybugs, Leafminers, Thrips, Waevils, Mites, Psylia
Apples, Plums, Peaches, Apricots, Avocados, Figs, Prunes, Nectarines,	Grasshoppers, Fruit Files, Aphids, San Jose Scale, H Scale, Orange Tortrix,
Pears, Mangoes, Chemies, Kiwifruit, Kumquats, Loquats, Quinces,	Codling Moth, Nematodes, Beetles, Pacific & Two Spotted Spider Mite, Leaf
Persimmons, and Palm.	rollers, Leafhoppers, Borers, Maggots, Leaf miner, and Sharpshooters.
Cucurbits, Watermelons, Gherkins, Squashes, Pumpkins, Gourds, Bitter	White Files, Budworms, Leafminers, Fell Armyworms, Pickle Worms, Cucumber
melans, Chayote, Okra, Asparagus, and Melans,	Beetles, Thrips, Aphilds, Nematodes, Beetles and Japanese Beetles.
Builb, Cole and Leafy vegetables such as Broccoll, Brussels sprouts,	Aphids, White Files, Cabbage Loopers, Diamondback Moths,
Cabbage, Celery, Lettuco, Endives, Kale, Parsley, Onions, Garlic,	Armyworms, Thrips, Leafminers, Onion Maggots, Weevils, Caterpillars, Whiteflies,
Shallots, Leeks, Kohirabi, Chard, Collard, Amaranth, and Spinach.	Nematodes and Grasshoppers.
Root and Tuber vegetables such as Beets, Carrots, Ginger, Radish,	Colorado Potato Beetle, Leaf hopper, Potato Fiea Beetles,
Horseradish, Potatoes, Turnips, Yams, Sweet Potatoes, Ginseng,	Aphids, White Flies, Psyllids, Nematodes, Armyworms, Thrips, Beetles, and
Rutabagas, Watercress, Artichokes and Turmeric.	Webworms.
Legumes such as Beans, Lentils, Peas, Peanuts, Soybeans and Garbanzo	Loopers, Learminers, Caterpillors, Psyllids, Weevils, Beanleaf Beetles, Mexicon
Beans.	Beetles, Thrips, Aphids, Nematodes, and Colorado Potato Boetles.
Cereals such as Barley, Corn, Millet, Rice, Rye, Sorghum, Triticale, Oats,	Weavils, Maggots, Wireworms, Corn Beetles, Flea Beetles, Thrips, Aphids,
and Wheat.	Cutworms, Barers, Armyworms, Rootworms, Leaf miners, Leafhappers,
Alfalfa and other feed & forage crops. Grasses	Caterpillars, Grasshoppers, Mites, Nematodes, Whiteflies, and Japanese beetles.
Fruiting vegetables such as Tomatoes, Tomatillos, Bell Peppers, Chili	Loopers, Leafminers, Cabbage Loopers, Catarpillars, Pepper, Nematodes
Peppers, and Eggplant.	Weevils, Aphids, Cucumber Beetles, Thrips, Whiteflies, and Grasshoppers
Berries such as, Blackberries, Blueberries, Currants, Strawberries, Cone	Gypsy Moths, Loopers, Grape Berry Moths, Leaf rollers, Fruit files,
berries, Gooseberries, Raspberries, Cranberries, Loganberries,	Thrips, Caterpillars, Tobacco Budworms, Nematodes, Alfalfa worm, Aphids,
Huckleberries, and Boysenberries	Phylloxers, Whiteflies, maggots, and Lygus Bug.
Grapes	Gypsy Moths, Loopers, Grape Berry Moths, Leaf rollers, Fruit files, Weevil,
Tropical fruits such as Banana, Cherimoyas, Mangoes, Guavas, Papayas	Thrips, Caterpillars, Tobacco Budworms, Aphilds, Alfalfa worm, Eelworm
and Pineappie.	Phylioxera, Mealy bugs, Whiteffles, Nematodes, Hoppers, and Sharpshooters.
Cotton and oilseeds such as Canola, Sesame, Rapeseed, Sunflower,	White Files, Fruit files, Thrips, Ballwarms, Leafininers, Ball Weevil, Grubs, Loopers,
Safflower, Olive, Peanut and Jojoba	Wireworms, Barers, Spider mites, Leaf perforators, Leafhoppers, Armyworms,
	Earworms, Nematodes, Caterpillars, Loopers, Cutworms, and Aphids.
Herbs & Spices including but not limited to Mints, Anise, Basil, Chives,	Aphids, Flea Beetles, Thrips, Nematodes, Whiteflies, and Leafhoppers
Coriander, Dill, Fennel, Markgold, Sage, and Thyme.	
Nuts such as Almonds, Pecans, Pistachios, Cashews, Chestnuts,	Casebearer, Aphids, Phylloxera, Leaf rollers, Weevils, Thrips, Whiteflies, Moths,
Macadamia, Hazelnuts, Brezil nut, Water chestnuts, and Walnuts.	Suckers Ants, Nematodes, Mealy bugs, Miles, Scales, Fagot worm, Bugs,
Coffee, Tea, Tobacco and Sugarcane	Caterplilars, Tobacco Budworms, Beedes, Grasshoppers, and Leafhoppers
Greenhouse Crops including ornamentals such as Chrysanthemums,	Aphids, Thrips, Leaf miners, Whitefiles, Nematodes, Fungus gnats and other
Poinsettias, Roses, Lilies, Geraniums, Daisles, Carnations, Salvias and	greenhouse pests.
Dahlias.	Prepare a 1% - 2% (1.25 oz - 2.5 dz./ Gal) solution of Agronaem in water and
All starter vegetables and fruit grops.	completely drench the plants. Repeat every 5 to 7 days till infestation clears.

FOR NEMATODE CONTROL, USE 2-4 GALLONS AGRONEEM IN 500 -1500 GALLONS WATER PER HALF ACRE AS A DRENCH OR DRIP.

Agroneem

Page 4 of 9

PLEASE READ THIS BOOKLET CAREFULLY BEFORE USING AGRONEEM

TANK MIXING:

'AGRONEEM' is compatible with commonly used insecticides, fungicides and foliar fertilizers, ilke all herbal products. Do not use Agroneem with Bordeaux mixture, lime sulfur or other highly alkaline materials.

It is advisable to pre-test the tank mix combination proposed on a few plants to observe phytotoxicity, if any. The climatic conditions and cultural practices may be different in the actual field than experimental conditions. Therefore the responsibility of damage due to tank mix, if any, remains with the user. This product is damaged by oxidants. Do not mix acids and bleaches with AGRONEEM.

MIXING DIRECTIONS:

- 1. Determine the acreage to be treated with AGRONEEM.
- 2. Fill tank with appropriate amount of water (see inside for application rates).
- 3. Add the pre-measured amount of AGRONEEM (see Inside for amount to be used).
- 4. Keep tank agitated with an agitation device. Proper agitation is required, especially when mixing AGRONEEM with other pesticides.

SPRAYING DIRECTIONS:

- Spray AGRONEEM using a boom system, propeller turbine system (indicated for orchards); pressure hose with gun sprayers, powered back sprayers and manual hand sprayers. Use a minimum boom pressure of 160 psi and maximum tractor speed of 2.0 miles per hour. Use horizontal nozzles at 10- inch spacing and two vertical drops, one at each end. Each vertical drop must have two nozzles, one at 90° angle and the other at 45° angle. The water used for mixing should have a pH at about 5.5 for AGRONEEM to be most effective. If necessary, use a buffer to bring the pH to the desired level.
- 2. DURING APPLICATION, MAKE SURE THAT THE SPRAY COVERS THE ENTIRE PLANT INCLUDING THE <u>UNDERSIDE OF LEAVES</u>.

GENERAL:

- AGRONEEM is more effective when used as a preventative.
- AGRONEEM is a broad-spectrum herbal insecticide.
- AGRONEEM affects the Insects at all stages of growth.
- SHAKE THE CONTAINER WELL, BEFORE USE.

MAXIMUM APPLICATION RATE IS 20 GRAMS OF ACTIVE INGREDIENT (530 FL. OZ. OR 16 QUARTS OF AGRONEEM E.C.) OR LESS PER ACRE
ACCORDING TO TOLERENCE EXEMPTION 40 CFR 180,1119

Page 5 of 9

WARRANTY:

AGRO LOGISTIC SYSTEMS, INC. warrants that material contained herein conforms to the description on the label and is reasonably useful for the purposes referred to in the directions for use. Method of application and timing, weather, cultural practices, insect problem, crop condition, and incompatibility with other chemicals not specifically recommended in the use of this product are beyond the control of the seller. It is the manufacturer's intention that the buyer assumes risk of use, storage, or handling of this material not in strict accordance with directions given herein.

FOR ADDITIONAL INFORMATION AND PRODUCT BROCHURE PLEASE CONTACT:

Agro Logistic Systems, Inc. P.O.Box: 5799 Diamond Bar, CA 91765 U.S.A.

TEL: (800) 425-3388 TEL: (909) 594-7222 FAX: (909) 594-7330 e-mail: info@agrologistic.com

website: www.agrologistic.com

Sub-label B: For Home Gardens

Page 6 of 9

For outdoor Use

Organic Insecticide From the Neem Tree

Ideal for Flower Beds, Vegetable Gardens, Lawns & Fruit Trees

Controls many insects including:

Red ants, Cutworms, Grubs, Chinch bugs, Crickets, Aphids, Whitefiles, Thrips, Beetles, Leafhoppers, Fruit worms, Spider mites, Caterpillars, loopers, Armyworms, Grasshoppers, Maggots, Weevils, Mites, spittlebugs, Webworms

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID		
If swallowed	- Call a poison control center or doctor immediately for treatment advice, - Have person signal glass of water if able to swallow.	
	-Do not induce vomiting unless taid to do so by a poison control center or doctor.	
If on sidn or dothing	- Do not give anything by mouth to an unconscious person. - Take off contaminated clothing.	
_	- Rinse skin immediately with plenty of weter for 15-20 minutes - Call a poison control center or decitor for treatment advice.	
If inhaled	- Move person to fresh air If person is not breakhing call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible Call a poison control centur or doctor for further treatment advict.	
If in eyes	- Hold eye open and rurse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, ofter the first 5 minutes, then continue rinsing eye Call poison control center or doctor for treatment advice.	
"Have the produc	t container or label with you when calling a poison control center or doctor, or going for treatment".	

SEE OTHER PANEL FOR PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:

Azadirachtin

0.15%

NEEM EXTRACT:

15.00%

OTHER INGREDIENTS:

84,85%

TOTAL:

100.00%

(4.8 GRAMS/GALLON ACTIVE INGREDIENT)

EPA REGISTRATION No. 70310-1

EPA ESTABLISHMENT No. 070310-IND-002

Agro Logistic Systems, Inc.

P.O. Box 5799

Dlamond Bar, CA 91765, U.S.A.

Tel: (800) 425-3388

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful If absorbed through skin or If Inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to inter-tidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not store near heat or open flame.

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

- 1. SHAKE THIS CONTAINER WELL BEFORE USE.
- 2. MIX 2 TABLESPOONS (OR 30 ml) OF AGRONEEM WITH 32 OZ. WATER IN A SPRAY CONTAINER. SHAKE WELL.
- 3. SPRAY THE MIXTURE ON THE LEAVES AND STEMS, INCLUDING THE UNDERSIDE OF LEAVES SO AS TO COMPLETELY DRENCH THE PLANT.
- 4. SPRAY AGRONEEM EARLY IN THE MORNING OR LATE AFTERNOON.
- 5. USE ALL OF THE DILUTED MATERIAL SOON AFTER MIXING. DO NOT SAVE MIXED MATERIAL.
- 6. FOR BEST RESULTS, APPLY WEEKLY (OR MORE FREQUENTLY IF INFESTATION PERSISTS).
- 7. USE AS A PREVENTATIVE TO KEEP INSECTS AWAY.
- 8. DO NOT WASH THE LEAVES FOR 24 HOURS AFTER SPRAYING AGRONEEM.
- 9. FOR LAWNCARE, APPLY 1.14 OZ (32 ml) OR LESS OF CONCENTRATE PER 100 SQ. FT. PER APPLICATION.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in original packaging in cool and dry place away from sunlight and foodstuffs. Do not refrigerate. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or [1-800-CLEANUP] for disposal instructions. Never place unused product down any indoor or outdoor drain,

WARRANTY STATEMENT

AGRO LOGISTIC SYSTEMS, INC. warrants that material contained herein conforms to the description on the label and is reasonably useful for the purposes referred to in the directions for use. Method of application and timing, weather, cultural practices, insect problem, crop condition, and incompatibility with other chemicals not specifically recommended in the use of this product are beyond the control of the seller. It is the manufacturer's intention that the buyer assumes risk of use, storage, or handling of this material not in strict accordance with directions given herein.

Manufacturing Date:

Expiry Date:

Sub-label C: For Golf Courses, Lawns, Playgrounds, Parks & Ornamentals

NEMATODE, AND TURF INSECT CONTROL

Controls many insects including:

Red ants, Cutworms, Grubs, Chinch bugs, Crickets, Aphids, Whitefiles, Thrips, Beetles, Leafhoppers, Fruit worms, Spider mites, Caterpillars, Cabbage loopers, Armyworms, Colorado potato beetles, Diamondback moths, Grasshoppers, Maggots, Weevils, Chafers, Mites, spittlebugs, Webworms

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID		
If swallowed	- Call a poison control center or doctor immediately for treatment advice Have person signal glass of water if able to swallowDo not induce verniting unless told to do so by a colson control center or doctor Do not give anything by mouth to an unconscious person.	
If an sidn or dollling	-Take off contaminated clothing Rinse sidn immediately with plenty of water for 15-20 minutes - Call a poison control center or doctor for treatment advice.	
If inhaled	- Move person to fresh elv If person is not breathing call 911 or an embuliance, then give extificial respiration, preferably mouth to mouth if possible Call a poison control center or doctor for further treatment advice.	
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call poison control center or doctor for treatment advice.	
"Have the produc	t container or label with you when calling a poison control center or doctor, or going for treatment".	

SEE OTHER PANEL FOR PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:

Azadirachtin

0.15%

NEEM EXTRACT:

15.00%

OTHER INGREDIENTS:

84.85%

TOTAL:

100.00%

(4.8 GRAMS/GALLON ACTIVE INGREDIENT)

EPA REGISTRATION No. 70310-1 EPA ESTABLISHMENT No. 070310-IND-002 Agro Logistic Systems, Inc.

P.O. Box 5799

Diamond Bar, CA 91765, U.S.A.

Tel: (800) 425-3388

Page 9 of 9

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful If absorbed through skin or if Inhaled. Causes moderate eye Irritation. Avoid contact with skin, eyes or clothing. Avoid breathing spray. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to inter-tidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

- 1. SHAKE THIS CONTAINER WELL BEFORE USE.
- 2. A. FOR NEMATODE CONTROL:

MIX 32 FL. OZ OF AGRONEEM WITH 2 TO 30 GALLONS WATER TO SPRAY ON 1000 SQUARE FEET. Using sprinklers, deliver additional amounts of water to push the material to 4 inches below ground level

Apply once every 3 to 6 months. Test the soil every month for nematode count. Repeat application if the nematode count is over the threshold limit.

B. FOR INSECT AND WORM CONTROL (see front panel for list):

MIX 2.5 TO 5.0 FL. OZ. OF AGRONEEM IN 2 TO 3 GALLONS OF WATER (or 1 to 2 gallon of Agroneem with 100 gallons water per acre). Ensure that ph of this mixture is around 5.5 by using a buffer. Spray the mixture on to 1000 square feet of turf. Using sprinklers or other means, deliver additional amounts of water to push the material to $\frac{1}{2}$ inch below ground level. For worm and insect control apply at the earliest sign of infestation. Repeat the application at 7-10 days interval if infestation persists.

- 3. SPRAY AGRONEEM IN THE EARLY MORNING OR LATE AFTERNOON.
- 4. USE ALL OF THE DILUTED MATERIAL SOON AFTER MIXING. DO NOT SAVE MIXED MATERIAL.
- 5. USE AS A PREVENTATIVE TO KEEP INSECTS AWAY.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep in original packaging in cool and dry place away from sunlight and foodstuffs. Do not refrigerate. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly, then securely wrap original container in several layers of newspaper and discard in trash.

WARRANTY STATEMENT

AGRO LOGISTIC SYSTEMS, INC. warrants that material contained herein conforms to the description on the label and is reasonably useful for the purposes referred to in the directions for use. Method of application and timing, weather, cultural practices, insect problem, crop condition, and incompatibility with other chemicals not specifically recommended in the use of this product are beyond the control of the seller. It is the manufacturer's intention that the buyer assumes risk of use, storage, or handling of this material not in strict accordance with directions given herein.

Manufacturing Date:

Explry Date: