

PM 04
70506-2
02-08-79

S 546092
PAGE 1 OF 14



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
70506-2

Date of Issuance:
FEB 8 1999

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
Lancer 90 SP

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):
United Phosphorous, Inc.
c/o John M. Wise, Agent
5 Victory Lane, Suite 201
Liberty, MO 64068

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes before you release the product for shipment:
 - a. Add the phrase, "EPA Reg. No. 70506-2".
 - b. On page 9 under general use precautions, correct the maximum application rate from 8 lbs. of LANCER 90 SP to 6 2/3 lbs. LANCER 90 SP per acre per crop cycle.
 - c. On page 9 in the directions for use against cutworms, correct the sentence, Use 1 lb. per acre of LANCER 75 SP to read: Use .8 lb. per acre of LANCER 90 SP.

Signature of Approving Official:
Michael Meant

Date:
FEB 8 1999

d. On page 2 under the PPE section for applicators and other handlers, add the use of waterproof gloves.

e. Qualify the directions to repeat application as necessary wherever it appears so that it is clear that applications must be limited to a maximum of 6 2/3 lbs. of LANCER 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.

f. Your cover letter of July 8, 1998 described the container packaging as consisting of individual five and twenty pound packages contained in a larger outer container or case. Your letter also stated that these small individual packages may be sold separately.

You are reminded that each of the individual packages contained within the larger outer container must also bear the complete label; not just the directions for use as was mentioned in your cover letter.

3. Submit or cite the following data in accordance with the specifications of the Data Call-In and within the time frames established under the respective DCIs for similar products. These data are the outstanding acephate generic data as of January 19, 1999 which are pertinent to your subject product as proposed for registration.

<u>Guideline #: Study</u>	<u>DCI *</u>
132-1(a);Foliar residue dissip.	C, D, E
132-1(b);Soil residue dissip.	**C
133-3;Dermal passiv dosimetry expo.	C, D, E
165-3;Accumulation;irrig. crop	** (C)
171-4(k);Tobacco	** (C)
171-4(l);Bean cannery residue	** (C ,F)
231;Estimation,dermal expos.	C, D
232;Estimation,inhalation expos.	C, D
233;Estimation,dermal expos.	C
234;Estimation,inhalation expos. mammal	C

* A through E: Refer to the attached DCI listing for acephate
 F: The Acephate Registration Standard of
 September 1987
 ** Waiver request pending with the Agency

page 3
EPA Reg. No. 70506-2

4. Submit two copies of the revised final printed label for both the individual product and the outer container before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

It is understood that the Confidential Statement of Formula(CSF) dated July 6, 1998 replaces all other CSFs submitted for the subject product.

Sincerely,

Marilyn Mautz
Biologist
Insecticide-Rodenticide Branch
Registration Division (7504C)

SR

LANCER 90 SP

Soluble Powder
(INSECTICIDE)

ACTIVE INGREDIENT	By Wt.
Acephate (O,S-Dimethyl acetylphosphoramidothioate)	90%
INERT INGREDIENTS	<u>10%</u>
TOTAL	100%

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist. Wash hands thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT: Acephate is an organophosphate, cholinesterase inhibitor.

If swallowed: drink 1 or 2 glasses of water (or milk) and induce vomiting by touching the back of the throat with finger. If possible contact a physician, Poison Control Center, or emergency center before inducing vomiting. Do not induce vomiting or give anything by mouth to an unconscious person. Take person and product container to the nearest emergency treatment center.

If in eyes: Wash eyes with fresh water for 15 minutes. If irritation continues, See a doctor.

If on skin: Wash skin with plenty of soap and water. If inhaled: Remove person from exposure area.

Note to Physicians: Acephate is a cholinesterase inhibitor. If signs of cholinesterase inhibition appear, atropine is antidotal. 2-PAM may also be used in conjunction with atropine but should not be used alone. Acephate is an organophosphate, cholinesterase inhibitor.

Net Contents: _____ Pounds

EPA Reg. No. 70506-

EPA Est. No. _____

Manufactured for:
United Phosphorus Ltd.
Exton, PA 19341

ACCEPTED
with COMMENTS
in EPA Letter dated:

FEB 8 1999

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, the above pesticide registered under EPA Reg. No. 70506-2

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Applicators and other handlers must wear: long-sleeved shirt and long pants, shoes plus socks and chemical-resistant headgear for overhead exposure.

Mixers and Loaders must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical-resistant headgear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or soil-incorporate spills.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas until restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks and chemical-resistant headgear for overhead exposure.

DIRECTIONS

TANK MIXES

NOTICE: Tank mixing of the product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of the user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

TABLE OF EQUIVALENTS

AMOUNT OF LANCER 90 SP PER ACRE	ACRES TREATED - BY A 5 LB. PACKAGE	ACRES TREATED BY A 20 LB. PACKAGE
2.5 oz.	32	128
3.2 oz.	25	100
3.25 oz.	24.6	98.4
0.25 lb.	20	80
0.28 lb.	17.8	71.4
0.5 lb.	10	40
0.56 lb.	8.9	35.6
0.8 lb.	6.25	25
1.0 lb.	5	20
1.1 lb.	4.5	18.2

GENERAL INFORMATION

Do not use this product through any type of irrigation system.

LANCER 90 SP is an insecticide which controls a wide range of important pests in cotton. The active ingredient is acephate, a water-soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by LANCER 90 SP through ingestion than by contact. LANCER 90 SP can be applied as hopper-box, in-furrow and foliar treatments in cotton. Application of LANCER 90 SP to maintain control should be repeated only as directed.

FAILURE TO FOLLOW THE DIRECTION FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to the product's nature, settling is likely to occur.

PREPARATION OF SPRAY SOLUTION

To prepare spray solution, empty the entire contents of package into a spray tank containing at least one-half of the total quantity of water required. Add remaining water with the agitator running until the spray volume is reached. **Do not add any liquid fertilizers, micronutrients or adjuvants to the spray solution until the LANCER 90 SP has completely dissolved.** LANCER 90 SP should completely dissolve in approximately five minutes. Dissolution rate may be slowed by cold water, lack of agitation, or water containing high concentrations of boron or sulfur.

COTTON

GENERAL USE PRECAUTIONS:

Do not use treated seed for food or feed purposes or process for oil.
 Do not feed gin trash to livestock.
 Do not allow animals to graze on treated areas.
 Do not apply more than 6% lbs. of LANCER 90 SP per acre per crop cycle.

**COTTON SEED HOPPER BOX TREATMENT
 (EXCEPT AZ, CA AND THE BLACK LANDS OF TEXAS)**

INSECTS	AMOUNT LANCER 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS
Black (Greasy) Cutworm Cotton Aphids Thrips	2.5 to 3.25 oz.	<p>Add dry LANCER 90 SP to seed in the planter box to achieve an application rate of 2.5 to 3.25 oz. per acre.</p> <p>Even coverage of seed is required for acceptable insect control. Plants grown from improperly treated seed may not exhibit insect control. To obtain even coverage, fill planter box 1/2 full of seed, add 1/2 of required LANCER 90 SP and mix thoroughly. Add remainder of seed and LANCER 90 SP to the hopper box and thoroughly mix.</p> <p>Improved distribution of LANCER 90 SP on seed can be achieved by premixing both in a separate container and then transferring treated seed to hopper box.</p> <p>Use of LANCER 90 SP with air planter is not recommended since loss of product will occur.</p> <p>Do not use treated seed for food or feed purposes or process for oil.</p>

USE PRECAUTIONS:

- Do not use treated seed for food or feed purposes or process for oil.
- LANCER 90S cotton seed hopper box treatment is not permitted in Arizona, California or the Black Lands of Texas.

COTTON (Continued)

COTTON IN-FURROW APPLICATION

INSECTS	AMOUNT LANCER 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS
<p>Aphids*</p> <p>Thrips</p> <p>Black (Greasy) Cutworm (Except CA)</p> <p>* Excluding Cotton Aphids in AZ & CA</p>	<p>0.55 to 1.1 lbs.</p> <p>NOTE: for the Blacklands of Texas use 1.1 lbs. per acre LANCER 90 SP</p>	<p>Apply LANCER 90 SP in 3 to 5 gals. of water per acre as an in-furrow spray at planting. LANCER 90 SP can be mixed with fungicides that are sprayed in-furrow for disease control.</p> <p>Flat-fan nozzles used for in-furrow application should be set so that the fanned spray pattern is aligned with the row to insure good spray deposition in the seed furrow. Cone type nozzles may not provide a spray pattern that insures maximum spray deposition in the seed furrow, and should be avoided. Spray systems that employ metal or plastic tubing for delivery of spray solution into the seed furrow should have tubing securely fastened to the furrow opener and should be checked frequently to insure that the tubing is properly positioned to deposit spray solution into the seed furrow.</p>

COTTON (Continued)

FOLIAR APPLICATION

INSECTS	AMOUNT LANCER 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Thrips	3.2 oz.	Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA and AZ) or 10 to 25 gals. spray per acre by ground.	21
Plant Bugs (Lygus)	0.25 to 1.0 lbs. 0.5 to 1.0 lbs. (CA and AZ)	Apply when eggs or insects first appear. Repeat as necessary to maintain control.	
Fleahopper	0.25 lb.	Lygus -- Use higher rate for lygus adults that have migrated into cotton.	
Cotton Aphid (excluding AZ and CA)	0.5 to 1.0 lbs.	Cotton Aphid -- This insect may develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current control recommendations.	
Whitefly (excluding Sweetpotato Whitefly/Silver Whitefly)	0.5 to 1.0 lbs.	Amyworms & Cabbage Looper -- Apply when eggs appear and repeat at 3 to 7 day spray intervals.	
Armyworms (excluding Beet Armyworm) Cabbage Looper	1.0 lbs.	Bollworm and Tobacco Budworm -- Early season light infestation use 0.5 lb. per acre. Mid and late season moderate to severe infestations use 0.8 to 1.0 lbs. per acre. Apply when eggs appear and repeat at 3 to 7 day spray interval.	
Bollworm Tobacco Budworm Adults Larvae Eggs (DEADHATCH®)	0.5 to 1.0 lbs. (East of Rockies) 1.0 lbs. (CA & AZ) Emerging Larvae 0.25 to 0.5 lbs.	Adult/Larvae: Moths are controlled when they come in direct contact with spray particles during application. Moth kill is most likely to occur when application is made late in the evenings during periods of peak activity. Emerging Larvae: 0.25 to 0.5 pounds per acre when when larvae first appear DEADHATCH: Control of emerging larvae by consumption of treated egg casings.	
Stinkbugs (AZ & CA)	0.8 lb.	Pink Bollworm -- Apply when insects appear and repeat at a 5 to 7 day spray interval.	
Pink Bollworm (AZ & CA)	1.0 lbs.		

COTTON (Continued)

FOLIAR APPLICATION (Continued)

INSECTS	AMOUNT LANCER 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Cutworms	0.8 lb.	<p>Cutworms -- Ground application is recommended. <i>Aerial applications are less effective, but may be used.</i> Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA and AZ) or 10 to 25 gals. spray per acre by ground. Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant. Apply when insects first appear or damage is first noted and repeat application as necessary to maintain control.</p>	21

COTTON TANK MIXES

GENERAL USE PRECAUTIONS:

Do not feed gin trash or treated forage to livestock.
 Do not allow livestock to graze on treated areas.
 Do not apply more than 8 lbs. of LANCER 90 SP per acre per crop cycle.

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions which appear on all product labels. The most restrictive labeling applies when using a tank mix.

RESISTANCE-MANAGEMENT

Cotton pest control programs, especially those for control of Silverleaf Whitefly populations, should employ a properly designed resistance-management strategy. Such resistance-management strategies include mixture or rotation of alternative classes of chemistry including organo-phosphates, carbamates, pyrethroids or insect growth regulators. Consult your state or area agricultural extension service for local resistance management strategies and advice on alternative insecticides.

INSECTS	AMOUNT LANCER 90 SP + LORSBAN® 4E PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Armyworms (excluding Beet Armyworm) Bollworms Cabbage Looper Cotton Aphid Cutworms Fleahopper Grasshoppers Pink Bollworm Plant Bugs (Lygus, Mirids) Salt Marsh Caterpillar Thrips Tobacco Budworm Whitefly (excluding Sweetpotato/Silverleaf Whitefly)	0.28 to 1.1 lbs. LANCER 90 SP + 1 to 2 pts. LORSBAN® 4E	User should comply with all applicable directions, restrictions and precautions on the EPA registered label for LORSBAN® 4 E. Cutworms: Use 1 lb. per acre of LANCER 75 SP. By Ground: Ground application is recommended. Apply in 10 to 25 gals. of spray per acre. Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant. By Air: Aerial applications are less effective, but may be used. Apply in 3 to 10 gals. spray per acre (minimum 5 gals spray per acre in CA). Apply when insects first appear or when damage is first noted and repeat application as necessary to maintain control.	21

INSECTS	AMOUNT LANCER + DANITOL® 2.4 EC PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Sweetpotato Whitefly (Silverleaf Whitefly)	0.5 to 1.0 lb. LANCER 90 SP + 10 3/4 to 16 fl. oz. DANITOL® 2.4 EC	Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA) or 10 to 25 gals. spray per acre by ground. User should comply with all applicable directions, restrictions and precautions on the EPA registered label for DANITOL® 2.4 EC Spray. Apply when insects first appear or when damage is first noted.	21

COTTON TANK MIXES WITH PYRETHROIDS

GENERAL USE PRECAUTIONS:

Do not feed gin trash or treated forage to livestock.
 Do not allow livestock to graze on treated areas.
 Do not apply more than 6½ lbs. of LANCER 90 SP per acre per crop cycle.

Synthetic Pyrethroids should be used within the guidelines of state and/or regional resistance management programs and recommendations.

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions which appear on all product labels. The most restrictive labeling applies when using a tank mix.

RESISTANCE-MANAGEMENT

Cotton pest control programs, especially those for control of Silverleaf Whitefly populations, should employ a properly designed resistance-management strategy. Such resistance-management strategies include mixture or rotation of alternative classes of chemistry including organo-phosphates, carbamates, pyrethroids or insect growth regulators. Consult your state or area agricultural extension service for local resistance management strategies and advice on alternative insecticides.

INSECTS	LANCER 90 SP AND TANK MIX PARTNER	AMOUNT OF LANCER 90 SP + TANK MIX PARTNER PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Aphids Bollworm Cabbage Looper Cotton Leaf Perforator Cutworms* Fall Armyworm Fleahoppers Pink Bollworm (AZ & CA) Plantbugs Stinkbugs Sweetpotato/ Silverleaf Whitefly Thrips Tobacco Budworm Whitefly	LANCER 90 SP + one of the following	0.5 to 1.0 lbs. per acre + one of the following	Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA) or 10 to 25 gals. spray per acre by ground. Begin applications when eggs or insects appear and apply as needed to maintain control. Stinkbugs: Use 0.8 lb. per acre of LANCER 90 SP. * Cutworms: Use 0.8 lb. per acre of LANCER 90 SP By Ground: Ground application is recommended. Control is most effective when ground applications is made in the evenings and sprays are directed toward the base and lower portion of plant. By Air: Aerial applications are less effective, but may be used. Apply when eggs or insects first appear or damage is first noted. Repeat application as necessary to maintain control.	21
	AMBUSH® 2 EC	Refer to the AMBUSH® 2 EC approved label for use instructions.		
	AMMO® 2.5 EC*	Refer to the AMMO® 2.5 EC approved label for use instructions.		
	ASANA® XL*	Refer to the ASANA® XL approved label for use instructions.		
	BAYTHROID® 2 EC	Refer to the BAYTHROID® 2 EC approved label for use instructions.		
	CAPTURE® 2 EC	Refer to the CAPTURE® 2 EC approved label for use instructions.		
	KARATE® 1 EC (Except CA)	Refer to the KARATE® 1 EC approved label for use instructions.		
	POUNCE® 3.2 EC*	Refer to the POUNCE® 3.2 EC approved label for use instructions.		
	SCOUT® X-TRA (Except CA)	Refer to the SCOUT® X-TRA approved label for use instructions.		

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE

- Keep pesticide in original container.
- Do not put concentrated or dilute into food or drink containers.
- Store in cool, dry place. Protect from excessive heat.
- Do not contaminate food or foodstuffs.
- Do not store or transport near feed or food.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

LIMITED WARRANTY:

1. United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. United Phosphorus does not make nor authorizes any agent or representative to make, any other warranty, guarantee or representations, express or implied, concerning this product. Specifically, **NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.**
2. Critical and unforeseeable factors beyond United Phosphorus's control prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are limited to, damage to plants and crops to which product is applied, lack of complete control over the handling and application of this product, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit under average conditions for the use stated on the labeling and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by United Phosphorus under 1. above) resulting from handling, storage and use of the product.
3. Precautions stated on the labeling should be followed to avoid hazardous exposure to the product.

- AMBUSH® -- Reg. TM of Zeneca Inc. Ag Products for permethrin insecticide.
- AMMO® -- Reg. TM of FMC Agri Chemical Group for cypermethrin insecticide.
- ASANA® -- Reg. TM of E.I. DuPont de Nemours & Co., Inc for esfenvalerate insecticide.
- BAYTHROID® -- Reg. TM of Miles, Inc. for cyfluthrin synthetic pyrethroid.
- CAPTURE® -- Reg. TM of FMC Agri Chemical Group for bifenthrin insecticide-miticide.
- DANITOL® -- Reg. TM of Sumotomo Chemical Company Ltd. for fenpropathrin Insecticide-miticide.
- KARATE® -- Reg. TM of Zeneca Ag Products U.K. for lambda-cyhalothrin insecticide.
- LORSBAN® -- Reg. TM of Dow Chemical Company for chlorpyrifos insecticide.
- POUNCE® -- Reg. TM of FMC Agri Chemical Group for permethrin Insecticide.
- SCOUT® -- Reg. TM of Hoechst-Roussel Agri-Vet Co. for tralomethrin pyrethroid insecticide.