

70506-2

02/22/2005

1/2

Please read instructions on reverse before completing form

Form Approved, OMB No. 2070-0060, Approval expires 5-31-98

	United States	<input type="checkbox"/> Registration	OPP Identifier Number <b>274841</b>
	Environmental Protection Agency	<input type="checkbox"/> Amendment	
	Washington, DC 20460	<input checked="" type="checkbox"/> Other	

Application for Pesticide - Section I		
1. Company/Product Number 70506-2	2. EPA Product Manager M. Mautz	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) United Phosphorus, Inc/Acephate 90SP	PM # 4	
5. Name and Address of Applicant (Include ZIP Code) United Phosphorus, Inc. 423 Riverview Plaza Trenton, NJ 08611	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg No. _____ Product Name _____	
<input type="checkbox"/> Check if this is a new address		

Section - II	
<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Other - Explain below
<b>NOTIFICATION FEB 22 2005</b>	
Explanation: Use additional page(s) if necessary. (For Section I and Section II.) Notification of additional packaging size: 8 pound bags.	
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.	

Section III			
Material This Product Will be Packaged in:			
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____
*Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		If "Yes" Package wgt.	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of label directions <input type="checkbox"/> On Label <input type="checkbox"/> On Label accompanying product
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled	<input type="checkbox"/> Other _____

Section IV			
1. Contact Person (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)			
Name Rebecca A. Clemmer	Title Regulatory Affairs Manager	Telephone No. (Include Area Code) 609-392-8200	
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law			6. Date Application Received  (Stamped)
2. Signature 		3. Title Regulatory Affairs Manager	
4. Typed Name Rebecca A. Clemmer		5. Date February 11, 2005	

a/p

# ACEPHATE 90 SP

Soluble Powder  
(INSECTICIDE)

NOTIFICATION  
FEB 22 2005

ACTIVE INGREDIENT		By Wt.
Acephate (O, S-Dimethyl acetylphosphoramidothioate)	.....	90%
OTHER INGREDIENTS	-----	10%
	TOTAL	100%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

FIRST AID	
Contains an organophosphate that inhibits cholinesterase.	
If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information.	
NOTE TO PHYSICIAN Acephate is a cholinesterase inhibitor. If signs of cholinesterase inhibition appear, atropine is antidotal. 2-PAM may also be used in conjunction with atropine but should not be used alone.	
For Chemical Emergency Assistance (Spill, Leak, Fire or Accident), Call CHEMTREC 1-800-424-9300	

EPA Reg. No. 70506-2

EPA Est. No. 33560-TN-1

Manufactured for:  
United Phosphorus, Inc.  
423 Riverview Plaza  
Trenton, NJ 08611  
1-800-247-1557 • www.upi-usa.com

Net Contents: 5, 8, 10, 20 LBS *new size*

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed. Causes eye irritation. Avoid contact with eyes, skin and clothing. Avoid breathing dust or spray mist. Wash hands thoroughly after handling.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE):

**Applicators and other handlers must wear:** long-sleeved shirt and long pants, shoes plus socks and chemical-resistant headgear for overhead exposure.

**Mixers and Loaders must wear:** long-sleeved shirt and long pants, waterproof gloves, shoes plus socks and chemical-resistant headgear.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### USER SAFETY RECOMMENDATIONS

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Cover or soil-incorporate spills.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

4/12

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers-of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas until restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, waterproof gloves, shoes plus socks and chemical resistant headgear for overhead exposure

---

**DIRECTIONS**

---

**TANK MIXES**

**NOTICE:** Tank mixing of this product with any other product which is not specifically and expressly authorized by the label shall be exclusive risk of the use, applicator, and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

TABLE OF EQUIVALENTS

AMOUNT OF ACEPHATE 90 SP PER ACRE	ACRES TREATED BY A 5 LB. PACKAGE	ACRES TREATED BY A 10 LB. PACKAGE	ACRES TREATED BY A 20 LB. PACKAGE
2.5 oz.	32	64	128
3.2 oz.	25	50	100
3.25 oz.	24.6	49.2	98.4
0.25 lb.	20	40	80
0.28 lb.	17.8	35.7	71.4
0.5 lb.	10	20	40
0.56 lb.	8.9	17.8	35.6
0.8 lb.	6.25	12.5	25
1.0 lb.	5	10	20
1.1 lb.	4.5	9.1	18.2

---

**GENERAL INFORMATION**


---

Do not use this product through any type of irrigation system

ACEPHATE 90 SP is an insecticide which controls a wide range of important pests in cotton. The active ingredient is acephate, a water-soluble insecticide readily absorbed by plant roots and foliage to give systemic control of feeding insects. Insect pests are generally controlled more effectively by ACEPHATE 90 SP through ingestion than by contact. ACEPHATE 90 SP can be applied as hopper-box, in-furrow and foliar treatments in cotton. Application of ACEPHATE 90 SP to maintain control should be repeated only as directed.

FAILURE TO FOLLOW THE DIRECTION FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, AND/OR ILLEGAL RESIDUES.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

NOTE: This product is sold by weight and package is full when packed, but due to the product's nature, settling is likely to occur.

---

**PREPARATION OF SPRAY SOLUTION**


---

To prepare spray solution, empty the entire contents of package into a spray tank containing at least one-half of the total quantity of water required. Add remaining water with the agitator running until the spray volume is reached. Do not add any liquid fertilizers, micronutrients or adjuvants to the spray solution until the ACEPHATE 90 SP has completely dissolved. ACEPHATE 90 SP should completely dissolve in approximately five minutes. Dissolution rate may be slowed by cold water, lack of agitation, or water containing high concentrations of boron or sulfur.

4/12

**COTTON**

**GENERAL USE PRECAUTIONS:**

- Do not use treated seed for food or feed purposes or process for oil.
- Do not feed gin trash to livestock.
- Do not allow animals to graze on treated areas.
- Do not apply more than 6 2/3 lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.

**COTTON SEED HOPPER BOX TREATMENT  
(EXCEPT AZ, CA AND THE BLACK LANDS OF TEXAS)**

INSECTS	AMOUNT ACEPHATE 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS
Black (Greasy) Cutworm Cotton Aphids Thrips	2.5 to 3.25 oz.	<p>Add dry ACEPHATE 90 SP to seed in the planter box to achieve an application rate of 2.5 to 3.25 oz. per acre.</p> <p>Even coverage of seed is required for acceptable insect control. Plants grown from improperly treated seed may not exhibit insect control. To obtain even coverage, fill planter box 1/2 full of seed, add 1/2 of required ACEPHATE 90 SP and mix thoroughly. Add remainder of seed and ACEPHATE 90 SP to the hopper box and thoroughly mix.</p> <p>Improved distribution of ACEPHATE 90 SP on seed can be achieved by premixing both in a separate container and then transferring treated seed to hopper box.</p> <p>Use of ACEPHATE 90 SP with an air planter is not recommended since loss of product will occur.</p>
<p><b>USE PRECAUTIONS:</b></p> <ul style="list-style-type: none"> <li>• Do not use treated seed for food or feed purposes or process for oil.</li> <li>• ACEPHATE 90 SP cotton seed hopper box treatment is not permitted in Arizona, California or the Black Lands of Texas.</li> </ul>		

**COTTON IN-FURROW APPLICATION**

INSECTS	AMOUNT ACEPHATE 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS
<p>Aphids* Thrips Black (Greasy) Cutworm (Except CA)</p> <p>* Excluding Cotton Aphids in AZ &amp; CA</p>	<p>0.55 to 1.1 lbs.</p> <p>NOTE: for the Blacklands of Texas use 1.1 lbs. per acre ACEPHATE 90 SP</p>	<p>Do not apply more than 6 2/3 lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.</p> <p>Apply ACEPHATE 90 SP in 3 to 5 gals. of water per acre as an in-furrow spray at planting. ACEPHATE 90 SP can be mixed with fungicides that are sprayed in-furrow for disease control.</p> <p>Flat-fan nozzles used for in-furrow application should be set so that the fanned spray pattern is aligned with the row to insure good spray deposition in the seed furrow. Cone type nozzles may not provide a spray pattern that insures maximum spray deposition in the seed furrow, and should be avoided. Spray systems that employ metal or plastic tubing for delivery of spray solution into the seed furrow should have tubing securely fastened to the furrow opener and should be checked frequently to insure that the tubing is properly positioned to deposit spray solution into the seed furrow.</p>

7/12

**COTTON (Continued)**

**FOLIAR APPLICATION**

INSECTS	AMOUNT ACEPHATE 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Thrips	3.2 oz.	Do not apply more than 6 <sup>2</sup> / <sub>3</sub> lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.	21
Plant Bugs (Lygus)	0.25 to 1.0 lbs.  0.5 to 1.0 lbs. (CA and AZ)	Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA and AZ) or 10 to 25 gals. spray per acre by ground.	
Fleahopper	0.25 lb.	Apply when eggs or insects first appear. Repeat as necessary to maintain control.	
Cotton Aphid (excluding AZ and CA)	0.5 to 1.0 lbs.	Lygus -- Use higher rate for lygus adults that have migrated into cotton.	
Whitefly (excluding Sweetpotato /Silverleaf Whitefly)	0.5 to 1.0 lbs.	Cotton Aphid -- This insect may develop resistance to various classes of insecticides. Consult your local Agricultural Extension Service for current control recommendations.	
Armyworms (excluding Beet Armyworm) Cabbage Looper	1.0 lbs.	Armyworms & Cabbage Looper -- Apply when eggs appear and repeat at 3 to 7 day spray intervals.	
Bollworm Tobacco Budworm Adults Larvae Eggs (DEADHATCH®)	0.5 to 1.0 lbs. (East of Rockies) 1.0 lbs. (CA & AZ) Emerging Larvae 0.25 to 0.5 lbs.	Bollworm and Tobacco Budworm -- Early season light infestation use 0.5 lb. per acre. Mid and late season moderate to severe infestations use 0.8 to 1.0 lbs. per acre. Apply when eggs appear and repeat at 3 to 7 day spray interval.	
Stinkbugs	0.8 lb.	Adult/Larvae: Moths are controlled when they come in direct contact with spray particles during application. Moth kill is most likely to occur when application is made late in the evenings during periods of peak activity.	
Pink Bollworm (AZ & CA)	1.0 lbs.	Emerging Larvae: 0.25 to 0.5 pounds per acre when larvae first appear  DEADHATCH®: Control of emerging larvae by consumption of treated egg casings.  Pink Bollworm -- Apply when insects appear and repeat at a 5 to 7 day spray interval.	

---

**COTTON (Continued)**


---

**FOLIAR APPLICATION (Continued)**

INSECTS	AMOUNT ACEPHATE 90 SP PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Cutworms	0.8 lb.	<p>Cutworms -- Ground application is recommended.</p> <p>Aerial applications are less effective, but may be used. Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA and AZ) or 10 to 25 gals. spray per acre by ground.</p> <p>Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant.</p> <p>Apply when insects first appear or damage is first noted and repeat application as necessary to maintain control. Do not apply more than 6 <sup>2</sup>/<sub>3</sub> lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.</p>	21

---

**COTTON TANK MIXES**


---

**GENERAL USE PRECAUTIONS:**

Do not feed gin trash or treated forage to livestock.

Do not allow livestock to graze on treated areas.

Do not apply more than 6 <sup>2</sup>/<sub>3</sub> lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions that appear on all product labels. The most restrictive labeling applies when using a tank mix.

**RESISTANCE-MANAGEMENT**

Cotton pest control programs, especially those for control of Sweetpotato/Silverleaf Whitefly populations, should employ a properly designed resistance-management strategy. Such resistance-management strategies include mixture or rotation of alternative classes of chemistry including organo-phosphates, carbamates, pyrethroids or insect growth regulators. Consult your state or area agricultural extension service for local resistance management strategies and advice on alternative insecticides.



9/12

**COTTON TANK MIXES (Continued)**

INSECTS	AMOUNT ACEPHATE 90 SP + LORSBAN® 4E PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Armyworms (excluding Beet Armyworm) Bollworms Cabbage Looper Cotton Aphid Cutworms Fleahopper Grasshoppers Pink Bollworm Plant Bugs (Lygus, Mirids) Salt Marsh Caterpillar Thrips Tobacco Budworm Whitefly (excluding Sweetpotato/ Silverleaf Whitefly)	0.28 to 1.1 lbs. ACEPHATE 90 SP + 1 to 2 pts. LORSBAN® 4E	<p>User should comply with all applicable directions, restrictions and precautions on the EPA registered label for LORSBAN® 4 E.</p> <p>Cutworms: Use 0.8 lb. per acre of ACEPHATE 90 SP.</p> <p>By Ground: Ground application is recommended. Apply in 10 to 25 gals. of spray per acre.</p> <p>Control is most effective when ground application is made in the evenings and sprays are directed toward the base and lower portion of plant.</p> <p>By Air: Aerial applications are less effective, but may be used. Apply in 3 to 10 gals. spray per acre (minimum 5 gals spray per acre in CA).</p> <p>Apply when insects first appear or when damage is first noted and repeat application as necessary to maintain control.</p> <p>Do not apply more than 6 <sup>2</sup>/<sub>3</sub> lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.</p>	21

INSECTS	AMOUNT ACEPHATE + DANITOL® 2.4 EC PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Sweetpotato/ Silverleaf Whitefly	0.5 to 1.0 lb. ACEPHATE 90 SP + 10 <sup>2</sup> / <sub>3</sub> to 16 fl. oz. DANITOL® 2.4 EC	<p>Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA) or 10 to 25 gals. spray per acre by ground.</p> <p>User should comply with all applicable directions, restrictions and precautions on the EPA registered label for DANITOL® 2.4 EC Spray.</p> <p>Apply when insects first appear or when damage is first noted.</p> <p>Do not apply more than 6 <sup>2</sup>/<sub>3</sub> lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.</p>	21

19/12

**COTTON TANK MIXES WITH PYRETHROIDS**

**GENERAL USE PRECAUTIONS:**

Do not feed gin trash or treated forage to livestock.

Do not allow livestock to graze on treated areas.

Do not apply more than 6 2/3 lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar.

Synthetic Pyrethroids should be used within the guidelines of state and/or regional resistance management programs and recommendations.

Always read and follow all label directions when using any pesticide alone or in tank mix combinations. Observe all restrictions and precautions that appear on all product labels. The most restrictive labeling applies when using a tank mix.

**RESISTANCE-MANAGEMENT**

Cotton pest control programs, especially those for control of Sweetpotato/Silverleaf Whitefly populations, should employ a properly designed resistance-management strategy. Such resistance-management strategies include mixture or rotation of alternative classes of chemistry including organo-phosphates, carbamates, pyrethroids or insect growth regulators. Consult your state or area agricultural extension service for local resistance management strategies and advice on alternative insecticides.

INSECTS	ACEPHATE 90 SP AND TANK MIX PARTNER	AMOUNT OF ACEPHATE 90 SP + TANK MIX PARTNER PER ACRE	ADDITIONAL USE INSTRUCTIONS	DAYS TO HARVEST
Aphids Bollworm Cabbage Looper Cotton Leaf Perforator Cutworms* Fall Armyworm Fleahoppers Pink Bollworm (AZ & CA) Plantbugs Stinkbugs Sweetpotato/Silverleaf Whitefly Thrips Tobacco Budworm Whitefly	ACEPHATE 90 SP + one of the following	0.5 to 1.0 lbs. per acre + one of the following	Do not apply more than 6 2/3 lbs. of ACEPHATE 90 SP per acre per crop cycle. This limitation includes all methods of application; i.e. hopper-box, in-furrow, and foliar. Apply in water at 3 to 10 gals. spray per acre by air (minimum 5 gals per acre in CA) or 10 to 25 gals. spray per acre by ground. Begin applications when eggs or insects appear and apply as needed to maintain control. Stinkbugs: Use 0.8 lb. per acre of ACEPHATE 90 SP. * Cutworms: Use 0.8 lb. per acre of ACEPHATE 90 SP By Ground: Ground application is recommended. Control is most effective when ground applications are made in the evenings and sprays are directed toward the base and lower portion of plant. By Air: Aerial applications are less effective, but may be used. Apply when eggs or insects first appear or damage is first noted. Repeat application as necessary to maintain control.	21
	AMMO® 2.5 EC*	Refer to the AMMO® 2.5 EC approved label for use instructions.		
	ASANA® XL*	Refer to the ASANA® XL approved label for use instructions.		
	BAYTHROID® 2 EC	Refer to the BAYTHROID® 2 EC approved label for use instructions.		
	CAPTURE® 2 EC	Refer to the CAPTURE® 2 EC approved label for use instructions.		
	KARATE® 1 EC (Except CA)	Refer to the KARATE® 1 EC approved label for use instructions.		
	SCOUT® X-TRA (Except CA)	Refer to the SCOUT® X-TRA approved label for use instructions.		

11/12

### STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

**PESTICIDE STORAGE:** Keep pesticide in original container. Do not put concentrated or dilute product into food or drink containers. Store in a cool, dry place. Protect from excessive heat. Do not contaminate food or feedstuffs. Do not store or transport near feed or food.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Do not reuse container. Dispose of empty bag in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT INFORMATION READ BEFORE USING PRODUCT

**LIMITED WARRANTY:**

1. United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for use under average conditions when used strictly in accordance with the directions on the labeling. United Phosphorus does not make nor authorizes any agent or representative to make any other warranty, guarantee or representations, express or implied, concerning this product. Specifically, NO IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR ANY PARTICULAR PURPOSE IS MADE.
2. Critical and unforeseeable factors beyond United Phosphorus' control prevent it from eliminating all risks in connection with the use of this product. Such risks include, but are not limited to, damage to plants and crops to which product is applied, lack of complete control over the handling and application of this product, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit under average conditions for the use state on the labeling and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by United Phosphorus under 1. above) resulting from handling, storage, and use of the product.
3. Precautions stated on the labeling should be followed to avoid hazardous exposure to the product.

AMMO® -- Reg. TM of FMC Corporation for cypermethrin insecticide.  
 ASANA® -- Reg. TM of E.I. duPont de Nemours & Co., Inc. for esfenvalerate insecticide.  
 BAYTHROID® -- Reg. TM of Bayer Crop Protection for cyfluthrin synthetic pyrethroid.  
 CAPTURE® -- Reg. TM of FMC Corporation for bifenthrin insecticide-miticide.  
 DANITOL® -- Reg. TM of Sumitomo Chemical Company Ltd. for fenpropathrin insecticide-miticide.  
 KARATE® -- Reg. TM of Syngenta Crop Protection, Inc. for lambdacyhalothrin insecticide.  
 LORSBAN® -- Reg. TM of Dow Agro Sciences for chlorpyrifos insecticide.  
 SCOUT® X-TRA -- Reg. TM of Hoechst-Roussel Agri for tralomethrin pyrethroid insecticide.

---

Approved by EPA 5/11/00  
 REVISED: 3-25-2002 for First Aid, Bee Statement;  
 Revised 8-22-02 for Precautionary Statements, Bee Statement.  
 EPA APPROVED 10-30-02  
 Revised 7/21/04 to update product name, other details

12/12



# United Phosphorus, Inc.

423 Riverview Plaza  
Trenton, NJ 08611  
(609) 392-8200 x 18 (phone)  
(609) 392-0808 (fax)

Rebecca A. Clemmer  
Regulatory Affairs Manager

February 11, 2005

Marilyn Mautz (PM 4)  
Document Processing Desk (NOTIF)  
Office of Pesticide Programs (7504C)  
U.S. Environmental Protections Agency  
1801 Bell Street  
Arlington, VA 22202

Re: Notification of Additional Packaging Size  
Acephate 90SP (EPA Reg. No. 70506-2)

Dear Ms. Mautz:

United Phosphorus, Inc. wishes to notify the Agency of an alternate packaging size for the subject product: 8 lb bags. Enclosed in support of this action, please find:

- EPA Form 8570-1
- One copy of the label, marked to show changes.

Please contact me if you have any questions.

Very truly yours,

A handwritten signature in cursive script that reads "Rebecca A. Clemmer".

Rebecca A. Clemmer  
[rclemmer@upi-usa.com](mailto:rclemmer@upi-usa.com)