



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 6, 2019

Crystal Layton
United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

Subject: Label Amendment – Adding squeeze and pour type container directions
Product Name: Tengard HG Termiticide/Insecticide
EPA Registration Number: 70506-7
Application Date: September 6, 2018
Decision Number: 544423

Dear Ms. Layton:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Autumn Metzger at 703-919-7453, or Metzger.autumn@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Elizabeth Fertich". The signature is fluid and cursive, with the first name and last name clearly distinguishable.

Elizabeth Fertich
Product Manager 04
Invertebrate & Vertebrate Branch 1
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

TENGARD HG TERMITICIDE/INSECTICIDE

Intended for outdoor homeowner use only

Active Ingredient


Permethrin*36.8%

Other Ingredients**63.2%

TOTAL 100.0%

*cis/trans ratio: Max 42% (±) cis and min. 58% (±) trans

** contains petroleum distillates



KEEP OUT OF THE REACH OF CHILDREN CAUTION

See [booklet on] [back/side/other panel(s)] for First Aid and additional Precautionary Statements [and Directions for Use]

FIRST AID	
IF SWALLOWED:	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
IF ON SKIN:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF IN EYES:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>NOTE TO PHYSICIAN: Contains petroleum distillates – vomiting may cause aspiration pneumonia.</p>	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency medical assistance, contact the Rocky Mountain Poison and Drug Center at 1-866-767-5089.</p>	

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300.

UPL NA Inc. 630 Freedom Business Center, Suite 402 King of Prussia PA 19406 1-800-438-6071	EPA Reg. No. 70506-7 EPA Est. No. _____
---	--

Net Contents: _____ **Fluid Ounces**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION

Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems. Use with care when applying to areas adjacent to any body of water. Do not apply directly to water. Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish ponds. Do not contaminate water when disposing of equipment wash waters.


This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds when bees are foraging the treatment area.

PHYSICAL OR CHEMICAL HAZARDS: Combustible. Do not use or store near heat or open flame.

DIRECTIONS FOR USE

[Reviewer notes: Container use directions when product is packaged in either the Tip and Pour or Squeeze and Pour container.]

[Optional text CONTAINER USE DIRECTIONS:



1. Remove the measuring chamber cap and induction seal. Replace cap and securely tighten. Tip container until liquid fills measuring chamber.
2. Return container to level position. No adjustment is needed.
3. Remove measuring chamber cap and dispense into proper application equipment.]

[OPTIONAL CONTAINER USE DIRECTIONS as listed below



1. Remove the measuring chamber cap and the seal. Squeeze container until liquid fills measuring chamber to the desired amount.
2. Dispense the product into the proper application equipment. Securely place the caps back on the bottle.]

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Failure to follow the directions for use and precautions on this label may result in poor pest control, plant injury or illegal residues.

Product Information

This product is non-staining to most home siding depending on age and cleanliness. However, before using in areas where the spray may contact home siding (vinyl siding, in particular), test in an inconspicuous area and recheck in a few hours. Do not use if any staining is observed

This product may be used to control insect pests outdoors, on ornamentals, lawns, trees, shrubs, and vines in landscape areas, as a perimeter treatment around residential buildings. Tengard Insecticide may also be used to control home-invading insects as a residual spray on outside surfaces, such as, garbage areas, window frames, and doorways. Not for use on plants being grown for commercial sale, or plants grown for seed production or sod farms and golf course turf, or in greenhouses or nurseries.

Application Instructions: Tengard Insecticide mixes readily with water and is specifically packaged for residential use as a diluted spray. Use a clean sprayer, such as, a tank type or hose-end sprayer to mix and apply the proper amount of concentrate with the desired volume of water. Ensure all storage and disposal instructions are followed. For perimeter and lawn applications measure the area to be treated to determine the application volume. For trees, shrubs and flowers apply as a thorough cover spray until foliage is wet. Refer to **Specific Application Instructions** in the rate Tables for additional instructions.

Product Use Restrictions

For use as a surface spray:

Do not allow adults, children, or pets to enter until sprays have dried.

All outdoor applications must be limited to spot or crack-and-crevice treatments only, except for the following permitted uses:

1. Treatment to soil or vegetation around structures;
2. Applications to lawns, turf, and other vegetation;
3. Applications to building foundations, up to a maximum height of 3 feet.

Other than applications to building foundations, all outdoor applications to impervious surfaces such as sidewalks, driveways, patios, porches, and structural surfaces (such as windows, doors, and eaves) are limited to spot and crack-and-crevice applications only.

For outdoor residential use:

Do not water the treated area to the point of runoff.

Do not make applications during rain.

Not for use in outdoor residential misting systems.

DO NOT use on vegetables or food crops.

DO NOT treat pets with this product.

DO NOT spray inside dwellings.

DO NOT treat soil that is water saturated or frozen or in any conditions where run-off or movement from the treatment area (site) is likely to occur.

DO NOT apply when wind velocity favors drift.

Do not apply directly to water.

Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish ponds.

Do not contaminate water when disposing of equipment wash waters.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

HOME-INVADING PESTS (OUTDOOR USE ONLY)

MIXING INSTRUCTIONS

Mix 1.6 fluid oz. (3¼ Tablespoons) of Tengard Insecticide concentrate in 1 gallon of water to make a 0.5% spray emulsion [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. [Optional text: Use provided [Squeeze/Tip] and Pour measuring chamber to measure the required amount of product.] Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Spot Treatment

- Apply a 0.5 % emulsion as a pin stream low-pressure spray or as a paint-on application with a paintbrush.
- Treat where pests are found or entry points of the structure such as window frames, doorways, porches, patios, and foundation.
- Spot-treatment may also be made to garages, garbage cans, and garbage areas.
- DO NOT apply indoors (including crawl spaces).

Perimeter Application

- Spray a 6 to 8 foot wide band of soil around the perimeter of the house. Treat the foundation wall to a height of 2 to 3 feet.
- Use a spray volume of 2 to 10 gallons of 0.5% emulsion per 1000 square feet. If mulch or debris is present, a higher volume application rate may be needed to ensure adequate coverage.
- Note: residual control will vary depending on exposure of the treated area to weathering conditions (temperature, sunlight, moisture). Retreat on a monthly schedule while insects are likely to be active.

Wood Destroying Pests

- Localized Areas Outdoors: for the control of insects such as termites, bees, carpenter ants and wood infesting beetles in localized above-ground areas of infested wood around structures, apply a 0.5% emulsion to voids and galleries in damaged wood and in spaces between wooden members of a structure and between wood and foundations where wood is vulnerable. Apply either as a spray or paint-on application. For termites the purpose of application is to destroy workers or winged reproductive forms, which may be present at the time of treatment. This product is not designed for preventative use and should only be applied when pests appear. This application is not a substitute for soil or foundation treatment. For active infestations inside a structure, consult a licensed Pest Control Service.
- Use a hose-end sprayer or sprinkling can to deliver a coarse drenching spray to treat stored lumber or woodpiles. Stored wood can be used for lumber one month after treatment. Do not apply inside structures.
- Repeat treatment as necessary but not more frequently than once a month. For active termite infestation consult a professional Pest Control Applicator.

HOME-INVADING PESTS (OUTDOOR USE ONLY)	
PEST	SPECIFIC APPLICATION INSTRUCTIONS
Ants (Incl. mounds) Armyworm Bark Beetles Bees Beetles Borers Boxelder Bugs Carpenter Ants Carpenter Bees Centipedes Chinchbugs Cockroaches Cockroaches (Asian) Crickets Earwigs Elm Leaf Fire Ants (Incl. mounds) Fleas Flies Firebrats Ground Beetles Gypsy Moths (adults & caterpillars) Millipedes Mole Crickets Pill Bugs Scorpions Silverfish Sod Webworm Sowbugs Spiders Ticks (including Deer Tick, Western Black-legged Tick known to transport Lyme disease and Rocky Mountain Spotted Fever) Wasps	<p>Boxelder Bugs, Elm Leaf Beetles and Gypsy Moth Caterpillars: Spray tree trunks, building siding or wherever pests congregate. Avoid excess application. Dampen surface thoroughly, but do not let runoff occur.</p> <p>Borers and Bark Beetles: To prevent infestation of ornamental trees and woody shrubs, spray the bark to the point of saturation, but do not let runoff occur. Repeat treatment every 21 days during early spring until mid-summer.</p> <p>Carpenter Bees: Apply as a pin stream, as a fine/course low pressure spray (20 psi or less) as a spot treatment.</p> <p>Fire Ants (mound drench): Apply 1 to 2 gallons of emulsion to each mound area by sprinkling the mound until it is wet and treat a 4-foot diameter circle around the mound. Use higher volume for mounds larger than 12 inches. For best results, apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day.</p> <p>Scorpions: remove accumulations of lumber, firewood, and other materials which serve as harborage sites and treat beneath and adjacent to infested sites.</p>

LAWN PESTS

Tengard HG Insecticide can be applied to lawns adjacent to or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, fence lines, storage sheds, barns, and other residential areas where pests congregate.

MIXING INSTRUCTIONS

Prepare an emulsion by mixing 0.4 to 0.8 fluid oz. (2½ to 5 teaspoons) of Tengard HG Insecticide Concentrate in sufficient volume of water to cover 1000 sq ft (approximately 4 to 25 gallons of water). [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Mound Drench Treatment (Ant and Fire Ant Mounds in Lawns)

Use a sprinkling can. Mix 1.6 fluid oz. (3¼ Tablespoons) of Tengard Insecticide Concentrate per 1 gallon of water to make a 0.5% spray emulsion [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. For large ant mounds, such as Fire Ant mounds, apply 1-2 gallons of a 0.5% emulsion to each mound area including a 4 foot diameter circle around the mound. Apply to smaller mounds until the mound is completely saturated. Use higher water volume for mounds larger than 12".

Lawn Broadcast Application

- Apply using a hose-end or pressure type sprayer to deliver a coarse spray of between 4 to 25 gallons of emulsion per 1,000 sq. ft. of lawn.
 - Use the higher label rate for heavy pest infestations.
 - Use sufficient volume to uniformly cover plant surface.
- Note: Use the higher volume of water when situations of thick thatch or dense sod exist, or in arid or drought conditions.

WHEN TO APPLY

- For best results treat in early morning or late evening hours, but not in the heat of the day.
- Spray evenly over the lawn when insects first appear.
- Repeat applications as necessary but not more frequently than every 10-14 days. Do not apply more than 1.6 fl oz/1000 sq ft per year

APPLICATION RATES FOR LAWN			
SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
Lawns , residential including but not limited to, St. Augustine, Bentgrass, Fescues, Buffalograss, Bahiagrass, Bluegrass, Perennial Ryegrass, and Centipede grass	Ants Armyworms Bilbugs Chiggers Chinchbugs Cricketts Cutworms Earwigs Fruit Flies Fleas Leafhoppers Millipedes Mole Crickets Sod Webworm Sowbugs Ticks (including Brown Dog Tick, and, Deer Tick, Western Black-legged Ticks known to transport Lyme disease and Rocky Mountain Spotted Fever). <i>For additional pests consult the list of pests under Outdoor Pests Controlled by Tengard HG Insecticide</i>	0.4 to 0.8 Fluid Ounces (2½ to 5 tsp) per 1000 sq. ft.	Apply using 4 – 25 gallons of spray volume/1000 sq. ft. of lawn. Use higher volumes on St. Augustine, Buffalograss, Bermuda and Bahiagrass lawns. Use the higher listed rate for fast knockdown and increased residual control. Dense and excessive (greater than 3 inches) lawn height and arid conditions may require higher volume application rates. Repeat applications may be necessary (10 to 14 day interval) to control heavy infestations during peak periods of insect feeding. Do not exceed a total of 1.6 fl oz/1000 sq ft per year (i.e. four (4) applications at the lower rate or two (2) applications at the higher rate).

ORNAMENTAL PESTS

MIXING INSTRUCTIONS

Mix 0.04 to 0.08 fluid oz. (= 1/8 to 1/4 teaspoons = 1-2 ml) of Tengard Insecticide Concentrate in 1 gallon of water for general pest control. Refer to the Ornamental Application Rate Table for additional mixing instructions and for specific directions for stem-infesting beetles. [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp) = 29.6 ml]. Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Ornamentals (Foliage and Flowering Plants)

- Apply using a hose-end or pressure type sprayer to deliver a coarse spray. Apply thoroughly to stem and foliage.
- Use the higher label rate (1/4 tsp. per gal of water) for heavy pest infestations. Note: Use the higher volumes of water when situations of dense foliage exists, or in arid or drought conditions.

WHEN TO APPLY

- For best results treat in early morning or late evening hours, but not in the heat of the day.
- Spray evenly over the lawn when insects first appear.
- To prevent insect reinfestation repeat treatment as necessary but not more frequently than once every 14 days.

APPLICATION RATES FOR USE ON ORNAMENTALS			
SITE	PEST	RATE	SPECIFIC APPLICATION INSTRUCTIONS
Ornamental foliage and flowering plants (Inc. Conifers, woody and herbaceous non-edible ornamentals.	Ants	0.04 to 0.08	Note: Direct application to blooms may cause browning of petals. Marginal leaf burn may occur on Salvia, Dieffenbachia and Pteris Fern.
	Aphids	Fluid	
	Bagworm	Ounces	
	Beet Armyworm	(1/8 to 1/4 tsp.)	Mosquitoes coming to rest on treated plants will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.
	Cabbage Looper		
	Cankerworms	(1 to 2 ml)	
	Citrus Thrips	Per gal.	¹ To control Webbing Coneworm make first application just prior to peak pollen flight. To control other Coneworms and Seed Bugs , make application 30 days prior to flower closure
	Coneworm (Incl. web) ¹		
	and Seed bugs		
	Fungus Gnat		
	Gypsy Moth		
	Caterpillars		
	Heliothis spp		
	Japanese Beetles		
	Lace Bug		
	Leaf Feeding		
	Caterpillars		
	Leaffooted Bugs		
	Leafminers		
	Leafhoppers		
	Leaf rollers		
	Lygus Bugs		
	Mealybugs		
	Mosquitoes		
	Nantucket PineTip		
	Moth		
	Oblique Leafroller		
	Pine Sawflies		
	Plant Bugs		
	Root Weevils (Adult)		
	Rosy Aphid		
	Rose Chafer		
	Spotted Leafminer		
Stink Bugs			
Tarnished Pant Bug			
Tent Caterpillars			
Webworms			
Whiteflies			
Zimmerman Pine Moth ¹			

	Bark Beetles and Boring insects, (Including but not limited to: Ash Borer, Bronze Birch Borer, Rhododendron Borer, Elm Bark Beetles and Turpentine Beetles)	0.3 to 0.6 fluid ounces (2 to 4 tsp) per gal	Stem-Infesting Beetles: Treat lower branches and trunk prior to adult emergence. Complete, uniform coverage is needed for best control. A repeat application may be necessary on a 3 to 4 week interval to prevent severe tree stem damage.
--	---	--	--

TERMITE CONTROL (SUBTERRANEAN)

Tengard HG Insecticide may be used to establish an exterior soil barrier against termites around structures. The chemical concentrate is mixed as an emulsion and adequately dispersed in the soil to provide a continuous exterior barrier between wood in the structure and the termite colonies in the soil.

Note: A complete termite barrier may require treatment with other EPA registered products to the inside perimeter of the foundation and to other interior critical areas. For these areas, and if termites are found inside of the home, consult a Licensed Professional Pest Control Service for treatment.

For use only around existing structures. Do not use inside dwelling. Not for Preconstruction use.

RESTRICTIONS:

When treating adjacent to an existing structure, the applicator must check the area to be treated, and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean up is completed.

MIXING INSTRUCTIONS

Using a sprinkler can add 3.2 fluid oz. (6.5 Tablespoons) to 2 gallons of water (to make a 0.5% emulsion) for each 5 linear feet of treatment to be provided [1 fluid oz. = 2 Tablespoons (T) = 6 teaspoons (tsp)]. Do not use utensils such as spoons and measuring cups for food purposes after use with pesticides.

HOW TO APPLY

Trench Treatment

- Remove all wood debris and wood forms. Around piers, chimney bases and along foundations of poured concrete, dig a trench 6 inches wide and 4 inches deep. If footing is more than 12 inches deep make holes with a steel rod or pipe one foot apart that extend from the bottom of the trench to the top of the footing. Make holes closer in hard packed clay soils. **Never dig below the top of the footing.**
- Apply a 0.5% emulsion at a rate of 4 gallons per 10 linear feet per foot of depth. Apply half of the emulsion when backfilling the trench. (One quart of Tengard Insecticide Concentrate will make 20 gallons of 0.5% emulsion and treat 50 linear feet.)

Restriction: DO NOT treat structures that contain cisterns or wells. Do not treat soil in the vicinity of wells, cisterns or ponds or structures with sump pumps or French drain. For those

situations, contact your County Cooperative Extension Agent or consult a commercial pest control service.

Fence Posts

- Pre-installation: Using a sprinkler can, pour ½ gallon of 0.5% spray emulsion into the fence post hole and add another ½ gallon in mixture with the soil when filling the hole so that all the soil is treated. Triple rinse sprinkler can after each use. Do not store any unused product in sprinkler can. Use and follow all directions for storage and disposal found on this label.
- Post-installation: installed poles and posts may be treated by first digging a trench around the post and applying 1 gallon of 0.5% emulsion per one foot depth of trench for posts less than 6 inches in diameter. For larger diameter posts increase the application rate to 4 gallon for each 10 linear feet of trench and per ft. of depth.

WHEN TO RE-APPLY

Retreatment for subterranean termites can only be performed if there is clear evidence of reinfestation or disruption of the barrier due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide barrier in the soil. These vulnerable or reinfestd areas may be retreated in accordance with application techniques described in this product's labeling. The timing and type of these retreatments will vary, depending on factors such as termite pressure, soil types, soil conditions and other factors, which may reduce the effectiveness of the barrier.

Annual retreatment of the structure is prohibited unless there is clear evidence that reinfestation or barrier disruption has occurred.

NOTICE: To the extent consistent with applicable law, buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

STORAGE AND DISPOSAL

Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal.

Pesticide Storage

Store at temperatures above 40°F (5°C).

Shake container well before using. If crystals form, warm to room temperature by placing container in a room at ambient temperature 70°F (21°C) until crystals dissolve.

Do not use or store near heat, open flame or hot surfaces.

Store in a dry place and avoid excess heat in storage. Store in original containers only.

Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. To confine spill, dike surrounding area or absorb with sand, cat litter, and commercial clay or gel absorbents. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

Pesticide Disposal

Pesticide wastes are toxic. Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council at www.acrecycle.org.

Triple rinse container promptly after emptying. If recycling is not available puncture and dispose of in a sanitary landfill or incineration or if allowed by state and local authorities by burning. If burned stay out of smoke.

For containers smaller than or equal to 5 gallons, triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

Rev. 02/26/2019, 09/24/2019, 10/29/2019 per EPA comments

[List of Optional Marketing Statements to be added to Final Print Label:

Brand Extensions

HG
H&G
Lawn
Outdoor

Use Statements

Insecticide
Insecticide/Termiticide
Termiticide
Lawn & Ornamental
Soil Insect & Termite Killer
Fire Ant Killer
Ant & Roach Killer
Outdoor Perimeter
For Outdoor Residential Use Only

Control Statements

Kills Ticks that may cause Lyme disease
Kills Listed Home Invading Pests (Outdoor)
Kills Named Insects
An easy to apply product to control Home Invading and Lawn Insects
An easy to apply product to control wood invading Insects
New Termiticide Formulation for Homeowner Use (Outdoor Use Only)
New Termite Insecticide (Outdoor Use Only)
New for Homeowner Use (Outdoor Use Only)
Controls Wood-Destroying Insects

Icons/Pictures

[Icon of People]
[Icon of Sprinkling Can]
[Icon of Hand Held Sprayer]
[Tree Icon]
[Plant Icon]
UPC symbology
Picture Fire Ant Mound
Picture of a residential home
Picture of a lawn
Pictures of one or more of the following insects:
termites, ants (Including Fire Ant mounds and Carpenter Ants),
cockroach, spider, scorpion, fleas, tick, mosquito

Other

Concentrate
Treats up to [# will depend on application] Square Feet
Made in the USA
Formulated and Packaged in the USA
How To Apply
When to Apply
Amount to Use]