


70506-20

2-11-2003

11/11

	U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460	EPA Reg. Number:  <b>70506-20</b>	Date of Issuance:  <b>FEB 11 2003</b>	
	<b>NOTICE OF PESTICIDE:</b> <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration	Term of Issuance:  <b>Conditional</b>		Name of Pesticide Product:  <b>Cypermethrin 2.5 EC AG Insecticide</b>
	(under FIFRA, as amended)	Name and Address of Registrant (include ZIP Code):  <b>United Phosphorus, Inc.          740 Springdale Drive, Suite 204          Exton, Pennsylvania 19341</b>		
<p><b>Note:</b> Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p><b>This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:</b></p> <ol style="list-style-type: none"> <li>Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.</li> <li>Make the labeling changes listed below before you release the product for shipment:           <ol style="list-style-type: none"> <li>Add the phase "EPA Registration No. 70506-20".</li> <li>Under Precautionary Statements change "Harmful if absorbed through skin, or swallowed." to "Harmful if absorbed through skin, inhaled, or swallowed. Causes eye irritation. Avoid breathing vapor or spray mist."</li> <li>Add the maximum usage instructions and table when using both zeta- cypermethrin and cypermethrin products. See EPA Registration Number 279-3027 for appropriate wording. Add the name of this product along with Ammo.</li> </ol> </li> </ol>				
Signature of Approving Official:		Date:		

page 2  
EPA Reg. No. 70506-20

2/11

3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

3/11

# CYPERMETHRIN 2.5 EC AG INSECTICIDE

## RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms.

For retail sale to, and use only by, Certified Applicators or persons under their direct supervision, and only for those users covered by the Certified Applicator's certification.

**Active Ingredient:**

*Cypermethrin**	.....30.6%
INERT INGREDIENTS***	.....69.4%
TOTAL	.....100%

\* (±)α-cyano-(3-phenoxyphenyl)methyl (±) *cis/trans*-3-(2,2-dichloroethenyl)-2,2-dimethylcyclopropane carboxylate

\*\*Cis/trans ratio: 47/53±10%

\*\*\*Contains xylene range aromatic solvents.

Contains 2.5 pounds cypermethrin per gallon

### KEEP OUT OF REACH OF CHILDREN

## CAUTION

### FIRST AID

**IF INHALED:**

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
- Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Do not give any liquid to the person.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**Note to Physician:** Contains petroleum distillates – vomiting may cause aspiration pneumonia.

### HOTLINE NUMBERS

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center at 1-800-858-7378 for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

United Phosphorous, Inc.  
423 Riverview Plaza, Trenton, NJ 08611

EPA Reg. No. 70506-

EPA Est. No.

Net Contents:

ACCEPTED  
with COMMENTS  
in EPA Letter Dated

FEB 11 2003

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

4/11  
✓

**CAUTION**

Harmful if absorbed through skin, <sup>inhalated</sup> or swallowed. Avoid contact with skin; may cause sensitization reaction in some individuals. Avoid contact with eyes or clothing.

Product should be opened and poured in a well ventilated area.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

**Handlers who may be exposed to the dilute through application or other tasks must wear:**

Coveralls over short-sleeved shirt and short pants, chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, shoes plus socks and protective eyewear.

**Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear:** Coveralls over short-sleeved shirt and short pants, chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton, shoes plus socks, protective eyewear, and chemical-resistant apron when mixing or loading.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over short-sleeved shirt and short pants.
- Chemical-resistant gloves, such as Barrier Laminate or Nitrile Rubber or Neoprene Rubber or Viton.
- Shoes plus socks.
- Protective eyewear.

5/11

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops if bees are visiting the treatment area.

**PHYSICAL/CHEMICAL HAZARDS**

Do not use or store near hear or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Resistance:** Some insects are known to develop resistance to products used repeatedly for control. Because the development of resistance cannot be predicted, the use of this product should conform to resistance management strategies established for the use area. Consult your local or state agricultural authorities for details.

If resistance to this product develops in your area, this product, or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of insect may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternative method of control for your area.

**CHEMIGATION USE DIRECTIONS**

Apply this product only through sprinkler, including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.

Crop injury, lack of effectiveness, or illegal residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump motor must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

6/11

CYPERMETHRIN 2.5 EC AG Insecticide should be applied continuously for the duration of the water application. CYPERMETHRIN 2.5 EC AG Insecticide should be diluted in sufficient volume to insure accurate application over the area to be treated. When using chemigation, a minimum of 0.5 inch per acre of irrigation water is recommended. Agitation is not required when a suitable diluent is used.

**GENERAL INSTRUCTIONS**

Use low rate under light to moderate infestation. Higher rates should be used under heavy insect pressure. The rate of application is variable according to insect pressure, timing of spray and field scouting.

**Rotational Crops:** With the exception of the crops listed below, rotational crops may not be planted within 30 days after the last application.

**Tank-Mixtures:** CYPERMETHRIN 2.5 EC AG Insecticide may be applied in tank mixtures with other products approved for use on cotton, brassicas, bulb vegetables, head lettuce and pecans. Observe all restrictions and precautions which appear on the labels of these products.

For improved control of aphids and mites on cotton, CYPERMETHRIN 2.5 EC AG Insecticide may be tank-mixed with Curacron® 8E. The tank-mixture is most effective when pest populations are first observed. Subsequent applications may be needed to maintain control.

**Spray Drift Precautions:** Do not make more than 10 synthetic pyrethroid applications (of one product or combination of products) to a cotton crop in one growing season.

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers.

**OBSERVE THE FOLLOWING PRECAUTIONS WHEN SPRAYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES AND COMMERCIAL FISH FARM PONDS.**

Do not apply by ground equipment within 25 feet, or by air within 150 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds. Increase the buffer zone to 450 feet when ultra low volume (ULV) application is made.

For aerial applications, the spray boom should be mounted on the aircraft so as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or rotor diameter.

Use the largest droplet size consistent with good pest control. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided.

Make aerial or ground applications when the wind velocity favors on-target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph.

Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Do not cultivate within 10 feet of the aquatic area so as to allow growth of a vegetative filter strip.

Low humidity and high temperatures increase the evaporation rate of spray droplets and therefore the likelihood of increased spray drift to aquatic areas. Avoid spraying during conditions of low humidity and/or high temperature.

Do not make aerial or ground applications during temperature inversions. Inversions are characterized by stable air and increasing temperatures with height above the ground. Mist or fog may indicate the presence of an inversion in humid areas. The applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

SC 1000 0000

7/11

**COTTON (14 DAY PHI)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL. OZ./A	
Pre-emergent Use: Cutworms	0.025 to 0.1	1.3 to 5	Use CYPERMETHRIN 2.5 EC AG Insecticide in the time period from 14 days prior to planting up to emergence of the crop. Apply as a broadcast spray by ground or air, banded (including T-band) or in-furrow spray using sufficient spray volume to achieve adequate coverage. Reduced volumes of water may be used with specialized equipment. Use a minimum of 1 gallon of water per acre by air. Use the higher rates of CYPERMETHRIN 2.5 EC when incorporating into the soil.
Foliar Use: Cutworms Tobacco Thrips Soybean (banded) thrips	0.025 to 0.1	1.3 to 5	CYPERMETHRIN 2.5 EC may be applied in water or refined vegetable oil. When water is used, apply a minimum of one gallon of finished spray per acre by air or five gallons of finished spray with ground equipment.
Foliar Use: Boll Weevil Cabbage Looper Cotton Bollworm Cotton Fleahopper Cotton Leaf Perforator European Corn Borer Fall Armyworm Lygus Bugs Other Plant Bugs Pink Bollworm Saltmarsh Caterpillar Tarnished Plant Bug Tobacco Budworm White Flies* Yellow Striped Armyworm	0.04 to 0.1	2 to 5	When applying in water by air, one quart of emulsified oil may be substituted for one quart of water in the finished spray. When using oil, use a minimum of one quart per acre in the finished spray. CYPERMETHRIN 2.5 EC may be injected into overhead sprinkler irrigation water provided 1) an anti-backflow check valve is present between the injection port and the water source, 2) a check valve is present in the line to prevent irrigation water from entering the chemical supply tank and 3) the irrigation injection system has interlocking on-off switches.
Foliar Use: Beet Armyworm**	0.06 to 0.1	3 to 5	For boll weevil control, apply CYPERMETHRIN 2.5 EC at a 3 to 4 day interval until pest numbers are reduced to acceptable levels. A maximum of 0.6 lb. active ingredient may be applied per acre per season. Do not graze or feed cotton for forage.

\*Aids in control

\*\*For control of beet armyworms only in the high plains of Texas, Arizona, and California.


**HEAD AND STEM BRASSICA VEGETABLES INCLUDING: BROCCOLI, CHINESE BROCCOLI (GAI LON, WHITE FLOWERING BROCCOLI); BRUSSELS SPROUTS; CAULIFLOWER; CAVALO BROCCOLO; KOHLRABI; CABBAGE; CHINESE CABBAGE (NAPA) (TIGHT-HEADING VARIETIES ONLY) (1 DAY PHI)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL. OZ./A	
Cutworms Corn Earworm Tobacco Budworm Lygus Bugs Saltmarsh Caterpillar Leafhoppers Flea Beetles Imported Cabbageworm Cucumber Beetles Aphids* Whiteflies*	0.05 to 0.1	2.5 to 5	Apply in water as necessary for insect control using a minimum of 15 gallon of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of CYPERMETHRIN 2.5 EC should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. Do not make applications less than 7 days apart.
Armyworms Loopers Stinkbugs Crickets Ground Beetles Onion Thrips Wireworm (adults)	0.075 to 0.1	3.75 to 5	A maximum of 0.6 lb. active ingredient may be applied per acre per season. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. Follow appropriate spray drift precautions. *Aids in control

8/11

**LEAFY BRASSICA GREENS INCLUDING: BROCCOLI RAAB (RAPINI); CHINESE CABBAGE (BOK CHOY); CHINESE MUSTARD CABBAGE (GAI CHOY); COLLARDS; KALE; MIZUNA; MUSTARD GREENS; MUSTARD SPINACH; RAPE GREENS (1 DAY PHI)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL. OZ./A	
Cutworms Corn Earworm Tobacco Budworm Lygus Bugs Saltmarsh Caterpillar Leafhoppers Flea Beetles Imported Cabbageworm Cucumber Beetles Aphids* Whiteflies*	0.05 to 0.1	2.5 to 5	Apply in water as necessary for insect control using a minimum of 15 gallon of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of CYPERMETHRIN 2.5 EC should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations. Do not make applications less than 7 days apart.
Armyworms Loopers Stinkbugs Crickets Ground Beetles Onion Thrips Wireworm (adults)	0.075 to 0.1	3.75 to 5	A maximum of 0.4 lb. active ingredient may be applied per acre per season. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. Follow appropriate spray drift precautions.  *Aids in control





9/11

**LETTUCE, HEAD (5 DAY PHI)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL. OZ./A	
Corn Earworm Cucumber Beetles Cutworms Diamondback Moth Flea Beetles Imported Cabbageworm Leafhoppers Lygus Bugs Saltmarsh Caterpillar Tobacco Budworm	0.05 to 0.1	2.5 to 5	Apply in water as necessary for insect control using a minimum of 15 gallons of finished spray per acre with ground equipment and 5 gallons per acre by air. Lower rates of CYPERMETHRIN 2.5 EC should be used under light to moderate insect pressure. Higher rates should be used to control heavy to extremely heavy insect populations.
Armyworms Crickets Loopers Onion Thrips Stinkbugs	0.075 to 0.1	3.75 to 5	A maximum of 0.6 lb. active ingredient may be applied per acre per season. In areas where arid climatic conditions persist, such as California and Arizona, higher than minimum recommended rates may be required. Follow appropriate spray drift precautions.

**BULB VEGETABLES (ALLIUM SPP.) INCLUDING: GARLIC; GARLIC, GREAT-HEADED (ELEPHANT); GREEN ESCHALOTS; JAPANESE BUNCHING ONIONS; LEEKS; ONION, DRY BULB AND GREEN; ONION, WELCH; SHALLOTS, DRY BULB AND GREEN; SPRING ONION OR SCALLIONS (7 DAY PHI)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL. OZ./A	
Onion Thrips	0.08 to 0.1	4 to 5	Apply CYPERMETHRIN 2.5 EC in a minimum of 20 gallons per acre with ground equipment or in a minimum of 3 gallons per acre by aircraft. Begin applications when pests appear. Do not make applications less than 7 days apart. To control Onion Thrips: Use higher rates as population increases and avoid rescue situations. Use of a crop oil concentrate at 16 fl. oz./A is recommended. A maximum of 0.5 lb. active ingredient may be applied per acre per season. Do not graze livestock in treated areas or cut treated crops for feed. Follow appropriate spray drift precautions.
Aphids Armyworms Cutworms Leafminers Onion Maggot Adults Stinkbugs	0.04 to 0.1	2.5 to 5	

**PECANS (21 DAY PHI)**

PEST	DOSAGE		REMARKS
	LB/A/A	FL. OZ./A	
Black Pecan Aphid Hickory Shuckworm Pecan Nut Casebearer Pecan Weevil Yellow Pecan Aphid	0.06 to 0.1	3 to 5	Applications at the lower rate should be made when pest populations are low. Rates should be increased as the pest pressure increases. Apply by ground equipment to the point of drip. Use 100 gallons of dilute spray per acre for smaller trees. For larger trees which require higher gallonage to achieve adequate coverage, apply in 200 to 300 gallons of water. In order to calculate the correct number of gallons of water needed to spray one acre of your trees, to the point of drip, you may need to conduct a test. If you do not know how to conduct such a test with your equipment, you should request assistance from your equipment dealer. Up to 0.6 lb. active ingredient per acre per season may be applied prior to shuck split. Do not graze livestock in treated orchards or cut treated cover crops for feed.

**DEALERS SHOULD SELL IN ORIGINAL PACKAGES ONLY.**

10/11

## **STORAGE AND DISPOSAL**

### **PESTICIDE STORAGE**

Do not store below 10°F (-12°C).

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food or feed by storage or disposal.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

To confine spill: If liquid, dike surrounding area or absorb with sand, cat litter or commercial clay. If dry material, cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

### **PESTICIDE DISPOSAL**

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

### **CONTAINER DISPOSAL**

Metal containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Do not cut or weld metal containers.

Plastic containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Returnable/Refillable Sealed Containers: Do not rinse container. Do not empty remaining formulated product. Do not break seals. Return intact to point of purchase.

11/11

---

**IMPORTANT INFORMATION  
READ BEFORE USING PRODUCT**

---

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorous, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorous Inc. and Seller harmless for any claims relating to such factors.

United Phosphorous, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorous, Inc., and Buyer and User assume the risk of any such use. **UNITED PHOSPHOROUS, INC. MAKES NEITHER WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

In no event shall United Phosphorous, Inc. or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHOROUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE, AT THE ELECTION OF UNITED PHOSPHOROUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT, OR COMPENSATION LIMITED TO DAMAGES NOT EXCEEDING THE FAIR MARKET PURCHASE PRICE, AND SHALL NOT INCLUDE INCIDENTAL OR CONSEQUENTIAL DAMAGES.**

United Phosphorous, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorous, Inc.

**Cypermethrin 2.5 EC AG RUP 30.6%.doc  
DRAFT 08-22-2002. Not EPA approved.**

