

70506-44

09-11-2009

1/26



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Ms. Rebecca A. Clemmer
Regulatory Manager
United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

SEP 11 2009

RE: Notification of Alternate Brand Name: PROKoZ Surflan AS Specialty Herbicide and other minor label changes
EPA Registration Number: 70506-44
Date of Submission: July 3, 2009

Dear Ms. Clemmer:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated July 3, 2009, for the above mention product. The Registration Division (RD) has conducted a preliminary screen of this request for its applicability under PRN 98-10 and finds that the action requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, please call me directly at 703-305-6249 or Joyce Edwards of my staff at 703-308-5479.

Sincerely,

A handwritten signature in black ink, appearing to read "Linda Arrington".

Linda Arrington
Notifications & Minor Formulations Team Leader
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
Washington, DC 20460

Registration
Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 70506-44	2. EPA Product Manager J. Miller	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) United Phosphorus, Inc/Surflan AS Specialty herbicide	PM # 23	
5. Name and Address of Applicant (Include ZIP Code) United Phosphorus, Inc. 630 Freedom Business Center, Suite 402 King of Prussia, PA 19406 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input checked="" type="checkbox"/> Notification - Explain below	<input type="checkbox"/> Other - Explain below

NOTIFICATION
SEP 11 2009

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)
Notification of alternate brand name (PROKoZ Surflan AS Specialty herbicide) and update of company and emergency contact information. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section III

1. Material This Product Will be Packaged in:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
*Certification must be submitted		If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of label directions <input type="checkbox"/> On Label <input type="checkbox"/> On Label accompanying product	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph	<input type="checkbox"/> Paper glued	<input type="checkbox"/> Stenciled	<input type="checkbox"/> Other _____

Section IV

1. Contact Person (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Rebecca A. Clemmer	Title Regulatory Manager	Telephone No. (Include Area Code) 610-491-2828
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		6. Date Application Received (Stamped) o
2. Signature <i>R.A. Clemmer</i>	3. Title Regulatory Manager	
4. Typed Name Rebecca A. Clemmer	5. Date July 3, 2009	



United Phosphorus, Inc.

630 Freedom Business Center
Suite 402
King of Prussia, PA 19406
(610) 491-2828 (phone)
(610) 491-2810 (fax)

Rebecca A. Clemmer
Regulatory Manager

July 3, 2009

Document Processing Desk (NOTIF)
Office of Pesticide Programs (H7504P)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., N.W.
Washington, D.C., 20460

Re: Surflan AS Specialty herbicide (EPA Reg. No. 70506-44)
Notification of Alternate Brand Name

To Whom It May Concern:

United Phosphorus, Inc. is notifying the Agency that this product may be marketed under the name PROKoZ Surflan AS Specialty herbicide in addition to its basic name. This alternate brand name is in addition to and does not replace the original product name.


The only other changes made to this label are the update of company and emergency contact information, the addition of the word 'exist' in the PPI washing instructions, and a trademark reference at the end of the label

In support of this Notification, enclosed please find EPA form 8570-1, and a copy of the label marked to show the changes

Please contact me if you have any questions.

Very truly yours,

Rebecca A. Clemmer
rebecca.clemmer@uniphos.com



4
26

PROKoZ[®] Surflan[®] AS Specialty herbicide

A selective preemergence surface-applied herbicide for control of annual grasses and many broadleaf weeds in:

- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Drainage Areas Under Shadehouse Benches
- Ornamental Bulbs
- Ground Covers/Perennials
- Christmas Tree Plantations
- Non-bearing fruit and nut trees and non-bearing vineyards
- Noncropland and Industrial Sites
- Established Warm Season Turf (including Bahiagrass, Bermudagrass, Buffalograss, Centipedegrass, St. Augustinegrass and Zoysiagrass)
- Tall Fescue (warm season areas)

Active Ingredient:

oryzalin: 3,5-dinitro- <i>N</i> ⁴ <i>N</i> ⁴ -dipropylsulfanilamide	40.4%
Inert Ingredients	59.6%
Total	100.0%

NOTIFICATION

SEP 11 2009

Contains 4.0 pounds of active ingredient per gallon.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before using this product, read Conditions of Sale and Limitation of Warranty and Liability at end of label booklet. If terms are unacceptable, return at once unopened.**

For emergency medical assistance, call the ~~National Pesticide Information Center~~ Rocky Mountain Poison Control Center at 1-800-858-7378/673-6671. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

Shake Well Before Using.


EPA Reg. No. 70506-44

EPA Est. No.



UPI

United Phosphorus, Inc.
423 Riverview Plaza 630 Freedom Business Center
Trenton, NJ 08611 King of Prussia, PA 19406
1-800-247-4557/438-6071 • www.upi-usa.com



5
28

(Page 1 through end):

Table of Contents	Page
Precautionary Statements	-
Hazards to Humans and Domestic Animals	-
First Aid	-
Environmental Hazards	-
Directions for Use	-
Agricultural Use Requirements	-
Storage and Disposal	-
General Information	-
Soil Preparation	-
Mixing Directions	-
Surflan AS Alone	-
Surflan AS Tank Mix Combinations	-
Mixing Order	-
Premixing	-
Equipment Cleaning	-
Activation and Cultivation	-
Weeds Controlled by Surflan AS	-
Weeds Suppressed by Surflan AS	-
Crop Specific Use Directions	-
Ornamental Plantings	-
Broadcast Application Rates	-
Tank Mix Combinations	-
Special Use Precautions	-
Recommended Species	-
Field and Container Grown Species	-
Non-bearing Trees and Vines	-
Ornamental Bulbs	-
Broadcast Application Rates	-
Special Use Precautions	-
Shadehouse Areas	-
Christmas Tree Plantations	-
Surflan AS Alone	-
Broadcast Application Rates	-
Tank Mix Combinations	-
Special Use Precautions	-
Noncropland Areas and Industrial Sites	-
Noncropland Areas--Tank Mix Combinations	-
Broadcast Application Rates	-
Industrial Sites--Tank Mix Combinations	-
Warm Season Turfgrasses	-
Annual Grasses Controlled by Surflan AS	-
Annual Broadleaf Weeds Controlled by Surflan AS	-
Broadleaf Weeds Suppressed by Surflan AS	-
Application Rates, Frequency and Timing of Application	-
Summer Annual Grasses and Broadleaf Weeds	-
Single Application Program	-
Split Application Program	-
Annual Bluegrass (<i>Poa annua</i>) and Winter Annual Broadleaf Weeds	-
Broadcast Application Rates	-
Weed Control in Florida	-
Application Equipment	-
Reseeding	-
Special Use Precautions	-
Conditions of Sale & Limitation of Warranty and Liability	-

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Causes Eye Irritation • Prolonged or Frequently Repeated Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Mixers and loaders must wear a chemical-resistant apron in addition to other PPE.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the ~~National Pesticide Information Center~~ Rocky Mountain Poison Control Center at 1-800-858-7378/673-6671 for emergency medical treatment information.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spills.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

7
26

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. **Exception:** If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Workers may enter treated areas without required PPE during the reentry interval following 1/2 to 1 inch of rainfall or irrigation, if they are performing tasks that do not involve contact with the soil subsurface; otherwise, PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Keep all persons, children and pets out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose of as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Surflan AS herbicide is a preemergence surface-applied product for the control of many annual grasses and broadleaf weeds in ornamental plantings, bulbs, ground covers/perennials, established warm-season turfgrass, Christmas tree plantations, non-bearing trees and vines, and noncropland and industrial sites.

Surflan AS is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Blazon or removed by spraying surface with water or washing with an industrial cleaner immediately after application. Surflan AS may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

Treatment of Plant Species Not Listed on the Label for Surflan AS

Users who wish to use Surflan AS on plant species not recommended on this label may determine the suitability for use by treating a small number of such plants at a recommended rate. Prior to treatment of larger areas, the treated plants should be observed for any sign of herbicidal injury during 30-60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. The user assumes responsibility for any plant damage or other liability resulting from use of Surflan AS on plant species not recommended on this label.

Aerial Application: Do not aerially apply this product.

Chemigation: Do not apply this product through any type of irrigation system.

For orchard crops, including citrus, pome fruits, stone fruits, and tree nuts, apply product only as a strip treatment in the tree rows; do not apply to row middles or drive rows.

Do not graze or feed forage from treated areas to livestock.

Precaution: Avoid spray drift to non-target areas when applying Surflan AS. Spray drift may result in reduced emergence of non-target plants adjacent to the treated area. Poor weed control may result if directions are not followed. Over-application may result in crop injury or excessive soil residue.

Application

Soil Preparation

Surflan AS controls weeds growing from seed. Surflan AS will not control emerged weeds. Surflan AS does not control established weeds; weeds growing from stolens, rhizomes, or root pieces. Therefore, areas to be treated should be free of emerged weeds. Weed residues, prunings, and trash should be thoroughly mixed into the soil or removed prior to treatment. In field applications, the soil should be in good till and free of clods at the time of application.

Ground Application: Apply Surflan AS as a directed spray to the soil surface or over the top of plants. Use only a properly calibrated, low-pressure, herbicide sprayer that will apply the spray uniformly. Use screens no finer than 50 mesh in nozzles and in-line strainers. Apply the appropriate rate of Surflan AS, as outlined in "Crop Specific Use Directions" section of this label. In all cases, use sufficient water volume to obtain uniform coverage and deliver the desired rate of Surflan AS to the treated area. The volume of water used is not critical, as long as the desired rate of Surflan AS is delivered uniformly across the area treated. When calibrating, determine the volume of water delivered by the sprayer to a given area (1,000 sq ft, 1 acre, etc.). Then mix the desired rate of Surflan AS in the amount of water required to cover the entire area to be treated. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to ensure proper calibration and uniform application. Maintain continuous agitation from mixing through application. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application.

Hand Held or Backpack Sprayer Application: The amount of water used to apply Surflan AS herbicide is not critical, but should be sufficient for uniform coverage of the target area. Calibrate by determining the volume of water required to treat 1000 square feet. Use this calibration volume to determine the amount of water and Surflan AS herbicide needed to treat the target area (see the following calibration example).

Note: Sprayer calibration (volume of spray needed to treat 1,000 square feet) will vary with each individual operator.

Steps in Calibration:

1. Mark an area of 1,000 square feet (i.e. 20 by 50 feet, or 25 by 40 feet).
2. Place the sprayer on a level surface and add water noting the final level of water in the spray tank.
3. Spray the marked area with a sufficient volume of water to provide uniform coverage. Refill the sprayer to the same level as before measuring the amount of water added. The measured water added to the sprayer is the volume needed to cover 1,000 square feet.

4. Determine the application rate (fl oz/1000 sq ft) for Surflan AS from the "Crop Specific Use Directions" section of this label.
5. To each volume of water used, as measured in step 3, add the amount of Surflan AS as determined in step 4.

Example: If the sprayer used 2 gallons of water to cover 1,000 square feet and the desired application rate of Surflan AS is 3 fluid oz/1,000 square feet, then you would add 3 fluid ounces of Surflan AS to every 2 gallons of water to be used.

Mixing Directions

Shake Well Before Using

Precaution: Do not allow the spray mixture to siphon back into water source.

Surflan AS - Alone

Make sure spray tank is clean and use only clean water. Fill spray tank 1/2 - 3/4 full. Start agitation and add the required amount of Surflan AS. Continue agitation and finish filling the spray tank. Maintain continuous agitation until application is completed.

Surflan AS - Tank Mix Combinations

Prior to mixing, read and carefully follow all label instructions and precautions for each product added to the tank mixture. Vigorous, continuous agitation is required for all tank mixes of Surflan AS. Sparger pipe agitators generally provide the best agitation in spray tanks.

Mixing Order: Fill the tank 3/4 full with clean water. Start agitation and add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: dry flowables (DF); wettable powders (WP); Surflan AS and other aqueous suspensions (AS), flowables (F), and liquids (L); solutions (S); and emulsifiable concentrates (EC).

Continue agitation and finish filling the spray tank with clean water. Maintain agitation until application is completed. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be completely resuspended before spraying is continued. A sparger agitator is particularly useful for this purpose.

Premixing: When tank mixing, initial mixing and dispersion of certain dry flowable or wettable powder products may be improved by premixing with water (slurrying). Adding the slurried material to the spray tank through a wetting screen of 20 or 35 mesh will help assure good initial dispersion.

Equipment Cleaning

If a buildup of material occurs on the walls of the spray tank, it should be removed between fillings by washing with soap and water and rinsing thoroughly. Tanks, lines, screens, and nozzles should be cleaned thoroughly after each use.

Activation and Cultivation

Surflan AS will remain stable on the soil surface up to 21 days following application. In the absence of timely rainfall, irrigation can be used to activate Surflan AS. A minimum of one-half (1/2) inch of rain or its equivalent in sprinkler irrigation is necessary to activate Surflan AS. If weeds begin to emerge due to lack of rainfall or irrigation, shallow cultivate 1-2 inches deep to destroy existing weeds, or remove them by hand. Shallow cultivation to a depth of 1-2 inches will enhance herbicidal effectiveness. Erratic weed control may result if Surflan AS is not activated by rainfall, irrigation, or cultivation within 21 days of application, or existing weeds have not been removed.

Weeds Controlled by Surflan AS

Annual Grasses:

Common Name	Scientific Name
-------------	-----------------

barley, little	<i>Hordeum pusillum</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
bluegrass, annual	<i>Poa annua</i>
crabgrass, large	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
crowfootgrass	<i>Dactyloctenium aegyptium</i>
cupgrass, southwestern	<i>Eriochloa gracilis</i>
foxtail, bristlegrass	<i>Setaria magna</i>
foxtail, giant	<i>Setaria faberi</i>
foxtail, green (pigeongrass)	<i>Setaria viridis</i>
foxtail, robust	<i>Setaria robusta</i>
foxtail, yellow	<i>Setaria glauca</i>
goosegrass (silver crabgrass)	<i>Eleusine indica</i>
Johnsongrass (seedling only)	<i>Sorghum halepense</i>
junglerice	<i>Echinochloa colonum</i>
lovegrass, Mexican	<i>Eragrostis mexicana</i>
lovegrass, orcutt	<i>Eragrostis orcuttiana</i>
oat, wild	<i>Avena fatua</i>
panicum, browntop	<i>Panicum fasciculatum</i>
panicum, fall (spreading panicgrass)	<i>Panicum dichotomiflorum</i>
panicum, Texas (buffalograss) (Coloradograss)	<i>Panicum texanum</i>
ryegrass, Italian	<i>Cenchrus incertus</i>
signalgrass (Brachiaria)	<i>Brachiaria</i> spp.
sprangletop, red	<i>Leptochloa filiformis</i>
witchgrass	<i>Panicum capillare</i>

Broadleaf Weeds:

Common Name	Scientific Name
bittercress	<i>Cardamine oligosperma</i>
carpetweed	<i>Mollugo verticillata</i>
chickweed, common	<i>Stellaria media</i>
fiddleneck, coast	<i>Amsinckia intermedia</i>
filaree, redstem	<i>Erodium cicutarium</i>
filaree, whitestem	<i>Erodium moschatum</i>
groundsel, common	<i>Senecio vulgaris</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambquarters	<i>Chenopodium album</i>
pigweed, prostrate	<i>Amaranthus blitoides</i>
pigweed, redroot	<i>Amaranthus retroflexus</i>
pigweed, spring	<i>Amaranthus hybridus</i>
pigweed, tumble	<i>Amaranthus albus</i>
puncturevine	<i>Tribulus terrestris</i>
purslane, common	<i>Portulaca oleracea</i>
pusley, Florida (Florida purslane) (Mexican clover) (pusley)	<i>Richardia scabra</i>
rocket, London	<i>Sisymbrium irio</i>
rockpurslane, desert	<i>Calandrinia ciliata</i>

11
26

shepherdspurse
spurge, prostrate
woodsorrel, yellow

Capsella bursa-pastoris
Euphorbia humistrata
Oxalis stricta

Weeds Suppressed by Surflan AS

Control of the following weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, and amount and timing of soil moisture:

Common Name	Scientific Name
horseweed	<i>Conyza canadensis</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca serriola</i>
mallow, common	<i>Malva neglecta</i>
milkweed, climbing	<i>Sarcostemma cynanchoides</i>
morningglory	<i>Ipomoea</i> spp.
mustard, black	<i>Brassica nigra</i>
mustard, wild	<i>Brassica kaber</i>
nightshade, black	<i>Solanum nigrum</i>
ragweed, common	<i>Ambrosia artemisiifolia</i>
smartweed	<i>Polygonum pennsylvanicum</i>
sowthistle, annual	<i>Sonchus oleraceus</i>
spurge, spotted	<i>Euphorbia maculata</i>
teaweed (prickly sida)	<i>Sida spinosa</i>
velvetleaf	<i>Abutilon theophrasti</i>
wheat, volunteer	<i>Triticum</i> spp.

Crop Specific Use Directions

Ornamental Plantings

Surflan AS is recommended for use on certain landscape container- and field-grown established ornamental plants including: trees, shrubs, ground covers/perennials, flowers, non-bearing fruit and nut trees, non-bearing vineyards; and in the production of ornamental bulbs (See "Ornamental Bulbs" section for special use directions).

Broadcast Application Rates

Labeled Use Site	Length of Control	Surflan AS		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
		(qt/acre)	(fl oz/1000 sq ft)		
Landscape Ornamentals	2 - 4 months	2	1.5	2	8
	3 - 6 months	3	2.2	4	12
	4 - 8 months	4	3	4	12
Field-grown and container-grown ornamentals	2 - 4 months	2	1.5	3	8
	3 - 6 months	3	2.2	3	9
	4 - 8 months	4	3	3	12

Tank Mix Combinations

Tank mix combinations of Surflan AS plus glyphosate, and many other labeled herbicides may be used to control undesirable vegetation in ornamental areas. Surflan AS may also be tank mixed with Gallery* herbicide and applied preemergence to broaden the spectrum of broadleaf weed control in ornamental areas. Applied as directed, these tank mixes of Surflan AS will provide control of susceptible weed

species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Surflan AS Plus glyphosate: Tank mix combinations of Surflan AS plus glyphosate are recommended to control existing undesirable vegetation. Applied as directed, Surflan AS plus glyphosate will provide postemergence control of susceptible weed species listed on the label for glyphosate and residual preemergence control of susceptible weed species listed on the label for Surflan AS. Refer to the label for glyphosate for specific use directions, precautions, and limitations before use.

Precautions: Do not apply sprays containing glyphosate over the top of ornamental plants. Extreme care must be exercised to prevent sprays containing glyphosate from coming in contact with foliage and stems of turfgrasses, trees, shrubs, or other desirable vegetation since severe damage or death may result. If spraying with glyphosate in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

Special Use Precautions:

Apply only to established plants that have been transplanted into their growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Rooted liners should be removed from their original growing containers and placed in new containers at least two weeks prior to treatment or injury may occur.

To avoid possible injury, do not apply Surflan AS to:

- Nursery, forest, or Christmas tree: seedling beds, cutting beds, or transplant beds.
- Unrooted liners or cuttings that have been planted in pots for the first time.
- Pots less than four inches wide.
- Ground covers until they are established and well rooted.
- Ornamental plantings where there is likelihood of runoff onto lawn areas.
- Areas containing dichondra or cool season turfgrass species.

On container grown ornamentals where weed seed germination continues for extended periods of time, do not make repeat applications of Surflan AS for at least 90 days or crop injury may occur.

Applications of Surflan AS over the top of plants with newly forming buds may cause injury. In this situation a directed spray is recommended.

For soils treated with Surflan AS during the previous season, plant only the ornamental species listed on this label or injury may occur.

Ice Plant: When establishing unrooted ice plant on coarse-textured soils in landscape plantings, do not exceed the 2 quart per acre rate of Surflan AS or crop injury may occur.

Note: Injury on the following plant species has been observed following applications of Surflan AS and use is not recommended:

- Deutzia gracilis* (slender deutzia)
- Pseudotsuga menziesii* (Douglas-fir)
- Thuja occidentalis* 'Techny' (Techny arborvitae)
- Tsuga canadensis* (eastern hemlock)
- Begonia* spp. (begonia)
- Coleus hybridus* (coleus)

Surflan AS May be Used on the Following Established Plant Species:
(Note: Limitations on recommended treatment methods).

Trees

Recommended Treatment Method

F = Field Grown

C = Container Grown

Scientific Name

Common Name

13
26

Scientific Name	Common Name	
<i>Abies balsamea</i>	Fir, balsam	F
<i>Abies concolor</i>	Fir, white	F
<i>Abies fraseri</i>	Fir, fraser	F
<i>Abies grandis</i>	Fir, grand	F
<i>Abies veitchi</i>	Fir, Vietch	F
<i>Abies lasiocarpa</i>	Fir, alpine	F
<i>Abutilon hybridum</i>	Albus-flowering maple	F
	Luteus-flowering maple	F
	Roseus-flowering maple	F
	Tangerine-flowering maple	F
	Vesuvius red-flowering maple	F
<i>Acer ginnala</i>	Flame maple	F
<i>Acer rubrum</i>	Red sunset maple	F
<i>Acer saccharinum</i>	Silver maple	F
<i>Acer spp.</i>	Maple	F
<i>Alsophila australis</i>	Australian tree fern	C,F
<i>Arecastrum romanzoffianum</i>	Queen palm	F
<i>Betula nigra</i>	Birch, river	F
<i>Betula papyrifera</i>	Paper birch	F
<i>Betula pendula</i>	Birch, white	F
<i>Bucida buceras</i>	Black olive	F
<i>Carya spp.</i>	Pecan, ornamental	C,F
<i>Cedrus, atlantica</i>	Atlas cedar	C,F
<i>Cedrus deodara</i>	Deodar cedar	C,F
<i>Ceratonia siliqua</i>	Carob	F
<i>Cercidium floridum</i>	Palo Verde, blue	F
<i>Cercis canadensis</i>	Redbud	C,F
<i>Chamaecyparis lawsoniana</i>	Falsecypress, Lawson	F
<i>Chamaecyparis obtusa</i>	Filicoides-fernspray cypress	F
<i>Chamaecyparis pisifera</i>	Gracilis-slender Hinoki cypress	F
	Sawara-false cypress	F
	Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	Cat Palm	F
<i>Chamaedorea costaricana</i>	Palm	F
<i>Chamaedorea elegans</i>	Parlor palm	F
<i>Citrus spp.</i>	Citrus, ornamental	C,F
<i>Cornus florida</i>	Dogwood, flowering	F
<i>Cryptomeria japonica</i>	Cryptomeria, Japanese	C,F
<i>Cupaniopsis anacardioides</i>	Carrot wood	F
<i>Cupressus arizonica (glabra)</i>	Cypress, Arizona	C,F
<i>Cupressus glabra</i>	Arizona cypress	C,F
<i>Cupressocyparis leylandii</i>	Leyland cypress	C,F
<i>Cupressus sempervirens</i>	Cypress, Italian	C,F
<i>Dicksonia antarctica</i>	Tasmanian tree fern	C,F
<i>Elaeagnus angustifolia</i>	Russian olive	C,F
<i>Eucalyptus camaldulensis</i>	Red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	Eucalyptus, mealy	F
	Silver dollar eucalyptus	F
<i>Eucalyptus nicholii</i>	Eucalyptus, narrow-leaved	F
<i>Eucalyptus sideroxylon</i>	Eucalyptus, red ironbark	F
<i>Ficus benjamina</i>	Ficus	F
<i>Fraxinus spp.</i>	Ash	F
<i>Ginkgo biloba</i>	Ginkgo (Maidenhair tree)	C,F
<i>Gleditsia triacanthos</i>	Honey locust	F
<i>Heteromeles arbutiflora</i>	Toyon	F
<i>Juniperus virginiana</i>	Redcedar, Eastern	F
<i>Koelreuteria paniculata</i>	Goldenrain tree	F

Scientific Name	Common Name	
<i>Liquidambar styraciflua</i>	Sweetgum, American	C,F
<i>Magnolia</i> spp.	Magnolia	F
<i>Malus</i> spp.	Crabapple	F
<i>Morus alba</i>	White mulberry	F
<i>Picea abies</i>	Pendula-weeping Norway spruce	F
	Repens-spreading Norway spruce	F
	Spruce, Norway	F
<i>Picea englemanni</i>	Spruce, Englemann	F
<i>Picea glauca</i>	Spruce, white	F
	Conica-dwarf Alberta spruce	F
<i>Picea glauca conica</i>	Dwarf Alberta spruce	F
<i>Picea mariana</i>	Spruce, black	F
<i>Picea pungens</i>	Glauca-Colorado blue spruce	F
	Hoopsii-Hoop's blue spruce	F
	Koster-Koster blue spruce	F
	Spruce, Colorado	C,F
<i>Pinus aristata</i>	Bristlecone pine	F
<i>Pinus canariensis</i>	Canary Island pine	F
<i>Pinus contorta</i>	Shore pine, beach pine	F
<i>Pinus eldarica</i>	Eldarica pine	F
<i>Pinus halepensis</i>	Aleppo pine	C,F
<i>Pinus radiata</i>	Monterey pine	F
<i>Pinus</i> spp.	Pine	C,F
<i>Pinus strobus</i>	Eastern white pine	F
<i>Pinus sylvestris</i>	Scotch pine	F
<i>Pinus thunbergiana</i>	Japanese black pine	F
<i>Platanus occidentalis</i>	American sycamore	F
<i>Platanus racemosa</i>	Californina sycamore	F
<i>Podocarpus</i> spp.	Podocarpus	F
<i>Populus deltoides</i>	Cottonwood	F
	Cottonwood (grown for pulp)	F
<i>Prunus caroliniana</i>	Laurelcherry, Carolina	F
<i>Prunus glandulosa</i>	Dwarf flowering almond	C,F
<i>Prunus laurocerasus</i>	Laurelcherry, English	F
<i>Prunus mahaleb</i>	Cherry, Mahaleb	F
<i>Prunus yedoensis</i>	Yoshino flowering cherry	F
<i>Pyrus communis</i>	Pear	F
<i>Quercus palustris</i>	Pin oak	F
<i>Quercus phellos</i>	Willow oak	F
<i>Quercus rubra</i>	Red oak	C,F
<i>Quercus</i> spp.	Oak	C,F
<i>Salix babylonica</i>	Babylon weeping willow	F
	Corkscrew willow	F
<i>Schinus molle</i>	California pepper tree	F
<i>Sequoia sempervirens</i>	Redwood, coast	F
<i>Sequoiadendron giganteum</i>	Giant sequoia	F
<i>Swietenia mahogani</i>	Mahogany	F
<i>Tabebuia caraiba</i>	Yellow tab	F
<i>Tilia cordata</i>	Linden, little leaf	C,F
<i>Ulmus parvifolia</i>	Chinese elm	F
<i>Umbellularia californica</i>	California laurel	F
<i>Washingtonia robusta</i>	Mexican fan palm	F

Shrubs**Recommended Treatment Method**

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Abelia grandiflora</i>	Glossy abelia	F
<i>Acacia redolens</i>	Acacia, prostrate	F
<i>Agave americana</i>	Century plant	F
<i>Agave macroculmis</i>	Agave	F
<i>Anisodontea hypomandarum</i>	Cape mallow	C,F
<i>Arctostaphylos stanfordiana</i>	Manzanita, Stanford	F
<i>Astilbe chinensis</i>	Astilbe/false spirea	C,F
<i>Baccharis pilularis</i>	Coyotebush	F
<i>Berberis thunbergii</i>	Aurea-golden Japanese barberry	C,F
	Crimson pygmy barberry	C,F
	Atropurea-redleaf Japanese barberry	C,F
	Barberry, Japanese	C,F
<i>Bougainvillea</i> spp.	Barbara Karst	F
	California gold	F
	Scarlet O'Hara	F
	Texas dawn	F
<i>Buddleia davidii</i>	Butterfly bush	C,F
<i>Buxus microphylla</i>	Littleleaf boxwood	F
<i>Buxus microphylla japonica</i>	Boxwood, Japanese	C,F
<i>Buxus sempervirens</i>	Boxwood, common	C,F
<i>Callistemon citrinus</i>	Bottlebrush, lemon	C,F
<i>Cassia artemisioides</i>	Cassia, feathery	F
<i>Ceanothus americanus</i>	Jerseytea, redroot	C,F
<i>Ceanothus</i> spp.	Wild lilac	C,F
<i>Chaenomeles japonica</i>	Flowering quince	C,F
<i>Chamaecyparis obtusa</i>	Kosteri cypress	F
	Nana-dwarf Hinoki cypress	F
	Torulosa cypress	F
<i>Chamaecyparis pisifera</i>	Squarrosa Minima cypress	F
<i>Chamaecyparis pisifera</i> spp.	Filifera-thread cypress	F
<i>Chrysalidocarpus lutescens</i>	Areca palm	F
<i>Clethra</i>	Summersweet	C,F
<i>Cleyera japonica</i>	Cleyera, Japanese	C,F
<i>Coleonema pulchrum</i>	Pink breath of heaven	C,F
<i>Cornus alba</i>	Sibirica-Siberian dogwood	F
<i>Cornus kousa</i>	Dogwood, kousa	C,F
<i>Cornus stolonifera</i>	Flaviramea-yellowtwig dogwood	F
<i>Cotoneaster adpressus</i>	Praecox-early cotoneaster	F
<i>Cotoneaster apiculatus</i>	Cotoneaster, cranberry	C,F
<i>Cotoneaster buxifolius</i>	Cotoneaster, brightbead	F
<i>Cotoneaster congestus</i>	Cotoneaster, Pyrenees	F
<i>Cotoneaster dammeri</i>	Cotoneaster, bearberry	C,F
<i>Cotoneaster himalayan</i>	Himalayan cotoneaster	F
<i>Cotoneaster horizontalis</i>	Cotoneaster, rock	C,F
<i>Cotoneaster lacteus</i>	Cotoneaster, parney	C,F
<i>Cotoneaster microphyllus</i>	Cotoneaster, rockspray	F
<i>Cotoneaster salicifolia</i>	Willowleaf cotoneaster	C,F
<i>Cytisus praecox</i>	Hollandia-warminster broom	F
<i>Cytisus scoparius</i>	Lena-Scotch broom	F
<i>Dasyliirion wheeleri</i>	Sotol, desert spoon	F
<i>Deutzia crenata</i>	Nakiana-dwarf deutzia	F
<i>Dodonaea viscosa</i>	Hopseedbush, clammy	F
	Hopseed bush	F
<i>Escallonia exoniensis</i>	Escallonia	C,F
<i>Euonymus alata</i>	Euonymus, winged	F
<i>Euonymus fortunei</i>	Canadale gold euonymus	C,F
	Emerald'n gold euonymus	C,F

Scientific Name	Common Name	
	Euonymus, stringybark	C,F
	Wintercreeper	C,F
<i>Euonymus japonica</i>	Euonymus, evergreen	C,F
	Silver king euonymus	F
<i>Euonymus kiatschovica</i>	Spreading euonymus	F
<i>Euonymus vegetus</i>	Bigleaf wintercreeper	C,F
<i>Fatsyhedera lizei</i>	Fatsyhedera	C,F
<i>Fatsia japonica</i>	Japanese aralia	C,F
<i>Felicia amelloides</i>	Blue marguerite	C,F
<i>Forsythia intermedia</i>	Forsythia, border	F
<i>Gardenia jasminoides</i>	Gardenia	C,F
<i>Genista pilosa</i>	Woadwaxen	F
<i>Hibiscus rosa-sinensis</i>	Ross Estey-hibiscus	F
	Hibiscus, Chinese	F
<i>Hibiscus syriacus</i>	Rose of Sharon, Red Bird	F
	Rose of Sharon, Red Heart	F
	Rose of Sharon, Woodbridge	F
	Rose-of-Sharon (Shrubalthea)	F
<i>Hydrangea macrophylla</i>	Hydrangea, French	C, F
<i>Hydrangea quercifolia</i>	Hydrangea, Oakleaf	C, F
<i>Ilex aquifolium</i>	Balkans holly	F
	Gold coast holly	F
	Holly, English	F
<i>Ilex aquipernyi</i>	San Jose holly	C,F
<i>Ilex cornuta</i>	Dwarf Burford holly	C,F
	Holly, Chinese	C,F
<i>Ilex crenata</i>	Compacta-dwarf Japanese holly	C,F
	Convexa holly	C,F
	Helleri-Heller's Japanese holly	C,F
	Holly, Japanese	C,F
<i>Ilex glabra</i>	Nordica-inkberry holly	F
<i>Ilex meserveae</i>	Blue boy holly	F
	Blue girl holly	F
	Ebony magic holly	F
<i>Ilex vomitoria</i>	Nana-dwarf yaupon holly	C,F
	Pendula-weeping yaupon holly	C,F
	yaupon holly	C,F
<i>Juniperus chinensis</i>	Media-old gold juniper	C,F
<i>Juniperus conferta</i>	Emerald sea shore juniper	F
<i>Juniperus horizontalis</i>	Huntington blue juniper	C,F
	Wiltonii-blue carpet juniper	C,F
<i>Juniperus procumbens</i>	Nana-dwarf Japanese garden juniper	C,F
<i>Juniperus prostrata</i>	Prostrata juniper	C,F
<i>Juniperus sabina</i>	Broadmoor juniper	F
	Foemina-Hicks juniper	F
	Tamariscifolia-Tam juniper	F
<i>Juniperus scopulorum</i>	Emerald green juniper	F
<i>Juniperus spp.</i>	Juniper	C,F
<i>Juniperus squamata</i>	Blue juniper	F
	Blue star juniper	F
	Parsonii juniper	F
<i>Justicia brandegeana</i>	Shrimp plant	C,F
<i>Justicia spicigera</i>	Honeysuckle, Mexican	F
<i>Kalmia latifolia</i>	Laurel, mountain	F
<i>Lagerstroemia indica</i>	Crape myrtle	C,F
<i>Lavandula angustifolia</i>	English lavender	C,F
<i>Leucothoe axillaris</i>	Leucothoe, coast	F

17
26

Scientific Name	Common Name	
<i>Leucothoe fontanesiana</i>	Leucothoe, drooping	F
<i>Ligustrum amurense</i>	Privet, amur	C,F
<i>Ligustrum japonicum</i>	Privet, Japanese	C,F
	yellow tip ligustrum	C,F
<i>Ligustrum lucidum</i>	Privet, glossy	C,F
<i>Ligustrum ovalifolium</i>	California privet	F
<i>Ligustrum texanum</i>	Howardi privet	F
	Wax leaf privet	F
<i>Ligustrum vicaryi</i>	Privet, golden	C,F
	Vicary golden privet	C,F
<i>Livistona chinensis</i>	Chinese fountain palm	F
<i>Lonicera fragrantissima</i>	Winter honeysuckle	F
<i>Lonicera periclymenum</i>	Flowering woodbine	F
	Serotina woodbine	F
<i>Lonicera sempervirens</i>	Trumpet honeysuckle	F
<i>Lorpetalum chinense</i>	(No common name)	C,F
<i>Mahonia aquifolium</i>	Oregon grape	F
<i>Myoporum parvifolium</i>	Myoporum, prostrate	F
<i>Myrtus communis</i>	Myrtle, true	C,F
<i>Nandina domestica</i>	Compacta-dwarf heavenly bamboo	C,F
	Harbour dwarf-heavenly bamboo	C,F
	Heavenly bamboo (Nandina)	C,F
	Nana compacta-heavenly bamboo	C,F
	Nana purpurea-heavenly bamboo	C,F
	Woods dwarf-heavenly bamboo	C,F
<i>Nerium oleander</i>	Hardy red oleander	C,F
	Oleander	C,F
	Ruby lace oleander	C,F
<i>Osmanthus heterophyllus</i>	Osmanthus, holly-leaf	F
<i>Pachysandra terminalis</i>	Japanese spurge	C,F
<i>Philadelphus spp.</i>	Mockorange	C,F
<i>Phoenix roebelenii</i>	Pigmy date palm	F
<i>Photinia fraseri</i>	Fraser's photinia	C,F
	Photinia	C,F
<i>Pieris japonica</i>	Lily-of-the-valley	F
	Snowdrift lily-of-the-valley	F
	Temple bells lily-of-the-valley	F
	Valley rose lily-of-the-valley	F
	Andromeda	C,F
<i>Pittosporum spp.</i>	Pittosporum	C,F
<i>Pittosporum tobira</i>	Green pittosporum	F
	Japanese pittosporum	F
	Tobira	F
	Wheeler's dwarf pittosporum	F
<i>Platycladus orientalis</i>	Arborvitae, Oriental	C,F
<i>Plumbago ariculata</i>	Blue cape plumbago	F
<i>Podocarpus macrophyllus</i>	Yewpine	C,F
<i>Potentilla fragiformis</i>	Cinquefoil	F
<i>Potentilla fruticosa</i>	Cinquefoil	C,F
<i>Protea neriifolia</i>	Protea	F
<i>Pyracantha coccinea</i>	Firethorn, scarlet	C,F
<i>Pyracantha fortuneana</i>	Lolondei Monrovia pyracantha	C,F
<i>Pyracantha fortuneana</i>	Monon pyracantha	C,F
	Red elf hybrid pyracantha	C,F
	Rutgers hybrid pyracantha	C,F
	Santa Cruz pyracantha	C,F
	Victory pyracantha	C,F

Scientific Name

Pyracantha skoidzumi
Pyracantha, fortuneana
Raphiolepis indica

Common Name

Firethorn, formosa C,F
 Firethorn C,F
 Enchantress-Moness raphiolepis F
 Raphiolepis (India hawthorn) C,F
 Springtime-Monme raphiolepis F

Raphiolepis ovata

Roundleaf raphiolepis F
 Eastercactus C,F

Rhipsalidopsis gaertneri

Rhododendron calendulaceum

Flame azalea F

Rhododendron campylocarpum

Butterfly rhododendron F

Rhododendron carolinianum x daurium

PJM rhododendron F

Rhododendron catawbiense

Catawba album rhododendron C,F

Catawba rhododendron C,F

Lord Roberts rhododendron C,F

Rocket rhododendron C,F

Rhododendron forrestii x griersonianum

Elizabeth rhododendron F

Rhododendron hybrid spp.

America rhododendron F

English Roseum rhododendron F

Nova Zembla rhododendron F

Scintillation rhododendron F

Rhododendron impeditum

Rhododendron F

Rhododendron indica

Formosa azalea C,F

Waucabusa azalea C,F

Rhododendron kerume

Coral bells azalea C,F

Hino crimson azalea C,F

Hino pink azalea C,F

Snow azalea C,F

Rhododendron maximum

Rhodie max (rosebay) C,F

Rhododendron mucronulatum

Rhododendron F

Rhododendron satuski

Gumpo pink azalea F

Higasa azalea F

Rhododendron spp.

Azalea C,F

Rhododendron C,F

Rhododendron spp. hybrids

Carror azalea C,F

Girard Roberta azalea F

Golden flare exbury azalea F

Rhus lancea

Sumac, African C,F

Rosa rugosa

Ramanas rose F

Rosmarinus officinalis

Rosemary F

Senecio cineraria

Dusty miller C,F

Spiraea vanhouttei

Bridal wreath F

Syringa vulgaris

Lilac, common C,F

Syzygium paniculata

Brush cherry C,F

Taxus cuspidata

Yew, Japanese F

Taxus media

Yew F

Thuja occidentalis

Arborvitae, American C,F

Emerald arborvitae F

Globosa-globe arborvitae F

Little giant-dwarf arborvitae F

Nigra-dark American arborvitae F

Pyramidalis arborvitae F

Rheingold arborvitae F

Woodwardii arborvitae F

Thuja orientalis

Aureus nana-dwarf golden arborvitae F

Minima glauca-dwarf arborvitae F

Thuja plicata

Red Cedar, Western F

Trachelospermum jasminoides

Star jasmine, Chinese F

Veitchia merrilli

Christmas palm F

19
26

Scientific Name	Common Name	
<i>Viburnum carlesii</i>	Koreanspice viburnum	C,F
<i>Viburnum davidii</i>	David viburnum	F
<i>Viburnum japonicum</i>	Viburnum	F
<i>Viburnum judd</i> (V X Judii)	Viburnum	C,F
<i>Viburnum opulus sterile</i>	Common snowball viburnum	F
<i>Viburnum plicatum tomentosum</i>	Doublefile viburnum	F
<i>Viburnum setigerum</i>	Tea viburnum	F
<i>Viburnum suspensum</i>	Virbumum, Sandankwa	F
<i>Viburnum tinus</i>	Viburnum, Laurustinus	C,F
	Compactum-spring bouquet viburnum	F
<i>Viburnum tinus compactum</i>	Spring bouquet viburnum	F
<i>Viburnum trilobum compactum</i>	Dwarf cranberry bush	F
<i>Viburnum x pragense</i>	Viburnum	F
<i>Weigela florida</i>	Bristol ruby weigela	F
	Java red weigela	F
	Minuet weigela	F
	Weigela, oldfashioned	F
<i>Xylosma congestum</i>	Xylosma	F
<i>Yucca elata</i>	Yucca, soaptree	C,F
<i>Yucca recurvifolia</i>	Yucca, pendulous	F

Groundcovers/Perennials**Recommended Treatment Method**

F = Field Grown

C = Container Grown

Scientific Name	Common Name	
<i>Agapanthus africanus</i>	Lily-of-the-Nile	C,F
<i>Ajuga</i> spp.	Carpet bugle	F
<i>Arctotheca calendula</i>	Cape weed	F
<i>Asparagus retrofractus</i>	(No common name)	C,F
<i>Asparagus varieegata</i>	Tree fern	C,F
<i>Aster novae-angliae</i>	New England aster	C,F
<i>Aster novi-belgii</i>	New York aster	C,F
<i>Athyrium nipponicum</i>	Japanese painter fern	C,F
<i>Brassica oleracea</i>	Wild cabbage	C,F
<i>Callistephus chinensis</i>	China aster	C,F
<i>Campanula elatines</i>	Bellflower	C,F
<i>Carpobrotus edulis</i>	Ice plant, largeleaf (see label)	F
<i>Clytostoma callistegioides</i>	Trumpet vine, violet	C,F
<i>Cortaderia selloana</i>	Pampas grass	F
<i>Cuphea hyssopifolia</i>	False Mexican heather	C,F
<i>Delosperma alba</i>	White iceplant	F
<i>Dietes vegeta</i>	Fortnight lily	C,F
<i>Digitalis mertonensis</i>	Foxglove	C,F
<i>Doronicum cordatum</i>	Leopard's bane	C,F
<i>Drosanthemum floribundum</i>	Trailing rosea iceplant	F
<i>Erianthus ravennae</i>	Hardy pampus grass	C,F
<i>Festuca ovina glauca</i>	Blue fescue	F
<i>Gaillardia grandiflora</i>	Blanket flower	C,F
<i>Gazania rigens leucolaena</i>	Gazania, trailing	C,F
<i>Gazania</i> spp.	Gazania	F
<i>Hedera canariensis</i>	Ivy, Algerian	F
<i>Hedera helix</i>	Ivy, English	F
<i>Heliotropium fragrans</i>	Common heliotrope	C,F
<i>Hemerocallis</i> spp.	Daylily	C,F
<i>Hosta lancifolia</i>	Albo-marginata hosta	C,F
<i>Hosta</i> spp.	Lily, plantain	C,F

Scientific Name*Heuchera micrantha**Hypericum* spp.*Iberis sempervirens**Lampranthus spectabilis**Leptospermum scaparium**Limonium perezii**Liriope gigantea**Liriope muscari**Lobelia erinus**Lonicera japonica**Mesembryanthemum crystallinum**Monarda didyma**Ophiopogon japonicus**Osteospermum fruticosum**Pachysandra terminalis**Pennisetum setaceum**Polystichum polyblepharum**Sedum brevifolium**Sedum kamtschaticum**Sedum spurium**Tulbaghia violacea**Verbena rigida**Veronica* spp.*Vinca major**Vinca minor***Common Name**

Coral bells

St. Johnswort

Evergreen candytuft

Trailing iceplant

New Zealand teatree/Manuka

Statice/Sea lavender

White lily turf

Lilac beauty lily turf

Majestic lily turf

Monroe white lily turf

Silvery sunproof lily turf

Variegated liriope lily turf

Big blue lily turf

Edging lobelia

Honeysuckle, Japanese

Ice plant (see label)

Bee Balm

Mondo grass

Daisy, trailing African

Japanese spurge

Fountaingrass

Tassel fern

Stonecrop

Stonecrop

Stonecrop, tworow

Society garlic

Veined verbena

Speedwell

Periwinkle, bigleaf

Periwinkle, dwarf

C,F

C,F

C,F

F

C,F

C,F

F

C,F

C,F

C,F

C,F

C,F

C,F

C,F

F

F

C,F

F

F

F

C,F

C,F

C,F

C,F

C,F

C,F

C,F

C,F

F

F

Flowers**Recommended Treatment Method****F = Field Grown****C = Container Grown****Scientific Name***Achillea* spp.*Antirrhinum majus**Caladium bicolor**Chrysanthemum* spp.

Mixed hybrid

*Cladium bicolor**Coreopsis lanceolata**Coreopsis verticulata**Dianthus barbatus**Dianthus gratianopolitanus**Dicentra spectabilis**Dimorphotheca* spp.*Echinacea purpurea**Evolvulus nuttallianus**Geum quellyon**Gladiolus hortulanus**Gypsophila paniculata**Impatiens wallerana**Iris* spp.*Liatris spicata**Pelargonium hortorum**Petunia* spp.**Common Name**

Yarrow

Snapdragon

Caladium, fancy leafed

Chrysanthemum

Dahlia

Fancy-leafed caladium

Coreopsis

Threadleaf coreopsis

Sweet William

Cheddar pink

Bleeding heart

Marigold, cape

Coneflower, purple

Blue daze

Geum

Gladiolus

Baby's breath

Impatiens (Busy lizzie)

Iris, bearded

Blazing star

Geranium

Petunia

C,F

F

F

C,F

C,F

F

F

C,F

F

C,F

C,F

F

C,F

C,F

F

F

F

F

F

C,F

F

C,F

21
26

Scientific Name	Common Name	
<i>Portulaca grandiflora</i>	Moss, rose	F
<i>Ranunculus asiaticus</i>	Ranunculus, Persian	F
<i>Rosa</i> spp.	Rose	F
<i>Rudbeckia fulgida</i>	Blackeyed susan	C,F
<i>Rudbeckia hirta</i>	Daisy, gloriosa (black-eyed Susan)	F
<i>Salvia</i> spp.	Salvia (Sage)	F
<i>Stokesia laevis</i>	Aster, stokes	F
<i>Strelitzia reginae</i>	Bird of paradise	F
<i>Tagetes</i> spp.	Marigold	F
<i>Viola wittrockiana</i>	Pansy	F
<i>Zinnia elegans</i>	Zinnia, common	F

Non-bearing Trees and Vines

Recommended Treatment Method

F = Field Grown

C = Container Grown

almond	F
apple	F
apricot	F
avocado	F
blackberry	F
blueberry	F
boysenberry	F
cherry, sour	F
cherry, sweet	F
currant	F
dewberry	F
elderberry	F
fig	F
filbert	F
gooseberry	F
grape, American	F
grape, European	F
grapefruit	F
kiwi	F
Kumquat	C,F
lemon	F
loganberry	F
macadamia nut	F
nectarine	F
olive	F
orange	C,F
peach	F
pear	F
pecan	C,F
pistachio	F
plum	F
pomegranate	F
prune	F
raspberry	F
walnut, black	F
walnut, English	F

† Non-bearing plants are defined as those that will not bear fruit for at least one year after treatment.

22
26

Ornamental Bulbs

Surflan AS may be applied for control of susceptible annual weeds in ornamental bulbs, e.g., bulbous iris, daffodil (narcissus), hyacinth, and tulip. Apply Surflan AS to the soil surface 2-4 weeks after planting, but prior to the emergence of annual weeds. For fall planted bulbs, apply Surflan AS again in late winter or early spring to weed-free soil surfaces.

Broadcast Application Rates

Time of Application	Soil Texture	Surflan AS		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
		(qt/acre)	(fl oz/1000 sq ft)		
Fall	Coarse	0.75	0.5	3	1.5
Fall	Medium and Fine	1.5	1.0	3	2.25
Feb. - March	All Soil Textures	0.75	0.5	3	2.25

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.
Do not apply to gladioli corms prior to emergence or less than one (1) inch in diameter.

Shadehouse Areas

Surflan AS may be applied to drainage areas under benches in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within 3 weeks prior to enclosure of greenhouse or poly-type structures.

Christmas Tree Plantations

Surflan AS Alone

Apply Surflan AS as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species, including fir (*Abies* spp.), pine (*Pinus* spp.), and spruce (*Picea* spp.). Follow all instructions provided in the "General Information" section of this label.

Broadcast Application Rates

Length of Control	Surflan AS		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
	(qt/acre)	(fl oz/1000 sq ft)		
2 - 4 months	2	1.5	2	8
4 - 8 months	4	3	2	8

Tank Mix Combinations

Tank mix combinations of Surflan AS plus other labeled herbicides may be used as directed or overtop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Surflan AS Plus glyphosate: Apply tank mix combinations of Surflan AS plus glyphosate only as directed sprays in Christmas tree plantings. When applied according to use directions, Surflan AS plus glyphosate will provide postemergence control of susceptible weed species listed on the label for glyphosate and residual preemergence control of susceptible weed species listed on the label for Surflan AS. Refer to the label for glyphosate for specific use directions, precautions, and limitations before use.

Special Use Precautions:

23
26

Do not apply to Douglas-fir (*Pseudotsuga menziesii*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plants that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation.

Noncropland Areas and Industrial Sites

Noncropland Areas - Tank Mix Combinations

Tank mix combinations of Surflan AS plus glyphosate and many other labeled herbicides may be used to control undesirable vegetation in noncropland areas such as roadsides, rights-of-way, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Broadcast Application Rates

Length of Control	Surflan AS		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (qt/acre)
	(qt/acre)	(fl oz/1000 sq ft)		
2 - 4 months	2	1.5	2	6
4 - 8 months	4	3	4	12
8 - 12 months	6	4.5	8	12

Industrial Sites - Tank Mix Combinations

Tank mix combinations of Surflan AS plus glyphosate, Spike* herbicide, and many other labeled herbicides may be used as overtop sprays to control existing vegetation on industrial sites such as utility substations, highway guard rails, sign posts, and delineators. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitation before use.

Warm Season Turfgrasses

Surflan AS may be applied as a preemergence treatment for control of annual grasses and certain broadleaf weeds in established warm season turf including bahiagrass, bermudagrass, buffalograss, centipedegrass, St. Augustinegrass, zoysiagrass, and established tall fescue growing in warm season areas. Established turf is defined as a dense turf having a well-anchored root system and healthy, vigorous top growth. Use Surflan AS only as a part of a total turf management program that includes good fertilization practices.

Surflan AS may be tank mixed with Gallery* herbicide (California registration pending) and applied preemergence to broaden the spectrum of broadleaf weed control in warm season turf. Refer to the label for Gallery for specific use directions, precautions, and limitations before use.

Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Surflan AS.

Surflan AS will not control emerged weeds. Successful preemergence control of weeds listed on this label requires that Surflan AS be applied prior to weed germination and be activated by at least one-half (1/2) inch of rainfall or irrigation within 21 days of application.

Surflan AS may injure turf that is not well established or is stressed or weakened due to unfavorable winter climatic conditions, drought, nematodes, or other factors which damage or weaken turf root systems. Apply Surflan AS only to healthy, well-established turf that has a well-anchored root system.

Do not apply Surflan AS in the spring or early summer to tall fescue turfgrass reseeded the previous fall. In such cases, apply Balan* 2.5G granular herbicide at 60-80 pounds per acre in early summer (Round 1) and Surflan AS at 1.5 quarts per acre approximately eight weeks later (Round 2). Do not apply Surflan AS

at the single application rate (2 quarts per acre) to established tall fescue; in such cases, apply 1.5 quarts per acre of Surflan AS in an initial application, followed by a second application of 1.5 quarts per acre 8-10 weeks later.

In bermudagrass areas that have been overseeded with winter grasses, a spring application of Surflan AS will thin the overseeded grasses.

Annual Grasses Controlled by Surflan AS

Summer Annuals:

Common Name	Scientific Name
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
crabgrass, large	<i>Digitaria sanguinalis</i>
crabgrass, smooth	<i>Digitaria ischaemum</i>
crabgrass	<i>Digitaria</i> spp.
crowfootgrass	<i>Dactyloctenium aegyptium</i>
foxtail, bristlegrass	<i>Setaria magna</i>
foxtail, giant	<i>Setaria faberi</i>
foxtail, green (pigeongrass)	<i>Setaria viridis</i>
foxtail, robust	<i>Setaria robusta</i>
foxtail, yellow	<i>Setaria glauca</i>
goosegrass (silver crabgrass)	<i>Eleusine indica</i>
Johnsongrass (seedling only)	<i>Sorghum halepense</i>
ryegrass, Italian	<i>Lolium multiflorum</i>
sandbur, field	<i>Cenchrus incertus</i>

Winter Annuals:

Common Name	Scientific Name
bluegrass, annual	<i>Poa annua</i>

Annual Broadleaf Weeds Controlled by Surflan AS

Summer Annuals:

Common Name	Scientific Name
carpetweed	<i>Mollugo verticillata</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
purslane, common	<i>Portulaca oleracea</i>

Winter Annuals:

Common Name	Scientific Name
chickweed, common	<i>Stellaria media</i>
henbit	<i>Lamium amplexicaule</i>

Broadleaf Weeds Suppressed by Surflan AS

Common Name	Scientific Name
groundsel, common	<i>Senecio vulgaris</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Application Rates, Frequency, and Timing of Application

Surflan AS can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for annual bluegrass (*Poa annua*) and winter annual broadleaf weed control.

25
26

Broadcast Application Rates (Warm Season Turfgrasses)

Use Area	Surflan AS		Minimum Time Between Applications	Total Amount Allowed Per Year
	(qt/acre)	(fl oz/1000 sq ft)	(months)	(qt/acre)
All, except Florida	1.5	1	3	6
Florida	2	1.5	3	6
Florida	1.5	1	3	4.5

1. Summer Annual Grasses and Broadleaf Weeds

Single Application Program: Apply 2 quarts per acre of Surflan AS in late winter or early spring, prior to the onset of conditions favorable for annual weed germination.

Split Application Program: As an alternative to a single application program, Surflan AS may be applied in a split application. This program is desirable when the initial application is made well in advance of weed germination and where weed control is desired for a longer period of time. Apply 1.5 quarts per acre of Surflan AS in an initial application, followed by a second application of 1.5 quarts per acre 8-10 weeks later.

The second treatment of the split application may follow application of a different preemergence grass herbicide in place of the initial application of Surflan AS.

2. Annual Bluegrass (*Poa annua*) and Winter Annual Broadleaf Weeds

In areas of heavy annual bluegrass infestation, its elimination will result in temporary thinning of turfgrass cover. Proper fertilization, irrigation, and soil incorporated reseeding should be employed to speed the restoration of desirable turfgrass cover in areas previously occupied by annual bluegrass (see section on reseeding).

Apply Surflan AS as a preemergence treatment in late summer or early fall, prior to the expected germination period for annual bluegrass and winter annual broadleaf weeds. If annual bluegrass infestation is severe and its elimination will result in thinning of turfgrass cover, apply Surflan AS at 1.5 quarts per acre. If thinning of turfgrass cover is not a potential problem, Surflan AS may be applied at 2 quarts per acre.

Weed Control in Florida

In Florida, apply 1.5 quarts per acre of Surflan AS three times per year, or every 90-100 days, in the fall, early spring, and early summer. Do not apply more than 1.5 quarts per acre of Surflan AS in any single application.

Application Equipment

Apply Surflan AS evenly over the turfgrass area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over-application. For best results, use application equipment designed to uniformly broadcast liquid herbicides. Calibrate application equipment prior to use, according to manufacturer's directions. Check equipment frequently to make sure it is working properly and distributing spray uniformly.

Reseeding

Herbicides that control annual weeds may also affect establishment of desirable turfgrass seedlings. Reseeding should be delayed for at least 90-120 days following application of Surflan AS. When reseeding, it is essential that proper cultural practices such as soil cultivation and seedbed preparation, irrigation, and fertilization be followed. For satisfactory reseeding results following use of Surflan AS, the seeding rate should be increased and equipment designed to place seed in full contact with soil (such as the Rogers Aero Seeder) should be employed.

Special Use Precautions:

To avoid possible injury, do not apply Surflan AS to:

- Cool season turfgrass species.

26
26

- Golf course putting greens and tees or lawns containing dichondra or cool season turfgrass species.
- Newly sprigged or sodded areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfgrasses are well established and have well-anchored root systems.
- Newly hydromulched areas of bermudagrass until such areas are well established.
- Bermudagrass variety "Sun Turf" when tank mixed with atrazine.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. **UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Surflan is a Registered Trademark of United Phosphorus, Inc.
PROKoZ is a Registered Trademark of PROKoZ, Inc.

EPA approved 12/23/05 .

Rev. 2/067/09