70506-46

००।यां २००७

	l		1	
0		I	٠	-

Please read instructions on revers	e before completing form	•	<u>Fo</u>	rm A				proval expires 5-31-98	
United States					Registra	ion	0	OPP Identifier Number	
≎EPA	Environmental		gency		Amendm	ent	27	74943	
	Washing	ton, DC 20460		X	Other				
	App	lication for Pe	esticide - Sectio	n I			'		
1. Company/Product Number			2. EPA Product M	_	jer	3.1	roposed C	Classification	
70506-46			Miller			H ,	٦	[] Dankistad	
4. Company/Product (Name) United Phosphorus, Inc/Surfla	n Dry Flowable		PM # 23				None	Restricted	
Name and Address of Appl	icant (Include ZIP Code)		6. Expedited Rev	iew	In accorda	ice wi	th FIFRA S	ection 3(c)(3)	
United Phosphorus, Inc.			(b)(i), my product is	s sim	ilar or identi	cal in	compositio	n and labeling	
423 Riverview Plaza Trenton, NJ 08611			to: EPA Reg No.			}			
				_					
Check if this is a new ad	dress		Product Name						
		Conti	ion - II						
The second secon		Secti		d lab	ele in roeno	ca to			
Amendment – Explain be	#IOW				els in respor	36 10			
Pasub mission in respons	no to Agonov lotter dated		Agency lette			NOTIFICATION			
Kesubmission in respons	se to Agency letter dated		Me 100 A	ppiic	allon		ECA	1 = 0000	
X Notification – Explain bel	ow		Other – Exp	dain	helow		FEB	1 5 2006	
Notification – Explain bei	OW		Outer = 2xp	, iaii i	DCION				
Explanation: Use additional p	page(s) if necessary (Fo	Section Land Sec	ction II)		-				
Change in name and mixing in	structions for tank mix pa	rtner already on la	ıbel.						
This notification is consistent v									
to the labeling or the confident false statement to EPA. I furth	ial statement of formula o	t this product. Tur	iderstand that it is a v	/iolat	ION Of 18 U.: of PR Notice	9.C. S	ec. 1001 to	Willfully make any	
product may be in violation of									
			<u> </u>						
		Sect	ion III			<u> </u>			
Material This Product Will be Child Decided to the Control of		Liac	tar Calubia Basinaia			1.	Tuna at C	-1-1-1	
Child-Resistant Packaging Yes	Unit Packaging Yes	🗸	ater Soluble Packagir Yes	ıg		2. Type of Container Metal			
		[]	No				Plastic		
No No	No					╝	Glass		
*Certification must	If "Yes"		Yes"		lo. per		Paper) No. 1 (18)	
be submitted	Unit Packaging wgt	container Pa	ckage wgt	۱ ۵	ontainer		Other (S	Specify)	
3. Location of Net Contents In	formation	4. Size(s) Retai	l Container		5. Lo	cation	of label di	rections	
Label Container					I	On La		i	
		1 1 1 1 1 1 1	·			On La	bel accom	panying product	
6. Manner in Which Label is A	mixed to Product	Lithograph Paper glue			Other	┼			
		Stenciled	·u						
Section IV									
1. Contact Person (Complete	items directly below for id	entification of indi	vidual to be contacted	i, if n	ecessary, to	proc	ess this app	olication.)	
							e Area Code)		
Rebecca A. Clemmer Regulatory Affairs Manager 609-39 Certification					92-82		e Application		
I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete.							ceived		
I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or									
both under applicable law 2. Signature 3 Title				+		(Stamped)			
			Affairs Manager					(-:milham)	
· illicitali.	being "		··-·- 3*·						
4 Typed Name 5 Date					 		7	•	
Rebecca A. Clemmer		January 16	, 2006					ļ	

EPA Form 8570-1 (Rev. 8-94) Previous editions are obsolete.

White - EPA File Copy (Original) Yellow - Applicant Copy

Surflan® Dry Flowable

A preemergence surface-applied herbicide for the control of most annual grasses and certain broadleaf weeds

Active Ingredient:

oryzalin: 3,5-dinitro-

N ⁴ , N ⁴ -dipropylsulfa	anilamide	85.0%
Inert Ingredients		
Total		100 0%

Contains 0.85 pounds of active ingredient per pound of product.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Avoid contact with eyes, skin or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- · Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.



User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.

 As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor immediately for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use, including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 70506-46

United Phosphorus, Inc.
423 Riverview Plaza
Trenton, NJ 08611
1-800-247-1557 • www.upi-usa.com

Herbicide-

Net Contents Ib

[Insert DOT shipping classification and diamond(s)]

Lot

418

(Datapack cover):

Surflan® Dry Flowable

A preemergence surface-applied herbicide for the control of most annual grasses and certain broadleaf weeds

Active Ingredient:

oryzalin: 3,5-dinitro-

 N⁴, N⁴-dipropylsulfanilamide
 85.0%

 Inert Ingredients
 15.0%

 Total
 100.0%

Contains 0.85 pounds of active ingredient per pound of product.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. Before buying or using this product, read "Warranty Disclaimer" and "Limitation of Remedies" inside label booklet.

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 70506-46

EPA Est.

United Phosphorus, Inc.
423 Riverview Plaza →
Trenton, NJ 08611 →
1-800-247-1557 • www.upi-usa.com

Herbicide*

Net Contents

lb

5/15

(Page 1 through end):

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes Moderate Eye Irritation • Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Avoid contact with eyes, skin or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- · Long-sleeved shirt and long pants
- · Chemical-resistant gloves
- · Shoes plus socks

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
 As soon as possible, wash thoroughly and change into clean clothing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor immediately for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.



Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. **Exception:** If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

Workers may enter treated areas without required PPE during the reentry interval following 1/2 to 1 inch of rainfall or irrigation, otherwise, PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- · Chemical-resistant gloves
- · Shoes plus socks

Storage and Disposal

Do not contaminate water, food or feed by storage or disposal.

Storage: Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Surflan Dry-Flowable herbicide is a preemergence, surface applied herbicide for the control of most annual grasses and certain broadleaf weeds. Surflan Dry Flowable controls susceptible annual weeds by disrupting plant growth processes during germination. Surflan Dry Flowable may be applied in liquid sprays of water or liquid fertilizer. Surflan Dry Flowable may be tank mixed with other herbicides to control existing vegetation or improve the spectrum of weeds controlled. Surflan Dry Flowable atone does not control established weeds.

General Use Precautions and Restrictions

Do not graze or feed forage from treated fields or orchards to livestock.

Do not over-apply Surflan Dry Flowable. Over-application may result in crop injury or soil residue.

Do not plant any root crop for 12 months following a Surflan Dry Flowable application.

Do not use Surflan Dry Flowable on soils containing more than 5% organic matter.

Page 6

7115

Apply Surflan Dry Flowable directly to the soil surface in orchards or vineyards. Carefully follow label directions to avoid poor weed control or crop injury.

Chemigation: Do not apply this product through any type of irrigation system.

Do not aerially apply this product.

Soil Preparation

Optimum herbicidal activity occurs when Surflan Dry Flowable is applied to the soil surface. Prior to application, existing unwanted vegetation should be destroyed by tillage or use of contact or translocated herbicides. Crop or weed residues, pruning and trash should be thoroughly mixed into the soil by tillage equipment or removed before treatment. Soil should be in good tilth and free of clods at the time of application.

Activation and Cultivation

A single 1/2 to 1-inch rainfall or sprinkler irrigation is required to activate Surflan Dry Flowable and move the herbicide into the zone of weed germination. Rainfall or irrigation of 1 inch or more is needed to activate Surflan Dry Flowable on fine-textured, high organic matter soils. If weeds begin to emerge, a shallow cultivation to a depth of 1 to 2 inches will destroy existing weeds and place Surflan Dry Flowable in the zone of weed germination.

Mixing Directions

Surflan Dry Flowable - Alone

Surflan Dry Flowable may be applied in water or most liquid fertilizer materials. Prior to mixing Surflan Dry Flowable in liquid fertilizer, refer to "Testing for Compatibility in Liquid Fertilizers" for test procedures to determine compatibility with the fertilizer product to be used. The combination of Surflan Dry Flowable with solution and suspension-type fertilizers provides annual weed control equal to Surflan Dry Flowable applied in water. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale.

Start with a clean spray tank. Fill the sprayer to 1/3 to 1/2 of the required spray volume. Start agitation. Add the correct amount of Surflan Dry Flowable, continue agitation, and fill spray tank to required spray volume. Maintain continuous agitation from mixing through application.

Precaution: Do not allow the mixture to siphon back into the water source.

Surflan Dry Flowable - Tank Mix Combinations

Surflan Dry Flowable may be tank mixed with other products and applied with water or most liquid fertilizer materials. Prior to mixing Surflan and another product in liquid fertilizer, refer to the tank mix product manufacturer's label to determine if application in liquid fertilizer is recommended. Also refer to the section in this label entitled "Testing for Compatibility in Liquid Fertilizers" for testing procedures to determine tank-mix compatibility with the liquid fertilizer product to be used. The combination of Surflan with solution and suspension-type fertilizers provides annual weed control equal to \$urflan applied in water. Individual state regulations relating to liquid fertilizer mixing, registration, labeling and application are the responsibility of the individual and/or company offering the fertilizer and chemical mixture for sale. Read and follow all label instructions for each material to be added to the spray tank.

Vigorous and continuous agitation is required for all tank mixes. Sparger pipe agitators generally provide the best agitation in spray tanks. To prevent foaming, keep the end of the fill pipe below the surface of the water in the spray tank during filling to prevent air from being stirred or splashed into the mixture.



Mixing Order: Fill the spray tank to 3/4 of the total spray volume required. Start agitation. Add different formulation types in the order indicated below, allowing time for complete mixing and dispersion after addition of each product. Allow extra mixing and dispersion time for dry flowable products.

Add different formulation types in the following order: Surflan Dry Flowable and other dry flowables (DF); wettable powders (WP); flowables (F); liquids (L) or aqueous suspensions (S); solutions (S); and emulsifiable concentrates (EC).

Fill the spray tank to final volume. Maintain continuous agitation during mixing, final filling and throughout application. If spraying and agitation must be stopped before the spray tank is empty, the materials may settle to the bottom. Settled materials must be suspended before spraying is resumed. A sparger agitator is particularly useful for this purpose. Settled materials may be more difficult to resuspend than when originally mixed.

Premixing: When tank mixing, initial mixing and dispersion of Surflan Dry Flowable or other dry flowable or wettable powder products may be improved by premixing with water (slurrying). Where recommended, follow product label instructions for each material. Adding the slurried material to the spray tank through a 20 to 35 mesh-wetting screen will help assure good initial dispersion. Line screens in the tank through a no finer than 50 mesh (100 mesh is finer than 50 mesh).

Testing for Compatibility in Liquid Fertilizers

Surflan Dry Flowable alone or in combination with other product formulations may not combine properly with some liquid fertilizer materials. Small quantities of such mixtures should always be tested before full-scale mixing. Follow the testing procedure below to determine if a compatibility agent is needed or which compatibility agent performs best in the mixture being applied. Nine compatibility agents listed at the end of this section have been thoroughly tested. Other surfactant materials available may not be formulated for use with liquid fertilizers.

Testing Procedure

- 1. Add 1 pint of liquid fertilizer to a 1-quart glass jar.
- 2. Add 1 to 4 teaspoonfuls (depending on application rate) of the products to be tank mixed in the liquid fertilizer. Add the materials in the order recommended in the "Mixing Directions" section of this label. Mix until evenly dispersed after addition of each formulation. If dry flowable or wettable powder formulations do not disperse well, it may be necessary to slurry the materials in a small amount of water before addition to the liquid fertilizer.
- 3. Observe the jar for about 10 minutes. If materials rise to the surface and form a thick layer (oil curds) which will not redisperse when agitated, a compatibility agent is needed. If the mixture is easily redispersed with slight agitation, a compatibility agent is not required, however, good agitation must be provided to maintain dispersion in the spray tank from mixing through application.
- 4. If the need for a compatibility agent is demonstrated in step 3, it may be tested as follows: Using a clean clear plastic or glass container, start at step 1 above and add 1/2 teaspoon of the compatibility agent to the liquid fertilizer mixture. Shake well and then repeat steps 2 and 3. An effective compatibility agent will cause the mixture to remain uniformly mixed with little or no separation for 1/2 hour or longer. If slight separation occurs, 2 to 3 inversions of the container should be sufficient to uniformly redisperse the mixture. If oily curds form which will not disperse, try adding additional compatibility agent or use an alternative compatibility agent to achieve a uniform mixture. A clean jar should be used for each test.

Compatibility Agents

The phosphate ester-type surfactants listed below are designed for use with liquid fertilizers and can be mixed at rates as low as 1 1/2 to 2 pints per ton of liquid fertilizer. They usually do not work well as compatibility agents for tank mixes in plain water. Add the compatibility agent just before adding herbicides. Read and follow label directions for the compatibility agent.

915

- 1. Sponto 168D (Witco Chemicals Co., Chicago, IL)
- 2. Compat (Farm Chemicals, Inc., Aberdeen, NC)
 Not registered for use in California.
- 3. Unite (Hopkins Ag Chemical, Madison, WI)
- 4. T-Mulz 734-2 (Thompson-Hayward Chemical Co., Kansas City, KS) Not registered for use in California.
- 5. Rigo Compatibility Agent (Rigo Company, Buckner, KY) Not registered for use in California.
- Amoco Spray Mate (Amoco Oil Co., Chicago, IL) Not registered for use in California.
- 7. Kem-Link (Universal Coop, Minneapolis, MN) Not registered for use in California.
- 8. Blendex (Helena Chemical, Cayce-West, Columbia, SC) Not registered for use in California.
- Spray-Aide (Miller Chemical and Fertilizer Corp., Hanover, PA)

Application Methods

Ground Broadcast Application

Apply Surflan Dry Flowable in a total spray volume of 20 to 40 gallons per acre (broadcast basis), using any properly calibrated low-pressure herbicide sprayer that will apply the spray uniformly. Use herbicide nozzle tips and screens no finer than 50 mesh for nozzle and in-line strainers. As the amount of spray volume per acre decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to insure proper calibration and uniform application. Avoid boom overlaps that will increase rates above those recommended.

Band Application

For band application, use the following formula to calculate the required amount of product per acre.

Band Width (inches)

X Broadcast rate per acre = Amount required per acre

Row Width (inches)

Weeds and Grasses Controlled in Fruit and Nut Crops and Vineyards

Annual Grasses

Common Name
barley, little
barnyardgrass (watergrass)
bluegrass, annual (poa)
brachiaria (signalgrass)
browntop panicum

Scientific Name
Hordeum pusillum
Echinochloa crus-galli
Poa annua
Brachiaria spp.
Panicum fasciculatum

Digitaria spp.

crabgrass (large crabgrass) .

(smooth crabgrass)

ootgrass

Dactyloctenium
aegyptium

cupgrass Eriochloa gracilis
downy brome Bromus tectorum
fall panicum Panicum dichotomiflorum

(spreading panicgrass)

foxtails Setaria spp.

(bottlegrass) (bristlegrass)



(giant foxtail) (green foxtail) (pigeongrass) (robust foxtail) (yellow foxtail) quineagrass

(narrowleaf panicum)

johnsongrass (seedling

only)

junglerice lovegrass, Mexican lovegrass, orcutt

oat, wild ryegrass, annual (Italian) sandbur, field sprangletop, red Texas panicum (buffalograss) (Coloradograss)

witchgrass

Panicum maximum

Sorghum halepense

Echinochioa colonum Eragrostis mexicana Eragrostis orcuttiana Avena fatua

Lolium multiflorum Cenchrus incertus Leptochloa filiformis Panicum texanum

Panicum capillare

Annual Broadleaf Weeds

Common Name Scientific Name carpet weed Mollugo verticillata chickweed Stellaria media cudweed Gnaphalium chilense fiddleneck, coast Amsinckia intermedia Florida pustey Richardia scabra (Florida purslane)

(Mexican clover)

(pusiey)

henbit Lamium amplexicaule knotweed, prostrate Polygonum aviculare lambsquarters Chenopodium album pigweeds Amaranthus spp.

(carelessweed) (prostrate pigweed) (redroot pigweed) (rough pigweed) (smooth pigweed) (spiny pigweed) (tumble pigweed)

Puncturevine Tribulus terrestris purslane, common Portulaca oleracea rockpurslane, redmaids Calandrinia caulescens shepherdspurse Capsella bursa-pastoris

Surflan Dry Flowable provides partial control or suppression of:

Common Name Scientific Name filaree, redstem Erodium cicutarium filaree, whitestem Erodium moschatum groundsel, common Senecio vulgaris ladysthumb Polygonum persicaria lettuce, prickly Lactuca serriola

mallow, common Malva neglecta

Sarcostemma cynanchoides milkweed, climbing

Ipomoea spp. morningglory, annual Brassica nigra mustard, black Sinapis arvensis mustard, wild Solanum nigrum nightshade, black prickly sida (teaweed) Sida spinosa

ragweed, common Ambrosia artemisiifolia

Ambrosia trifida ragweed, giant Sisymbrium irio rocket, London smartweed, annual Polygonum spp. sowthistle, annual Sonchus oleraceus spurge, prostrate Euphorbia humistrata spurge, spotted Euphorbia maculata velvetleaf Abutilon theophrasti

Approved Crops

Fruit and Nut Crops and Vineyards (Non-bearing and Bearing)

Surflan Dry Flowable - Alone

Apply Surflan Dry Flowable to fruit and nut crops and vineyards as a preemergence treatment to control annual grasses and broadleaf weeds listed in "General Information". Surflan Dry Flowable is approved for use on the following crops:

almond grapefruit pear apple kiwi pecan apricot lemon pistachio avocado 1 macadamia nut plum pomegranate

cherry nectarine

fig olive prune

filbert walnut, English orange

grape peach

Small Fruits:

blackberry currant gooseberry blueberry† dewberry loganberry boysen**berry** elderberry raspberry

†Do not apply Surflan Dry Flowable to lowbush blueberries.

General Use Directions

Existing vegetation should be destroyed by tillage or use of a contact or translocated herbicide in tank mix combination with Surflan Dry Flowable. To insure tank mix compatibility of Surflan Dry Flowable with contact or translocated herbicides, mix the products together in correct proportions in a small container of water. After mixing, let stand for 5 minutes and check for compatibility. If the combination remains mixed or can be resuspended easily, the mixture is compatible.

Application Methods: Surflan Dry Flowable may be applied in liquid sprays of water or liquid fertilizer. Refer to the "General Information" section of this label for mixing directions and details on application methods and precautions before use.

Activation: A single 1/2 to 1 inch rainfall or sprinkler irrigation is required to activate Surflan Dry The value and move the herbicide into the zone of weed seed germination. If weeds begin to emerge, a



shallow cultivation 1 to 2 inches deep will destroy existing weeds and place Surflan Dry Flowable in the zone of weed seed germination. Rainfall or sprinkler irrigation of 1 inch or more is required to activate Surflan Dry Flowable on fine-textured or high organic matter soil.

Broadcast Application Rates

Soil Texture	Length of Control	Surflan Dry Flowable (lb/acre)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (lb/acre)
All Soil	Short Term (2 - 4 months)	2.4	2.5	14
Textures	Long Term (6 - 8 months)	4.7	2.5	14
	(8 - 12 months)	7.1	2.5	14

Precautions:

Poor weed control may result if directions are not carefully followed. Over-application may result in crop injury or excessive soil residue. Apply spray directly to the soil surface of the orchard or vineyard.

Surflan Dry Flowable Tank Mix Recommendations

The following products are compatible with Surflan Dry Flowable when used in tank mix combination. Performance and risk of carryover from products listed below when used in combination with Surflan Dry Flowable at recommended rates is the same as when each product is used separately.

Follow tank mixing instructions provided in the "General Information" section of this label when mixing Surflan Dry Flowable with other products.

Users should always consult the manufacturer's label for the product(s) to be tank mixed with Surflan Dry Flowable for specific information on use rates, additional weeds controlled, rotational crop restrictions or risk of carryover, special tank mix instructions, additional use directions, precautions and limitations before use.

Note: Some crop uses of tank mix products are limited to or excluded from certain states.

Recommended Surflan Dry Flowable Tank Mix Combinations:

Surflan Dry Flowable plus Goai

Surflan Dry Flowable plus Goal plus Gramoxone Max*

Surfian Dry Flowable plus Gramoxone Max*

Surflan Dry Flowable plus Princep (simazine)

Surflan Dry Flowable plus glyphosate

Surflan Dry Flowable plus Solicam

Surflan Dry Flowable - Tank Mix Reference Table:

The following table lists approved crops for Surflan Dry Flowable and products for tank mixes with Surflan Dry Flowable

(** = Approved Crop). Refer to the table to determine if the herbicides recommended for tank mixing can be applied to your crop. For use on a given crop, two asterisks must be indicated for Surflan Dry Flowable and the product with which it is to be tank mixed.

^{*} Tank mixtures of Surflan Dry Flowable and Gramoxone Max may be incompatible. To improve the quality of the spray mixture, fill the spray tank with water to % of desired final volume, start agitation, add Surflan Dry Flowable to the spray tank, add an ammonium sulfate water conditioner/compatibility agent to the spray mixture then add Gramoxone Max to the spray tank. Once all ingredients have been added to the spray tank, finish filling the spray tank with water to the final desired volume.



	Surflan Dry Flowable	Goal	Gramoxone Max	Princep (simazine)	glyphosate	Solica m
Almond	**	**	44	**	**	**
Apple	**	44	**	**	#0	**
Apricot	**	**	# b	-	**	*
Avocado	8-2	**	re .	**	89	-
Cherry	**	**	**	**	**	-
Fig	**	**	**	-	**	-
Filbert	**	**	**	**	*1	**
Grape	6-5	**	**	44	**	pá
Grapefruit	**	-	**	**	**	**
Kiwi	**	**	**	-	**	-
Lemon	**	•	**	**	**	**
Macadamia nut	**	**	**	**	**	-
Nectarine	**	••	**	-	**	**
Olive	••	**	**	**	**	-
Orange	**	-	**	**	**	**
Peach	••	**	**	**	**	**
Pear	**	**	**	**	**	**
Pecan	**	**	**	**	**	**
Pistachio	**	**	••	-	**	•
Plum	89	**	**	ė e	**	**
Pomegranate	••	**	-	-	**	-
Prune	**	••	**	-	**	**
Walnut,	**	**	••	-	**	**
English						
Small Fruits:						, <u>-</u>
Blackberry	**	-	**	**	**	**
Blueberry	**	-	**	**	**	**
Boysenberry	**	-	**	**	**	
Currants	••	•		-	**	•
Dewberry	**	-	-	•	•	-
Elderberry	**		•	_	••	
Gooseberry	#4	-			**	-
Loganberry	**	-	-	••	**	-
Raspberry	**	-	•	**	**	••

Warranty Disclaimer

United Phosphorus, Inc. (UPI) warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. UPI MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Inherent Risks of Use

It is impossible to eliminate all risks associated with use of this product. Crop injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperatures, soil conditions, etc.), abnormal conditions (such as excessive rainfall drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors all of which are beyond the control of UPI or the seller. All such risks shall be assumed by buyer.



Limitation of Remedies

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at UPI's election, one of the following:

- 1. Refund of purchase price paid by buyer or user for product bought, or
- 2. Replacement of amount of product used.

UPI shall not be liable for losses or damages resulting from handling or use of this product unless UPI is promptly notified of such loss or damage in writing. In no case shall UPI be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of UPI or the seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

Goal is a trademark of Dow AgroSciences, LLC Gramoxone, Princep, Solicam are trademarks of Syngenta Crop Protection

EPA-accepted: 11/30/1999

Rev.1/16/06



United Phosphorus, Inc.

423 Riverview Plaza Trenton, NJ 08611(609) 392-8200 x 1018 (phone)
(609) 392-0808 (fax)

Rebecca A. Clemmer Regulatory Affairs Manager

January 16, 2006

Joanne Miller (PM 23)
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504C)
U.S. Environmental Protection Agency
1801 Bell Street
Arlington, VA 22202

Re:

Surflan® Dry Flowable (EPA Reg. No. 70506-46)

Notification of minor label change

Dear Ms. Miller:

Enclosed please find notification of an amendment to the subject label. We have altered the name of one of the tank mix partners in order to match a change in that product's name (Gramoxone Super is changed to Gramoxone Max), and we have added additional instructions for when Surflan DF is tank mixed with that product. In support of this action, enclosed please find:

- EPA Form 8570-1:
- One copy of the label, marked to show changes;
- One 'clean' copy of the label; and
- One postcard to return to indicate acceptance of this Notification.

Please contact me if you have any questions.

Victorial Continuer

Very truly yours,

Rebecca A. Clemmer

reiemmer/aupi-usa.com