

70506-47

12/28/2005

1/16

(Container label)

Surflan® 0.25 A.S.

A Selective Preemergence Surface-Applied Herbicide for Use in Established Ornamentals and Southern Turfgrasses for the Control of Annual Grasses and Many Broadleaf Weeds in:

- Container Grown Ornamental Plants
- Ground Covers
- Noncropland Areas
- Ornamental Trees
- Ornamental Plants (Including Around Highway and Industrial Sites)
- Ornamental Woody Shrubs
- Southern Turfgrasses (Including Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, Tall Fescue and Zoysiagrass)

Active Ingredient:

oryzalin: 3,5-dinitro-N ⁴ ,N ⁴ -dipropylsulfanilamide	3%
Inert Ingredients	97%
Total	100%

Contains 0.25 pounds active ingredient per gallon.

Avoid Freezing--Store Above 40°F

Keep Out of Reach of Children

CAUTION

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • May Cause Skin Allergies To Develop

Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- In addition, mixers and loaders must also wear a chemical-resistant apron.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ACCEPTED

DEC 28 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

70506-47

2/16

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment wash waters or rinsate.

See label booklet for Directions for Use, Including Storage and Disposal.

Notice: Read entire label. Use only according to label directions. **Before buying this product, read Conditions of Sale and Limitation of Warranty and Liability elsewhere on this label.**

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

(Logo) United Phosphorus, Inc.
United Phosphorus, Inc.
423 Riverview Plaza
Trenton, NJ 08611
1-800-247-1557 • www.upi-usa.com

EPA Reg. No. 70506-47

EPA Est. _____

Herbicide

Net Contents - ____

3/16

(Label Booklet)

Surflan® 0.25 A.S.

A Selective Preemergence Surface-Applied Herbicide for Use in Established Ornamentals and Southern Turfgrasses for the Control of Annual Grasses and Many Broadleaf Weeds in:

- Container Grown Ornamental Plants
- Ground Covers
- Noncropland Areas
- Ornamental Trees
- Ornamental Plants (Including Around Highway and Industrial Sites)
- Ornamental Woody Shrubs
- Southern Turfgrasses (Including Bahiagrass, Bermudagrass, Centipedegrass, St. Augustinegrass, Tall Fescue and Zoysiagrass)

Active Ingredient:

oryzalin: 3,5-dinitro- <i>N</i> ⁴ , <i>N</i> ⁴ -dipropylsulfanilamide	3%
Inert Ingredients	97%
Total	100%

Contains 0.25 pounds active ingredient per gallon.

Avoid Freezing--Store Above 40°F

Keep Out of Reach of Children

CAUTION

See label booklet for Directions for Use including Storage and Disposal.

Notice: Read entire label. Use only according to label directions. **Before buying this product, read Limited Warranty and Disclaimer Section elsewhere on this label.**

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Do Not Ship or Store with Food, Feeds, Drugs, or Clothing.

(Logo) United Phosphorus, Inc.
 United Phosphorus, Inc.
 423 Riverview Plaza
 Trenton, NJ 08611
 1-800-247-1557 • www.upi-usa.com

EPA Reg. No. 70506-47

EPA Est. _____

Herbicide

Net Contents - ____

4/16

Precautionary Statements

Hazards to Humans and Domestic Animals

Keep Out of Reach of Children

CAUTION

Causes Moderate Eye Irritation • May Cause Skin Allergies To Develop

Avoid contact with skin, eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- In addition, mixers and loaders must also wear a chemical-resistant apron.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside.
- Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water when disposing of equipment wash waters or rinsate.

5/16

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all directions for use carefully before applying.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Avoid freezing. Store above 40°F. Store in original container only. In case of leak or spill, use absorbent materials to contain liquids and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

General Information

Surflan 0.25 A.S. is a preemergence surface-applied herbicide for the control of annual grasses and many broadleaf weeds.

Soil Preparation: Surflan 0.25 A.S. will not control established weeds. Therefore, areas to be treated should be free of established weeds. Weed residues, prunings and trash should be thoroughly mixed into the soil or removed prior to treatment. The soil should be in good tilth and free of clods at the time of application.

Application Restrictions:

- Keep all persons, children, and pets out of treated area until sprays have dried.
- Do not graze or feed forage from treated areas to livestock.
- Do not aerially apply this product.
- **Chemigation:** Do not apply this product through any type of irrigation system.

Ornamentals

General Use Precautions and Restrictions for Ornamentals

Apply only to established plantings. Rooted liners should be established in containers two weeks prior to treatment or crop injury may occur.

6/16

Do not apply Surflan 0.25 A.S. in ornamental plantings where the likelihood of runoff onto lawn areas containing dichondra or cool-season turfgrass species exists as severe injury or death may occur. Poor weed control may result if directions are not followed. Over application may result in crop injury or excessive soil residue.

Plant only ornamental species listed on this label into soil treated the previous season with Surflan 0.25 A.S. or injury may occur.

Grasses Controlled by Surflan 0.25 A.S.

Common Name	Scientific Name
barley, little	<i>Hordeum pusillum</i>
barnyardgrass (Watergrass)	<i>Echinochloa crusgalli</i>
bluegrass, annual	<i>Poa annua</i>
brachiaria (signalgrass)	<i>Brachiaria</i> spp.
crabgrasses (large crabgrass) (smooth crabgrass)	<i>Digitaria</i> spp.
crowfootgrass	<i>Dactyloctenium aegyptium</i>
cupgrass, southwestern	<i>Eriochloa gracilis</i>
foxtails	<i>Setaria</i> spp.
(bottlegrass)	
(bristlegrass)	
(giant foxtail)	
(green foxtail)	
(pigeongrass)	
(robust foxtail)	
(yellow foxtail)	
goosegrass (Silver crabgrass)	<i>Eleusine indica</i>
johnsongrass (Seedling only)	<i>Sorghum halepense</i>
junglerice	<i>Echinochloa colonum</i>
lovegrass, Mexican	<i>Eragrostis mexicana</i>
lovegrass, orcutt	<i>Eragrostis orcuttiana</i>
oat, wild	<i>Avena fatua</i>
panicum, browntop	<i>Panicum fasciculatum</i>
panicum, fall (Spreading panicgrass)	<i>Panicum dichotomiflorum</i>
panicum, Texas	<i>Panicum texanum</i>
buffalograss	
coloradograss	
ryegrass, Italian	<i>Lolium multiflorum</i>
sandbur, field	<i>Cenchrus incertus</i>
sprangletop, red	<i>Leptochloa filiformis</i>
witchgrass	<i>Panicum capillare</i>

Broadleaf Weeds Controlled by Surflan 0.25 A.S.

Common Name	Scientific Name
bittercress	<i>Cardamine</i> spp.
carpetweed	<i>Mollugo verticillata</i>
chickweed, common	<i>Stellaria media</i>
fiddleneck, coast	<i>Amsinckia intermedia</i>
filaree, redstem	<i>Erodium cicutarium</i>
filaree, whitestem	<i>Erodium moschatum</i>
groundsel, common	<i>Senecio vulgaris</i>
henbit	<i>Lamium amplexicaule</i>
knotweed, prostrate	<i>Polygonum aviculare</i>

7/16

lambsquarters	<i>Chenopodium album</i>
pigweeds	<i>Amaranthus spp.</i>
(carelessweed)	
(prostrate pigweed)	
(redroot)	
(rough pigweed)	
(spring pigweed)	
(tumble pigweed)	
puncturevine	<i>Tribulus terrestris</i>
purslane, common	<i>Portulaca oleracea</i>
pusley, Florida	<i>Richardia scabra</i>
(Florida purslane)	
(Mexican clover)	
(pusley)	
rocket, London	<i>Sisymbrium irio</i>
rockpurslane, desert	<i>Calandrinia ciliata</i>
shepherdspurse	<i>Capsella bursa pastoris</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
woodsorrel, yellow	<i>Oxalis stricta</i>

Suppression of the following weeds may be erratic, ranging from poor to excellent, depending upon soil temperature, time of germination, depth of seed in the soil, amount and timing of soil moisture.

Common Name	Common Name
annual sowthistle	morningglory
black mustard	prickly lettuce
black nightshade	smartweed
climbing milkweed	spotted spurge
common mallow	teaweed (prickly sida)
common ragweed	velvetleaf
horseweed	volunteer wheat
ladysthumb	wild mustard

Surflan 0.25 A.S. will not control established weeds.

Mixing Directions

Start with a clean spray tank. Fill sprayer one-third ($1/3$) to one-half ($1/2$) full with clean water. Start agitation. Add correct quantity of Surflan 0.25 A.S., continue agitation and finish filling the tank. Maintain agitation from mixing through application.

If you see a buildup of material on the walls of the spray tank, wash the tank with soapy water between fillings. Rinse and continue the spraying operation. Clean the tank, lines, and screens thoroughly after use.

Application

Ground Application: Apply Surflan 0.25 A.S. as a directed spray to the soil surface or as an overtop spray. Apply Surflan 0.25 A.S. in sufficient water per acre to uniformly treat the area. Use only a properly calibrated, low-pressure herbicide sprayer that will apply the spray uniformly. Use herbicide tips and screens no finer than 50 mesh in nozzle and in-line strainers. As the amount of water used (spray volume) decreases, the importance of accurate calibration and uniform application increases. Check the sprayer daily to ensure proper calibration and uniform application. Maintain agitation from mixing through application. Avoid boom overlaps that will increase rates above those recommended.

9/16

Activation and Cultivation

Surflan 0.25 A.S. will remain stable on the soil surface up to 21 days. In the absence of timely rainfall, irrigation can be used to move Surflan 0.25 A.S. into soil surface. A one-half (1/2) inch rain or its equivalent in sprinkler irrigation is necessary to activate Surflan 0.25 A.S. If weeds begin to emerge due to lack of rainfall or irrigation, shallow cultivate (1 to 2 inches) to destroy existing weeds or remove by hand in containers and place Surflan 0.25 A.S. in the weed germination zone. Shallow cultivation (1 to 2 inches) will not reduce herbicidal effectiveness. If Surflan 0.25 A.S. is not activated by rainfall, irrigation or cultivation within 21 days of application, erratic weed control may result.

Recommended Species

Surflan 0.25 A.S. may be used on the Following Ornamental Plant Species:

Common Name	Scientific Name
abelia, glossy	<i>Abelia grandiflora</i>
apple	<i>Malus</i> spp.
arborvitae, Giant	<i>Thuja plicata</i>
arborvitae, Oriental	<i>Platycladus orientalis</i>
azalea	<i>Rhododendron</i> spp.
barberry, Japanese	<i>Berberis thunbergii</i>
begonia	<i>Begonia semper florans</i>
bird of paradise	<i>Strelitzia reginae</i>
bottlebrush, lemon	<i>Callistemon citrinus</i>
boxwood, common	<i>Buxus sempervirens</i>
boxwood, Japanese	<i>Buxus microphylla japonica</i>
brush cherry	<i>Syzygium paniculata</i>
California laurel	<i>Umbellularia californica</i>
cape marigold	<i>Dimorphotheca</i> spp.
cherry, Mahaleb	<i>Prunus mahaleb</i>
cherry, sweet	<i>Prunus avium</i>
chrysanthemum, florists	<i>Chrysanthemum morifolium</i>
cleyera, Japanese	<i>Cleyera japonica</i>
cotoneaster, bearberry	<i>Cotoneaster dammeri</i>
cotoneaster, brightbead	<i>Cotoneaster buxifolius</i>
cotoneaster, cranberry	<i>Cotoneaster apiculatus</i>
cotoneaster, Pyrenees	<i>Cotoneaster congestus</i>
cotoneaster, rock	<i>Cotoneaster horizontalis</i>
cotoneaster, rockspray	<i>Cotoneaster microphyllus</i>
coyotebush, dwarf	<i>Baccharis pilularis</i>
crape Myrtle, common	<i>Lagerstroemia indica</i>
cryptomeria, Japanese	<i>Cryptomeria japonica</i>
cypress, Italian	<i>Cupressus sempervirens</i>
daisy, trailing African	<i>Osteospermum fruticosum</i>
douglasfir	<i>Pseudotsuga taxifolia</i>
eastercactus	<i>Rhipsalidopsis gaertneri</i>
escallonia	<i>Escallonia exoniensis</i>
eucalyptus, mealy	<i>Eucalyptus cinerea</i>
eucalyptus, narrow-leaved	<i>Eucalyptus nicholii</i>
eucalyptus, red	<i>Eucalyptus sideroxylon</i>
euonymus, evergreen	<i>Euonymus japonica</i>
euonymus, stringybark	<i>Euonymus fortunei</i>
euonymus, winged	<i>Euonymus alata</i>
falsecypress, Lawson	<i>Chamaecyparis lawsoniana</i>
fatshedera	<i>Fatshedera lizei</i>

9/16

fir, alpine	<i>Abies lasiocarpa</i>
fir, balsam	<i>Abies balsamea</i>
fir, fraser	<i>Abies fraseri</i>
fir, grand	<i>Abies grandis</i>
fir, Vietch	<i>Abies veitchi</i>
fir, white	<i>Abies concolor</i>
firethorn	<i>Pyracantha, fortuneana</i>
firethorn, formosa	<i>Pyracantha skoidzumi</i>
firethorn, scarlet	<i>Pyracantha coccinea</i>
forsythia, border	<i>Forsythia intermedia</i>
gazania, trailing	<i>Gazania rigens leucolaena</i>
geranium (Pelargonium)	<i>Pelargonium hortorum</i>
ginkgo	<i>Ginkgo biloba</i>
hibiscus, Chinese	<i>Hibiscus rosa-sinesis</i>
holly, Chinese	<i>Ilex cornuta</i>
holly, English	<i>Ilex aquifolium</i>
holly, Japanese	<i>Ilex crenata</i>
honeysuckle, Japanese	<i>Lonicera japonica</i>
hopseedbush, clammy	<i>Dodonaea viscosa</i>
ice plant	<i>Mesembryanthemum crystallinum</i>
(See precautions for ornamental plantings)	
ice plant, largeleaf	<i>Carpobrotus edulis</i>
(See precautions for ornamental plantings)	
ivy, Algerian	<i>Hedera canariensis</i>
ivy, English	<i>Hedera helix</i>
jerseytea, redroot	<i>Ceanothus americanus</i>
juniper	<i>Juniperus spp.</i>
laurelcherry, Carolina	<i>Prunus caroliniana</i>
laurelcherry, English	<i>Prunus laurocerasus</i>
leucothoe, coast	<i>Leucothoe axillaris</i>
leucothoe, drooping	<i>Leucothoe fontanesiana</i>
lilac, common	<i>Syringa vulgaris</i>
lily, plantain	<i>Hosta spp.</i>
lilyturf, bigblue	<i>Liriope muscari</i>
lily-of-the-Nile	<i>Agapanthus africanus</i>
linden, little leaf	<i>Tilia cordata</i>
magnolia, Southern	<i>Magnolia grandiflora</i>
manzanita, Stanford	<i>Arctostaphylos stanfordiana</i>
maple	<i>Acer spp.</i>
marigold	<i>Tagetes spp.</i>
mockorange	<i>Philadelphus spp.</i>
moss, rose	<i>Portulaca grandiflora</i>
myoporum, prostrate	<i>Myoporum parvifolium</i>
myrtle, true	<i>Myrtus communis</i>
oak <i>Quercus</i> spp.	
oleander	<i>Nerium oleander</i>
orange, ornamental	<i>Citrus spp.</i>
oregon grape	<i>Mahonia aquifolium</i>
osmanthus, holly-leaf	<i>Osmanthus heterophyllus</i>
palo verde, blue	<i>Cercidium floridum</i>
pansy	<i>Viola wittrockiana</i>
pear	<i>Pyrus communis</i>
pecan, ornamental	<i>Carya spp.</i>
periwinkle, bigleaf	<i>Vinca major</i>

10/11

periwinkle, dwarf	<i>Vinca minor</i>
petunia	<i>Petunia</i> spp.
photinia	<i>Photinia fraseri</i>
pine	<i>Pinus</i> spp.
pittosporum	<i>Pittosporum</i> spp.
privet, amur	<i>Ligustrum amurense</i>
privet, glossy	<i>Ligustrum lucidum</i>
privet, golden	<i>Ligustrum vicaryi</i>
privet, Japanese	<i>Ligustrum japonicum</i>
protea	<i>Protea neriifolia</i>
ranunculus, Persian	<i>Ranunculus asiaticus</i>
redbud	<i>Cercis canadensis</i>
redcedar, eastern	<i>Juniperus virginiana</i>
redcedar, western	<i>Thuja plicata</i>
redwood, coast	<i>Sequoia sempervirens</i>
raphiolepis (India hawthorn)	<i>Raphiolepis indica</i>
rhododendron	<i>Rhododendron</i> spp.
rose	<i>Rosa</i> spp.
rose-of-Sharon (Shrubalthea)	<i>Hibiscus syriacus</i>
Russian olive	<i>Elaeagnus angustifolia</i>
sage	<i>Salvia</i> spp.
shrimp plant	<i>Justicia brandegeana</i>
snapdragon	<i>Antirrhinum majus</i>
sotol, desert spoon	<i>Dasyllirion wheeleri</i>
spruce, black	<i>Picea mariana</i>
spruce, Colorado	<i>Picea pungens</i>
spruce, Englemann	<i>Picea englemanni</i>
spruce, Norway	<i>Picea abies</i>
spruce, white	<i>Picea glauca</i>
star jasmine, Chinese	<i>Trachelospermum jasminoides</i>
stonecrop	<i>Sedum brevifolium</i>
sumac, African	<i>Rhus lancea</i>
sweetgum, American	<i>Liquidambar styraciflua</i>
sweet William	<i>Dianthus barbatus</i>
tobira	<i>Pittosporum tobira</i>
trumpet vine, violet	<i>Clytostoma callistegioides</i>
viburnum, Laurustinus	<i>Viburnum tinus</i>
virburnum, Sandankwa	<i>Virburnum suspensum</i>
weigela, oldfashioned	<i>Weigela florida</i>
wintercreeper	<i>Euonymus fortunei</i>
xylosma, Japanese	<i>Xylosma congestum</i>
yarrow	<i>Achillea</i> spp.
yaupon	<i>Ilex vomitoria</i>
yew <i>Taxus media</i>	
yew, Japanese	<i>Taxus cuspidata</i>
yewpine	<i>Podocarpus macrophyllus</i>
yucca, pendulous	<i>Yucca recurvifolia</i>
yucca, soaptree	<i>Yucca elata</i>
zinnia, common	<i>Zinnia elegans</i>

Surflan 0.25 A.S. may be used on the Following Container-Grown Plant Species:

Common Name	Scientific Name
andromeda	<i>Pieris japonica</i>
arborvitae, American	<i>Thuja occidentalis</i>
arborvitae, Oriental	<i>Platyclusus orientalis</i>

11/16

astilbe/false spirea	<i>Astilbe chinensis</i> and <i>A. chinensis</i> hybrids
barberry, Japanese	<i>Berberis thunbergii</i>
bellflower	<i>Campanula elatines</i>
blazing star	<i>Liatris spicata</i>
bleeding heart	<i>Dicentra spectabilis</i>
bottlebrush, lemon	<i>Callistemon citrinus</i>
boxwood, common	<i>Buxus sempervirens</i>
brush cherry	<i>Syzygium paniculata</i>
cleysera, Japanese	<i>Cleyera japonica</i>
cotoneaster, bearberry	<i>Cotoneaster dammeri</i>
cotoneaster, cranberry	<i>Cotoneaster apiculatus</i>
cotoneaster, parney	<i>Cotoneaster lacteus</i>
cotoneaster, rock	<i>Cotoneaster horizontalis</i>
crape Myrtle, common	<i>Lagerstroemia indica</i>
cryptomeria, Japanese	<i>Cryptomeria japonica</i>
cypress, Arizona	<i>Cupressus arizonica (glabra)</i>
cypress, Italian	<i>Cupressus sempervirens</i>
daylily	<i> Hemerocallis</i> spp.
dogwood, kousa	<i>Cornus kousa</i>
eastercactus	<i>Rhipsalidopsis gaertneri</i>
escallonia	<i>Escallonia exoniensis</i>
euonymus, evergreen	<i>Euonymus japonica</i>
euonymus, stringybark	<i>Euonymus fortunei</i>
fatshedera	<i>Fatshedera lizei</i>
firethorn	<i>Pyracantha, fortuneana</i>
firethorn, formosa	<i>Pyracantha skoidzumi</i>
firethorn, scarlet	<i>Pyracantha coccinea</i>
gardenia	<i>Gardenia jasminoides</i>
ginkgo	<i>Ginkgo biloba</i>
holly, Chinese	<i>Ilex cornuta</i>
holly, Japanese	<i>Ilex crenata</i>
jerseytea, redroot	<i>Ceanothus americanus</i>
juniper	<i>Juniperus</i> spp.
kumquat	<i>Fortunella</i> spp.
lilac, common	<i>Syringa vulgaris</i>
lilyturf, bigblue	<i>Liriope muscari</i>
lily-of-the-Nile	<i>Agapanthus africanus</i>
linden, little leaf	<i>Tilia cordata</i>
mockorange	<i>Philadelphus</i> spp.
myrtle, true	<i>Myrtus communis</i>
oak <i>Quercus</i> spp.	
oleander	<i>Nerium oleander</i>
orange, ornamental	<i>Citrus</i> spp.
pecan, ornamental	<i>Carya</i> spp.
photinia	<i>Photinia fraseri</i>
pine	<i>Pinus</i> spp.
pittosporum	<i>Pittosporum</i> spp.
privet, amur	<i>Ligustrum amurense</i>
privet, glossy	<i>Ligustrum lucidum</i>
privet, golden	<i>Ligustrum vicaryi</i>
privet, Japanese	<i>Ligustrum japonicum</i>
redbud	<i>Cercis canadensis</i>
raphiolepis (India hawthorn)	<i>Raphiolepis indica</i>
rhododendron	<i>Rhododendron</i> spp.
russian olive	<i>Elaeagnus angustifolia</i>

12/16

- | | |
|-----------------------|-----------------------------------|
| shrimp plant | <i>Justicia brandegeana</i> |
| spruce, Colorado | <i>Picea pungens</i> |
| sumac, African | <i>Rhus lancea</i> |
| sweetgum, American | <i>Liquidambar styraciflua</i> |
| trumpet vine, violet | <i>Clytostoma callistegioides</i> |
| viburnum, Laurustinus | <i>Viburnum tinus</i> |
| wintercreeper | <i>Euonymus fortunei</i> |
| yaupon | <i>Ilex vomitoria</i> |
| yucca, soaptree | <i>Yucca elata</i> |

Precaution

Ice Plant: When establishing unrooted ice plant on coarse soils in landscape plantings, use only the 1.5 pint of Surflan 0.25 A.S. or crop injury may occur. After the ice plant is well established, a second application may be made.

Broadcast Application Rates

Labeled Use Site	Length of Control	Surflan 0.25 A.S.	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (pt/1000 sq ft)
		(pt/1000 sq ft)		
Landscape ornamentals	2 - 4 months	1.1 to 1.5	2	5.9
	3 - 6 months	2.2	4	8.8
	4 - 8 months	3	4	8.8
Field-grown and container-grown ornamentals	2 - 4 months	1.5	3	5.9
	3 - 6 months	2.2	3	6.6
	4 - 8 months	3	3	8.8

Southern Turfgrass Areas

Surflan 0.25 A.S. is a preemergence herbicide, which is applied to established Southern turf including bahiagrass, bermudagrass, centipedegrass, established tall fescue, St. Augustinegrass and zoysiagrass for the control of annual grasses and some broadleaf weeds. Surflan 0.25 A.S. works by controlling weeds as the seeds germinate.

Surflan 0.25 A.S. will not harm nearby established ornamental trees and shrubs. Do not apply Surflan 0.25 A.S. on golf course putting greens.

General Use Precautions and Restrictions for Southern Turfgrass

Surflan 0.25 A.S. will not control established weeds.

Surflan 0.25 A.S. may injure turf that is under weakened conditions due to degree of establishment or winter climatic conditions. Apply Surflan 0.25 A.S. to healthy, established turf only.

Use Surflan 0.25 A.S. only as a part of a total turf management program that includes good fertilization practices.

Surflan 0.25 A.S. should not be applied in the spring or summer to tall fescue turfgrass reseeded the previous fall. In bermudagrass areas that have been overseeded with winter grasses, a spring application of Surflan 0.25 A.S. will thin the overseeded grasses.

Surflan 0.25 A.S. should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until these turfs are well established.

13/16

Surflan 0.25 A.S. should not be applied to newly hydromulched areas of bermudagrass until these areas are well established.

Do not spray Surflan 0.25 A.S. in combination with atrazine on turf.

Areas planted with bermudagrass variety "Sun Turf" in Northern Oklahoma have shown sensitivity to Surflan 0.25 A.S. treatments.

Do not apply Surflan 0.25 A.S. on golf course putting greens.

Grasses Controlled by Surflan 0.25 A.S.

Summer Annuals

Common Name	Scientific Name
barley, little	<i>Hordeum pusillum</i>
barnyardgrass (watergrass)	<i>Echinochloa crus-galli</i>
bluegrass, annual (poa)	<i>Poa annua</i>
brachiaria (signalgrass)	<i>Brachiaria</i> spp.
browntop panicum	<i>Panicum fasciculatum</i>
crabgrass	<i>Digitaria</i> spp.
(large crabgrass)	
(smooth crabgrass)	
crowfootgrass	<i>Dactyloctenium aegyptium</i>
cupgrass	<i>Eriochloa gracilis</i>
downy brome	<i>Bromus tectorum</i>
fall panicum	<i>Panicum dichotomiflorum</i>
(spreading panicgrass)	
foxtails	<i>Setaria</i> spp.
(bottlegrass)	
(bristlegrass)	
(giant foxtail)	
(green foxtail)	
(pigeongrass)	
(robust foxtail)	
(yellow foxtail)	
guineagrass	<i>Panicum maximum</i>
(narrowleaf panicum)	
johnsongrass (seedling only)	<i>Sorghum halepense</i>
jungerice	<i>Echinochloa colonum</i>
lovegrass, Mexican	<i>Eragrostis mexicana</i>
lovegrass, orcutt	<i>Eragrostis orcuttiana</i>
oat, wild	<i>Avena fatua</i>
ryegrass, annual (Italian)	<i>Lolium multiflorum</i>
sandbur, field	<i>Cenchrus incertus</i>
sprangletop, red	<i>Leptochloa filiformis</i>
Texas panicum	<i>Panicum texanum</i>
(buffalograss)	
(coloradograss)	
witchgrass	<i>Panicum capillare</i>

Winter Annuals

Common Name	Scientific Name
bluegrass, annual	<i>Poa annua</i>

14/16

Broadleaf Weeds Controlled by Surflan 0.25 A.S.

Summer Annuals

Common Name	Scientific Name
carpetweed	Mollugo verticillata
knotweed, prostrate	Polygonum aviculare
purslane, common	Portulaca oleracea

Winter Annuals

Common Name	Scientific Name
chickweed, common	Stellaria media
henbit	Lamium amplexicaule

Broadleaf Weeds Partially Controlled by Surflan 0.25 A.S.

Common Name	Scientific Name
groundsel, common	Senecio vulgaris
spurge, prostrate	Euphorbia humistrata
woodsorrel, yellow	Oxalis stricta

Surflan 0.25 A.S. will not control established weeds.

General Use Directions

Rates, Frequency and Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply Surflan 0.25 A.S. prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. In the absence of timely rainfall, irrigation can be used to move Surflan 0.25 A.S. into the soil surface. Any cultural practices that disturb the soil, such as aerification or verticutting, should be done prior to application of Surflan 0.25 A.S.

Application Rates, Number of Applications Allowed and Minimum Interval between Applications:

Warm Season Turf Areas	Surflan 0.25 A.S.		Minimum Time Between Applications (months)	Total Amount Allowed Per Year (pt/1000 sq ft)
	(pt/1000 sq ft)	(fl oz/1000 sq ft)		
All, except Florida	1.1 - 1.5	18 - 24	3	4.4
Florida	1.1	18	3	3.3

Application Directions

Surflan 0.25 A.S. can be applied in the spring for summer annual grass and broadleaf weed control, and in the fall for *Poa annua* and winter annual broadleaf weed control. Do not apply more than 1.1 pt (18 fl oz)/1000 sq ft per application in the state of Florida.

1. Spring Germinating Summer Annual Grasses and Broadleaf Weeds

For preemergence control, apply 1.5 pints (24 ounces) Surflan 0.25 A.S. per 1000 square feet per acre in late winter or early spring prior to the onset of conditions favorable for annual weed grass germination.

157/16

For application to smaller areas, apply the following amounts of Surflan 0.25 A.S.:

Amount Per 100 sq. ft.	Amount Per 500 sq. ft.	Amount Per 1000 sq. ft.
2 1/2 oz	12 1/2 oz	24 oz

2. Annual Bluegrass (*Poa Annua*) and Broadleaf Winter Annuals

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in these thin areas previously occupied by annual bluegrass. For preemergence control, apply 18 ounces Surflan 0.25 A.S. per 1000 square feet in the late summer or early fall prior to the expected germination of annual bluegrass and broadleaf winter annual weeds.

For application to smaller areas, apply the following amounts of Surflan 0.25 A.S.:

Amount Per 100 sq. ft.	Amount Per 500 sq. ft.	Amount Per 1000 sq. ft.
1 3/4 oz	8 3/4 oz	18 oz

Application Equipment

Apply Surflan 0.25 A.S. evenly over the turfgrass areas without skips, misses, or overlaps. For best results use equipment designed to apply liquid herbicides and insecticides. Calibrate application equipment prior to use, according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing spray uniformly.

Surflan 0.25 A.S. is an orange chemical. No marker dyes are needed as this orange characteristic will serve as a marker dye.

Reseeding

Chemicals that effectively control annual weeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 90-120 days after applying Surflan 0.25 A.S.. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation and fertilization be followed. For satisfactory reseeding results after using Surflan 0.25 A.S., the amount of seed used should be increased and equipment designed to place seed in contact with the soil (such as the Rogers Aero Seeder) should be employed.

Noncropland Areas

Surflan 0.25 A.S. plus Roundup and many other labeled herbicides may be used to control undesirable vegetation in noncropland areas such as roadsides, rights-of-way, etc. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions, and limitations before use.

Broadcast Application Rates

Labeled Use Site	Length of Control	Surflan 0.25 A.S.	Minimum Time Between Applications	Total Amount Allowed Per Year
		(pt/1000 sq ft)	(months)	(pt/1000 sq ft)
Noncropland	2 - 4 months	1.5	2	4.4
	6 - 8 months	3	4	8.8
	8 - 12 months	4.4	8	8.8

16/16

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. **UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.