

70506-53

01/03/2006

1/25



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

JAN 3 - 2006

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

ANN TILLMAN, Ph.D.
UNITED PHOSPHORUS INC.
423 RIVERVIEW PLAZA
TRENTON, NJ 08611

Dear Dr. Tillman:

SUBJECT: LABEL AMENDMENT - Oryzalin TRED
Snapshot 80 DF
EPA Registration No. 70506-53
Application Date: November 14, 2005

The label amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you add the following statement to the **Directions for Use**:

"For orchard crop, including citrus, pome fruits, stone fruits and tree nuts, apply product only as a strip treatment in the tree rows; do not apply to row middles or drive rows."

A stamped copy of labeling is enclosed for your records. Please submit one(1) copy of your final printed labeling before you release the product for shipment. **This amended labeling supersedes all previously accepted ones.**

Sincerely yours,

Joanne I. Miller
Joanne I. Miller
Product Manager (23)

Herbicide Branch
Registration Division (7505C)

Enclosure

2/25

Snapshot®

80 Dry Flowable

A selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Ground Covers/Perennials
- Ornamental Bulbs
- Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards
- Christmas Tree Plantations
- Non-Cropland

Active Ingredients:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomer	20.0%
oryzalin: 3,5-dinitro-N ⁴ ,N ⁴ -dipropylsulfanilamide	60.0%
Inert Ingredients	20.0%
Total	100.0%

Contains 0.80 pounds active ingredient per pound.

ACCEPTED
with **COMMENTS**
In EPA Letter Dated:

JAN 3 - 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

20506-53

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Precautionary Statements

Hazards to Humans and Domestic Animals

Causes Moderate Eye Irritation • Prolonged or frequently repeated contact may cause allergic reactions in some individuals.

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

3/25

Engineering Controls Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove personal protective equipment immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

First Aid

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to label booklet for Directions for Use, including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read Conditions of Sale and Limitation of Warranty and Liability inside label booklet.**

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 70506-53

EPA Est. _____

**UPI**

United Phosphorus, Inc.

423 Riverview Plaza

Trenton, NJ 08611

1-800-247-1557 • www.upi-usa.com

Specialty Herbicide

Net Contents

4/25

(Datapack cover):

Snapshot[®]

80 Dry Flowable

A selective preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

- Landscape Ornamentals
- Container Grown Ornamentals
- Field Grown Ornamentals
- Ground Covers/Perennials
- Ornamental Bulbs
- Non-Bearing Fruit and Nut Trees and Non-Bearing Vineyards
- Christmas Tree Plantations
- Non-Cropland

Active Ingredients:

isoxaben: N-[3-(1-ethyl-1-methylpropyl)-5-isoxazolyl]-2,6-dimethoxybenzamide and isomer	20.0%
oryzalin: 3,5-dinitro-N ⁴ ,N ⁴ -dipropylsulfanilamide	60.0%
Inert Ingredients	<u>20.0%</u>
Total	100.0%

Contains 0.80 pounds active ingredient per pound.

Keep Out of Reach of Children

CAUTION PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Agricultural Use Requirements
 Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

Refer to inside of label booklet for additional precautionary information including Personal Protective Equipment (PPE), User Safety Recommendations and Directions for Use including Storage and Disposal.

Notice: Read the entire label. Use only according to label directions. **Before buying or using this product, read Conditions of Sale and Limitation of Warranty and Liability inside label booklet.**

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300. Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

EPA Reg. No. 70506-53

Specialty Herbicide

EPA Est. _____
Net Contents

S/as

(Page 1 through end):

Table of Contents	Page
Precautionary Statements	
Hazards to Humans and Domestic Animals	
Personal Protective Equipment (PPE)	
User Safety Recommendations	
First Aid	
Environmental Hazards	
Directions for Use	
Agricultural Use Requirements	
Storage and Disposal	
General Information	
General Use Precautions	
Mixing and Application Directions	
Snapshot 80 Dry Flowable - Alone	
Snapshot 80 Dry Flowable - Tank Mixes	
Application	
Application Rates and Weeds Controlled or Suppressed by Snapshot 80 Dry Flowable	
Approved Uses:	
Special Use Precautions	
Ornamental Trees	
Ornamental Shrubs	
Ornamental Ground Covers	
Ornamental Bulbs	
Christmas Tree Plantations	
Noncropland Areas	
Conditions of Sale and Limitation of Warranty and Liability	

6/25

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes Moderate Eye Irritation • Prolonged or frequently repeated contact may cause allergic reactions in some individuals.

Avoid contact with eyes or clothing.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls Statements

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Users should remove personal protective equipment immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Wash the outside of gloves before removing.

First Aid

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Environmental Hazards

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate. Drift may result in reduced germination or emergence of non-target plants adjacent to treated area.


Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

NOTICE: The container cap for Snapshot 80 Dry Flowable herbicide may be used to measure the amount of product required to treat small areas. Use 2 to 4 measured container caps per 1000 sq ft.

7125

Please refer to Application Rates and Weeds Controlled section of label booklet for complete use directions.



When filled to this level, container cap contains approximately 1/2 oz of Snapshot 80 Dry Flowable.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for Agricultural Pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Entry Restrictions for Non-WPS Uses: Keep all persons, children and pets out of treated area until sprays have dried.

Storage and Disposal

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container. Do not store in direct sunlight. Do not store at temperatures above 120°F. In case of leak or spill, contain material and dispose as waste.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Dispose of in a sanitary landfill, or by incineration, in accordance with applicable regulations, or by burning if allowed by state and local authorities. If burned, stay out of smoke.

8/25

General Information

Snapshot 80 Dry Flowable herbicide is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in:

Landscape Ornamentals
 Container Grown Ornamentals
 Field Grown Ornamentals
 Ground Covers / Perennials
 Ornamental Bulbs

Non-Bearing Fruit and Nut Trees and
 Non-Bearing Vineyards
 Christmas Tree Plantations
 Non-cropland

Apply Snapshot 80 Dry Flowable anytime prior to germination of target weeds, or immediately after cultivation.

General Use Precautions

Snapshot 80 Dry Flowable controls weeds growing from seed. Snapshot 80 Dry Flowable does not control established weeds, weeds growing from stolens, rhizomes, or root pieces. Poor weed control may result if directions are not followed. Over-application may result in crop injury or excessive soil residue. Existing weeds should be controlled by cultivation or with postemergence herbicides. Weed residues, prunings and trash should be removed or thoroughly mixed into soil prior to treatment. Soil should be in good condition and free of clods at the time of application. A single rainfall or sprinkler irrigation of 0.5 inches or more, or flood irrigation is required to activate Snapshot 80 Dry Flowable. Optimum weed control is obtained when Snapshot 80 Dry Flowable is activated within 21 days of application. If rainfall or irrigation does not occur within 21 days of application and tillage is possible, Snapshot 80 Dry Flowable may be activated using cultivation equipment capable of uniformly mixing the herbicide into the upper 1 to 2 inches of soil.

Snapshot 80 Dry Flowable is orange in color and may cause temporary discoloration of sprayed surfaces. If this discoloration is undesirable, it may be altered by using a commercially available colorant such as Blazon or removed by spraying surface with water or washing with an industrial cleaner immediately after application. Snapshot 80 Dry Flowable may also be applied with mulch colorants, such as Mulch Magic or Nu-Mulch.

Treatment of Plant Species Not Listed on the Label for Snapshot 80 Dry Flowable

Users who wish to use Snapshot 80 Dry Flowable on plant species not recommended on this label may determine the suitability for such uses by making a trial application of Snapshot 80 Dry Flowable at a recommended rate to a small number of plants. Prior to using Snapshot 80 Dry Flowable on a larger number of plants, the treated plants should be observed for any sign of herbicidal injury during 30 to 60 days of normal growing conditions to determine if the treatment is non-injurious to the target plant species. **The user assumes responsibility for any plant damage or other liability resulting from use of Snapshot 80 Dry Flowable on plant species not recommended on this label.**

Apply Snapshot 80 Dry Flowable only to established plantings. **Do not apply** Snapshot 80 Dry Flowable to newly transplanted ornamentals, nursery stock, ground covers, ornamental bulbs, non-bearing fruit[†] and nut trees and non-bearing vineyards or Christmas trees until soil or potting media has been settled by packing and irrigation or rainfall and no cracks are present or plant injury may occur.

[†]**Note:** Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Do not apply Snapshot 80 Dry Flowable to turf.

Do not aerially apply this product.

9/25

Do not graze or feed forage from treated areas to livestock.

In Arizona: The state of Arizona has not approved Snapshot 80 Dry Flowable for use on plants grown for commercial production such as: turf sod farms, ornamentals and nursery stock grown for resale, and non-bearing fruit and nut trees and vineyards.

Application Directions

Apply Snapshot 80 Dry Flowable as a directed spray to the soil surface or overtop of plants using a properly calibrated low pressure herbicide sprayer that provides a uniform distribution. Spray tip screens should be no finer than 50 mesh. In-line screens and strainers should be no finer than 16 mesh (50 mesh is finer than 16 mesh). Apply Snapshot 80 Dry Flowable in 10 to 200 gallons of water carrier per acre (0.25 to 5 gallons per 1000 sq ft). As spray volume decreases, the importance of accurate calibration and uniform application increases. Calibrate application equipment prior to use, according to manufacturer's directions to ensure proper calibration and uniform application. Maintain continuous agitation from mixing through application. Take precautions to avoid spray drift when applying Snapshot 80 Dry Flowable. Drift may result in reduced germination or emergence of non-target plants adjacent to the treated area. Avoid spray pattern skips and overlaps that may result in incomplete coverage or over application.

Chemigation: Do not apply Snapshot 80 Dry Flowable through any type of irrigation system.

Snapshot 80 Dry Flowable - Alone

Check to be sure spray equipment is clean and not contaminated. Using clean water, fill spray tank to 1/2 the final volume required and start agitation. Slowly add the correct quantity of Snapshot 80 Dry Flowable to the spray tank, continue agitation and complete filling the tank. Maintain agitation during filling and through application. Agitation should be sufficient to create a rippling or rolling action on the liquid surface. A sparger pipe generally provides the best agitation. If spraying and agitation is interrupted, Snapshot 80 Dry Flowable may settle to the bottom of the spray tank. If settling occurs, material must be resuspended before spraying continues. Clean spray tank, lines, screens and nozzles thoroughly after use.

Tank Mixes with Snapshot 80 Dry Flowable

Tank mix combinations of Snapshot 80 Dry Flowable plus glyphosate or other labeled herbicides may be used to control undesirable vegetation. Applied as directed, these tank mixes will provide control of susceptible weed species listed on the respective labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Tank Mixing

Add Snapshot 80 Dry Flowable to the spray tank as described above and fill the spray tank to 3/4 of final volume required. Add other product formulations in the following order: (1) dry flowables, (2) wettable powders, (3) aqueous suspensions, (4) flowables, (5) liquids, (6) solutions and emulsifiable or liquid concentrates. To insure proper mixing, allow two to three minutes agitation between the addition of each product, and agitate continuously until the product(s) are completely dispersed in water. Maintain agitation during filling and through application. If a buildup of material is observed on the walls of the spray tank, wash the tank with soapy water between fillings, rinse and then continue the spraying operation. Clean spray tank, lines, screens and nozzles thoroughly after use. Follow label directions for each material added to the tank.

Snapshot 80 Dry Flowable Plus glyphosate: Tank mix combinations of Snapshot 80 Dry Flowable plus glyphosate are recommended to control existing undesirable vegetation. Applied as directed, Snapshot 80 Dry Flowable plus glyphosate will provide postemergence control of susceptible weed species listed on the label for glyphosate and residual preemergence control of susceptible weed species listed on the label for Snapshot 80 Dry Flowable. Refer to the label for glyphosate for specific use directions, precautions and limitations before use.

10/25

Precautions:

Do not apply sprays containing glyphosate over the top of ornamental plants. Extreme care must be exercised to prevent contact of sprays containing glyphosate with foliage and stems of trees, shrubs, or other desirable vegetation since severe damage or death may result. Note: If spraying with glyphosate in areas adjacent to desirable plants, use a shield to prevent spray from contacting foliage and stems of desirable plants.

Application Rates and Weeds Controlled or Suppressed by Snapshot 80 Dry Flowable

Snapshot 80 Dry Flowable Broadcast Rates				
per acre		per 1,000 sq ft		
Pounds	Bottles	Ounces	Measured Caps	Tablespoons
2.5	2	1.0	2	4
3.75	3	1.5	3	6
5.0	4	2.0	4	8

Weeds controlled when applied at 2.5 pounds (2 bottles) per acre (1.0 oz or 2 measured container caps per 1000 sq ft).

Common Name	Scientific Name
aster, slender	<i>Aster exilis</i>
barnyardgrass	<i>Echinochloa crus-galli</i>
bluegrass, annual	<i>Poa annua</i>
bursage, annual	<i>Ambrosia acanthicarpa</i>
celery, wild	<i>Apium eptophyllum</i>
chickweed, common	<i>Stellaria media</i>
clover, white	<i>Trifolium repens</i>
crowfootgrass	<i>Dactyloctenium aegyptium</i>
cudweed, purple	<i>Gnaphalium purpureum</i>
fiddleneck, coast	<i>Amsinckia intermedia</i>
filaree, redstem	<i>Erodium cicutarium</i>
fleabane, blackleaved	<i>Conyza bonariensis</i>
fleabane, dwarf	<i>Conyza ramosissima</i>
foxtail	<i>Setaria spp.</i>
groundcherry, lanceleaf	<i>Physalis lanceifolia</i>
henbit	<i>Lamium amplexicaule</i>
horseweed	<i>Conyza canadensis</i>
knotweed, prostrate	<i>Polygonum aviculare</i>
lambsquarters, common	<i>Chenopodium album</i>
mallow, little	<i>Malva parviflora</i>
mustard, Indian	<i>Brassica juncea</i>
mustard, wild	<i>Sinapis arvensis</i>
nightshade, black	<i>Solanum nigrum</i>
pepperweed, Virginia	<i>Lepidium virginicum</i>
pigweed	<i>Amaranthus spp.</i>
pineappleweed	<i>Matricaria matricarioides</i>
plantain, slender	<i>Plantago elongata</i>
purslane, common	<i>Portulaca oleracea</i>
radish, wild	<i>Raphanus raphanistrum</i>
ragweed, common	<i>Ambrosia artemisiifolia</i>
rocket, London	<i>Sisymbrium irio</i>
rockpurslane, desert	<i>Calandrinia ciliata</i>
shepherdspurse	<i>Capsella bursa-pastoris</i>

sibara	<i>Sibara virginica</i>
smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
sowthistle, annual	<i>Sonchus oleraceus</i>
speedwell, purslane	<i>Veronica peregrina</i>
telegraphplant	<i>Heterotheca grandiflora</i>
thistle, Russian	<i>Salsola iberica</i>

In addition to the weeds listed at the 2.5 pound per acre rate, the following weeds will be controlled at 3.75 pounds (3 bottles) per acre (1.5 oz or 3 measured container caps per 1000 sq ft).

Common Name	Scientific Name
aster, heath	<i>Aster ericoides</i>
bittercress	<i>Cardamine oligosperma</i>
bittercress, hairy	<i>Cardamine hirsuta</i>
brassbuttons, southern	<i>Cotula australis</i>
carrot, wild	<i>Daucus carota</i>
chamberbitter	<i>Phyllanthus urinaria</i>
chickweed, mouseear	<i>Cerastium vulgatum</i>
crabgrass	<i>Digitaria</i> spp.
dandelion	<i>Taraxacum officinale</i>
eclipta	<i>Eclipta prostrata</i>
galinsoga, hairy	<i>Galinsoga ciliata</i>
geranium, Carolina	<i>Geranium carolinianum</i>
goosegrass	<i>Eleusine indica</i>
groundsel, common	<i>Senecio vulgaris</i>
jugleryce	<i>Echinochloa colonum</i>
ladysthumb	<i>Polygonum persicaria</i>
lettuce, prickly	<i>Lactuca scariola</i>
mallow, dwarf	<i>Malva rotundifolia</i>
maretail	<i>Hippuris vulgaris</i>
mayweed	<i>Anthemis cotula</i>
morningglory, ivyleaf	<i>Ipomoea hederacea</i>
mustard, black	<i>Brassica nigra</i>
oat, wild	<i>Avena fatua</i>
pennywort	<i>Hydrocotyle</i> spp.
phyllanthus, long-stalk	<i>Phyllanthus tenellus</i>
plantain, bracted	<i>Plantago aristata</i>
plantain, broadleaf	<i>Plantago major</i>
plantain, buckhorn	<i>Plantago lanceolata</i>
pokeweed, common	<i>Phytolacca americana</i>
rockpurslane, redmaids	<i>Calandrinia ciliata</i> var. <i>menziesii</i>
ryegrass, annual	<i>Lolium multiflorum</i>
sida, prickly	<i>Sida spinosa</i>
signalgrass	<i>Bracharia</i> spp.
sorrell, red	<i>Rumex acetosella</i>
speedwell, thymeleaf	<i>Veronica serpyllifolia</i>
sprangletop, red	<i>Leptochloa filiformis</i>
spurge, hyssop	<i>Euphorbia hyssopifolia</i>
spurge, spotted	<i>Euphorbia maculata</i>
sweetclover, yellow	<i>Melilotus officinalis</i>
tansymustard, green	<i>Descurainia pinnata</i> spp. <i>brachycarpa</i>
velvetleaf	<i>Abutilon theophrasti</i>
witchgrass	<i>Panicum capillare</i>

12/25

woodsorrel, yellow *Oxalis stricta*

In addition to the weeds listed at the 2.5 pound per acre and 3.75 pound per acre rates, the following weeds will be controlled at 5 pounds (4 bottles) per acre (2.0 oz or 4 measured container caps per 1000 sq ft).

Common Name	Scientific Name
barley, little	<i>Hordeum leporinum</i>
burclover, California	<i>Medicago polymorpha</i>
carpetweed	<i>Mollugo verticillata</i>
cupgrass, Southwestern	<i>Eriochloa gracilis</i>
dogfennel	<i>Eupatorium capillifolium</i>
eveningprimrose	<i>Oenothera</i> spp.
fescue, rattail	<i>Vulpia myuros</i>
filaree, whitestem	<i>Erodium moschatum</i>
goosefoot, nettleleaf	<i>Chenopodium murale</i>
jimsonweed	<i>Datura stramonium</i>
knotweed, silversheath	<i>Polygonum argyrocoleon</i>
kochia	<i>Kochia scoparia</i>
medic, black	<i>Medicago lupulina</i>
mullein, turkey	<i>Eremocarpus setigerus</i>
nettle, burning	<i>Urtica urens</i>
ox tongue, bristly	<i>Picris echioides</i>
panicum, browntop	<i>Panicum fasciculatum</i>
panicum, fall	<i>Panicum dichotomiflorum</i>
pimpernel, scarlet	<i>Anagallis arvensis</i>
puncturevine	<i>Tribulus terrestris</i>
pusley, Florida	<i>Richardia scabra</i>
sandbur, field	<i>Cenchrus incertus</i>
sowthistle, spiny	<i>Sonchus asper</i>
spurge, petty	<i>Euphorbia pepi</i>
spurge, prostrate	<i>Euphorbia humistrata</i>
stinkgrass	<i>Eragrostis cilianensis</i>
sunflower	<i>Helianthus</i> spp.
swinecress	<i>Coronopus didymus</i>
thistle, musk	<i>Carduus nutans</i>
willoweed, panicle	<i>Epilobium paniculatum</i>
woodsorrel, creeping	<i>Oxalis corniculata</i>

In addition to the weeds controlled, the following weeds will be partially controlled or suppressed at 5.0 pounds (4 bottles) per acre (2.0 oz or 4 measured container caps per 1000 sq ft).

Common Name	Scientific Name
bindweed, field	<i>Convolvulus arvensis</i>
dock, curly	<i>Rumex crispus</i>
johnsongrass (seedling)	<i>Sorghum halapense</i>
mallow, common	<i>Malva neglecta</i>
mallow, Venice	<i>Hibiscus trionum</i>
milkweed, climbing	<i>Sarcostemma cynanchoides</i>
milkweed, honeyvine	<i>Ampelamus albidus</i>
morningglory, tall	<i>Ipomoea purpurea</i>
panicum, Texas	<i>Panicum texanum</i>
wheat, volunteer	<i>Triticum</i> spp.

13/25

Approved Uses

Ornamentals

Snapshot 80 Dry Flowable herbicide is a preemergence herbicide for control of certain annual grasses and broadleaf weeds in: landscape ornamentals; container grown ornamentals; field grown ornamentals; ground covers/perennials; non-bearing fruit and nut trees and non-bearing vineyards. Refer to the "General Information" section of this label for general use precautions, mixing and application directions, and information on application rates and weeds controlled.

Special Use Precautions

To avoid possible injury, do not apply Snapshot 80 Dry Flowable to:

- Nursery, forest, or Christmas tree; seedling beds, cutting beds, or transplant beds.
- Unrooted liners or cuttings that have been planted in pots for the first time.
- Rooted liners until they have been removed from their original growing containers and placed in new containers at least two weeks prior to treatment.
- Pots less than four inches wide.
- Bedding plants or areas where bedding plants will be planted or transplanted within one year following application.
- Ground covers until they are established and well rooted.
- Ornamental plantings where the likelihood of runoff onto turf areas exists.

Application of Snapshot 80 Dry Flowable over the top of plants with newly forming buds may cause injury. In this situation a directed spray application is recommended.

Where loss of stand has occurred in field grown ornamentals, use untreated soil as fill around roots when replacing plants or injury may occur.

Labeled Use Site	Snapshot 80 Dry Flowable (lb/acre)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (lb/acre)
• Landscape Ornamentals	2.5	2	13.3
• Field-grown and container-grown ornamentals	3.75	3	15
	5	4	15

Note: Injury on the following plant species has been observed following applications of Snapshot 80 Dry Flowable and use is not recommended.

Ajuga spp. (bugle)
Brassica spp. (mustard)
Deutzia gracilis (slender deutzia)
Euonymus alatus 'compacta' (dwarf burning bush)
Euphorbia spp. (spurge)
Hydrangea spp. (hydrangea)
Iberis spp. (candytuft)
Juniperus horizontalis 'Prince of Wales' (Prince of Wales Juniper)
Melaleuca quinquenervia (cajeput tree)
Pseudotsuga menziesii (Douglas-fir)
Rhododendron carolinianum (Carolina rhododendron)
Rhododendron catawbiense 'Roseum elegans' (Roseum elegans rhododendron)

M/25

Sedum spp. (stonecrop)
Thuja occidentalis 'Techny' (Techny arborvitae)
Tsuga canadensis (eastern hemlock)
Yucca recurvifolia (green yucca)

Ice Plant: When establishing unrooted ice plant on coarse textured soils, do not exceed the 3.75 pound per acre rate of Snapshot 80 Dry Flowable or injury may occur.

**Snapshot 80 Dry Flowable may be used on
the following established ornamental plant species:**

(Note: Limitations on recommended treatment methods).

Trees

Recommended Treatment Method:

C = Container Grown

F = Field Grown

Scientific Name	Common Name	
<i>Abies balsamea</i>	Fir, balsam	C,F
<i>Abies concolor</i>	Fir, white	F
<i>Abutilon hybridum</i>	Albus-flowering maple	C,F
	Luteus-flowering maple	C,F
	Roseus-flowering maple	C,F
	Tangerine-flowering maple	C,F
	Vesuvius red-flowering maple	F
<i>Acer ginnala</i>	Flame maple	F
<i>Acer rubrum</i>	Red maple	F
	Red sunset maple	F
<i>Acer saccharinum</i>	Silver maple	C,F
<i>Areacastrum romanzoffianum</i>	Queen palm	C,F
<i>Betula nigra</i>	Birch, river	C,F
<i>Betula papyrifera</i>	Paper birch	F
<i>Brachychiton populneus</i>	Bottle tree	C,F
<i>Bucida buceras</i>	Black olive	F
<i>Ceratonia siliqua</i>	Carob †	F
<i>Cercis canadensis</i>	Redbud	C,F
<i>Chamaecyparis obtusa</i>	Filicoides-fernspray cypress	F
<i>Chamaecyparis obtusa</i>	Gracilis-slender Hinoki cypress	F
<i>Chamaecyparis pisifera</i>	Sawara-false cypress	F
	Squarrosa-moss cypress	F
<i>Chamaedorea cataractarum</i>	Cat Palm	F
	Palm	C,F
<i>Chamaedorea costaricana</i>	Palm	C,F
<i>Chamaedorea elegans</i>	Parlor palm	C,F
<i>Cornus florida</i>	Cloud nine dogwood	C,F
	Dogwood, flowering	C,F
<i>Crataegus viridis</i>	Green hawthorn	F
<i>Cryptomeria japonica</i>	Cryptomeria, Japanese	C,F
<i>Cupaniopsis anacardioides</i>	Carrot wood	F
<i>Cupressus glabra</i>	Arizona cypress	F
<i>Elaeagnus angustifolia</i>	Russian olive	C,F
<i>Eucalyptus camaldulensis</i>	Red gum eucalyptus	F
<i>Eucalyptus cinerea</i>	Eucalyptus, mealy	F
	Silver dollar eucalyptus	F
<i>Eucalyptus microtheca</i>	Coolibah tree	C,F
<i>Eucalyptus sideroxylon</i>	Eucalyptus, red ironbark	F
<i>Ficus benjamina</i>	Ficus	C,F

15/25

	Mini ficus	C,F
Fraxinus udhei	Shamel ash	C,F
Ginkgo biloba	Ginkgo (Maidenhair tree)	F
Gleditsia triacanthos	Honey locust	F
	Shademaster honey locust	F
Heteromeles arbutiflora	Toyon	F
Illicium floridanum	Florida anise-tree	C,F
Juniperus virginiana	Redcedar, Eastern	C,F
Liquidambar styraciflua	Sweetgum, American	F
Magnolia grandiflora	Magnolia, Southern	C,F
Morus alba	White mulberry	F
Musa acuminata	Banana †	C,F
Oxydendrum arboreum	Sourwood	C,F
Picea abies	Pendula-weeping Norway spruce	C,F
	Repens-spreading Norway spruce	C,F
	Spruce, Norway	C,F
Picea glauca	Conica-dwarf Alberta spruce	C,F
Picea glauca conica	Dwarf Alberta spruce	F
Picea pungens	Dwarf globe blue spruce	C,F
	Glauca-Colorado blue spruce	C,F
	Hoopsii-Hoop's blue spruce	C,F
	Koster-Koster blue spruce	F
Pinus aristata	Bristlecone pine	F
Pinus canariensis	Canary Island pine	F
Pinus contorta	Shore pine, beach pine	F
Pinus eldarica	Eldarica pine	F
Pinus leucodermis	Bosnian pine	C,F
Pinus mugo	Pumilio- shrubby Swiss mountain pine	C,F
Pinus nigra	Austrian black pine	C,F
Pinus radiata	Monterey pine	F
Pinus strobus	Eastern white pine	C,F
	White pine	C,F
Pinus sylvestris	Columnar Scotch pine	C,F
	Scotch pine	C,F
Pinus thunbergiana	Japanese black pine	C,F
Platanus occidentalis	American sycamore	F
Platanus racemosa	Califorina sycamore	F
Podocarpus spp.	Podocarpus	F
Populus deltoides	Cottonwood	F
Prosopis chilensis	Chilean mesquite	C,F
Prunus yedoensis	Yoshino flowering cherry	F
Quercus ilicifolia	Bear oak	F
Quercus palustris	Pin oak	F
Quercus phellos	Willow oak	C,F
Quercus rubra	Red oak	C,F
Quercus virginiana	Live oak	C,F
Salix babylonica	Babylon weeping willow	F
	Corkscrew willow	F
Sequoiadendron giganteum	Giant sequoia	F
Swietenia mahogani	Mahogany	F
Tabebuia caraiba	Yellow tab	F
Ulmus parvifolia	Chinese elm	F
Washingtonia robusta	Mexican fan palm	F

† No portion of the plants from treated area may be used for food or feed.

16/25

Shrubs**Recommended Treatment Method:****C = Container Grown****F = Field Grown**

Scientific Name	Common Name	
Abelia grandiflora	Edward Goucher abelia	C,F
	Glossy abelia	C,F
Acacia abyssinica	Abyssinica acacia	C,F
Acacia redolens	Acacia, prostrate	C,F
Acacia stenophylla	Shoestring acacia	C,F
Acalypha wilkesiana	Copper leaf	C,F
Acer ginnala	Amur maple	C,F
Acer palmatum	Coral bark Japanese maple	C,F
Acer palmatum	Dwarf Japanese maple	C,F
Agave americana	Century plant	F
Astilbe arendsii	False spiraea	C,F
Astilbe chinensis	Astilbe/false spirea	C,F
Athyrium nipponicum	Japanese painted fern	C,F
Baccharis pilularis	Coyotebush	F
Berberis gladwynensis	William Penn barberry	C,F
Berberis mentorensis	Mentor barberry	C,F
Berberis thunbergii	Aurea-golden Japanese barberry	F
	Crimson pygmy barberry	C,F
	Atropurea-redleaf Japanese barberry	F
Bougainvillea spp.	Barbara Karst	C,F
	California gold	C,F
	Pink pixie	C,F
	Scarlet O'Hara	C,F
	Temple fire	C,F
	Texas dawn	C,F
Buxus microphylla japonica	Boxwood, Japanese	C,F
Buxus microphylla Koreana	Korean boxwood	F
Buxus sempervirens	Boxwood, common	C,F
Callistemon citrinus	Bottlebrush, lemon	F
Callistemon viminalis	Weeping bottlebrush	C,F
Calluna vulgaris	Spring torch scotch heather	C,F
Camellia japonica	Camellia	C,F
Cassia artemisioides	Cassia, feathery	C,F
Ceanothus spp.	Wild lilac	F
Cephalotaxus drupacea	Plum yew	C,F
Cerastium tomentosum	Snow-in-summer	C,F
Chamaecyparis obtusa	Kosteri cypress	C,F
	Nana-dwarf Hinoki cypress	C,F
	Torulosa cypress	C,F
Chamaecyparis pisifera	Baileyi-dogwood	F
	Flaviramea-dogwood	F
	Squarrosa Minima cypress	C,F
	Filifera-thread cypress	C,F
Chrysalidocarpus lutescens	Areca palm	F
Clethra alnifolia	Summersweet	C,F
Cleyera japonica	Cleyera, Japanese	C,F
Cornus alba	Sibirica-Siberian dogwood	C,F
Cornus stolonifera	Baileyi-red-osier dogwood	F
	Flaviramea-yellowtwig dogwood	F
Cotinus coggygria	Royal purple smoke tree	C,F
Cotinus dammeri	Coral beauty smoke tree	C,F

17/25

	Eichholz smoke tree	C,F
Cotoneaster adpressus	Praecox-early cotoneaster	C,F
Cotoneaster apiculatus	Cotoneaster, cranberry	C,F
Cotoneaster congestus	Cotoneaster, Pyrenees	C,F
Cotoneaster dammeri	Cotoneaster, bearberry	C,F
Cotoneaster himalayan	Himalayan cotoneaster	C,F
Cotoneaster horizontalis	Cotoneaster, rock	C,F
Cycas revoluta	Sago palm	C,F
Cytisus praecox	Hollandia-warminster broom	C,F
Cytisus scoparius	Lena-Scotch broom	C,F
Cytisus spp.	Holandia-Scotch broom	F
Daphne odora	Fragrant daphne	C,F
Deutzia crenata	Nakiana-dwarf deutzia	C,F
Dodonea viscosa	Hopseed bush	F
Elaeagnus pungens	Fruitland silver berry	C,F
Erica cinerea	Purple bell heather	C,F
Erica vagans	Cornish heather	C,F
Erica x darleyensa	Mediterranean pink heather	C,F
Euonymus fortunei	Canadale gold euonymus	C,F
	Emerald'n gold euonymus	F
	Sunspot euonymus	C,F
Euonymus japonica	Silver king euonymus	F
	Variogated evergreen euonymus	C,F
Euonymus kiatschovica	Spreading euonymus	C,F
Forsythia intermedia	Forsythia, border	F
Gardenia jasminoides	August beauty gardenia	C,F
	Gardenia	C,F
	Radican gardenia	C,F
Gaultheria shallon	Salal/lemon leaf	C,F
Gelsemium sempervirens	Carolina jessamine	C,F
Genista pilosa	Woadwaxen	C,F
Hibiscus rosa-sinensis	Ross Estey-hibiscus	C,F
Hibiscus syriacus	Rose of Sharon, Red Bird	C,F
	Rose of Sharon, Red Heart	F
	Rose of Sharon, Woodbridge	C,F
Ilex aquifolium	Balkans holly	F
	Gold coast holly	F
Ilex aquipernyi	San Jose holly	C,F
Ilex attenuata	Savannah holly	C,F
Ilex cornuta	Burford holly	C,F
	Dwarf Burford holly	C,F
	Needle point holly	C,F
Ilex crenata	Compacta-dwarf Japanese holly	C,F
	Convexa holly	C,F
	Dwarf Chinese holly	C,F
	Green luster holly	C,F
	Helleri-Heller's Japanese holly	C,F
	Hetzii's Japanese holly	C,F
	Stokesii Japanese holly	C,F
Ilex glabra	Compacta-compact inkberry holly	C,F
	Nordica-inkberry holly	C,F
Ilex meserveae	Blue boy holly	C,F
	Blue girl holly	C,F
	Ebony magic holly	F
Ilex vomitoria	Nana-dwarf yaupon holly	C,F

18/25

	Pendula-weeping yaupon holly	C,F
	yaupon holly	C,F
Illicium annisatum	Mystery gardenia	C,F
Itea ilicifolia	Henry Garnet holly leaf sweetspire	C,F
Ixora collinea	Ixora	C,F
Juniperus chinensis	hollywood juniper	C,F
	Media-old gold juniper	C,F
	pfitzer juniper	C,F
	Pfitzerana glauca-blue juniper	C,F
	Pfitzerana-pfitzer juniper	C,F
	Sea green juniper	F
	Torulosa-hollywood juniper	C,F
Juniperus conferta	Emerald sea shore juniper	C,F
	Shore juniper	C,F
Juniperus horizontalis	Andorra juniper	C,F
	Bar Harbor juniper	C,F
	Blue rug juniper	C,F
	Creeping juniper	C,F
	Dwarf Andorra juniper	C,F
	Huntington blue juniper	C,F
	Plumosa-Andorra juniper	C,F
	Wiltonii-blue carpet juniper	C,F
Juniperus procumbens	Nana-dwarf Japaneses garden juniper	C,F
Juniperus prostrata	Prostrata juniper	C,F
Juniperus sabina	Broadmoor juniper	C,F
	Foemina-Hicks juniper	C,F
	Savin juniper	C,F
	Tamariscifolia-Tam juniper	C,F
Juniperus scopulorum	Emerald green juniper	F
Juniperus squamata	Blue juniper	C,F
	Blue star juniper	C,F
	Parsonii juniper	C,F
Kalmia latifolia	Laurel, mountain	C,F
Lagerstroemia indica	Crape myrtle	C,F
Lantana spp.	Lantana	C,F
Leucothoe axillaris	Leucothoe, coast	C,F
Leucothoe fontanesiana	Leucothoe, drooping	C,F
Ligustrum japonicum	Privet, Japanese	C,F
	Wax ligustrum	C,F
	Yellow tip ligustrum	C,F
Ligustrum lucidum	Privet, glossy	C,F
Ligustrum ovalifolium	California privet	F
Ligustrum texanum	Howardi privet	C,F
	Wax leaf privet	C,F
Ligustrum vicaryi	Privet, golden	F
	Vicary golden privet	F
Ligustrum vulgare	Lodense privet	C,F
Livistona chinensis	Chinese fountain palm	F
Lonicera fragrantissima	Winter honeysuckle	C,F
Lonicera periclymenum	Flowering woodbine	C,F
	Serotina woodbine	C,F
Lonicera sempervirens	Trumpet honeysuckle	C,F
Mahonia bealei	Leather leaf mahonia	C,F
Mahonia repens	Creeping mahonia	C,F
Myrica cerifera	Wax myrtle	C,F

19/05

Nandina domestica	Compacta-dwarf heavenly bamboo	C,F
	Harbour dwarf-heavenly bamboo	C,F
	Heavenly bamboo (Nandina)	C,F
	Nana compacta-heavenly bamboo	C,F
	Nana purpurea-heavenly bamboo	C,F
	Woods dwarf-heavenly bamboo	C,F
Nerium oleander	Hardy red oleander	C,F
	Oleander	C,F
	Ruby lace oleander	C,F
Osmanthus fortunei	Fortunes osmanthus	C,F
Phoenix roeeloelenii	Pigmy date palm	C,F
Photinia fraseri	Fraser's photinia	C,F
Pieris japonica	Lily-of-the-valley	C,F
	Mountain fire lily-of-the-valley	C,F
	Snowdrift lily-of-the-valley	C,F
	Temple bells lily-of-the-valley	C,F
	Valley rose lily-of-the-valley	C,F
	Valley valentine lily-of-the-valley	C,F
Pieris japonica x forestii	Forest flame lily-of-the-valley	C,F
Pinus mugo	Mugo-mugho pine	C,F
Pittosporum tobira	Green pittosporum	C,F
	Wheeler's dwarf pittosporum	C,F
Plumbago ariculata	Blue cape plumbago	F
Plumbago capensis	Plumbago	C,F
Podocarpus macrophyllus	Yewpine	C,F
Polystichum polyblepharum	Tassel fern	C,F
Potentilla fragiformis	Cinquefoil	F
Potentilla fruticosa	Cinquefoil	C,F
	Gold drop potentilla	F
	Goldfinger potentilla	C,F
	Red ace potentilla	C,F
	Sunset potentilla	C,F
	Tangerine potentilla	C,F
Potentilla verna	Spring cinquefoil	C,F
Prunus glandulosa	Dwarf pink flowering almond	C,F
Pyracantha fortuneana	Lolendei Monrovia pyracantha	C,F
	Monon pyracantha	F
	Red elf hybrid pyracantha	F
	Rutgers hybrid pyracantha	C,F
	Santa Cruz pyracantha	C,F
	Victory pyracantha	F
Raphiolepis indica	Charisma-Monruce raphiolepis	C,F
	Enchantress-Moness raphiolepis	F
	Raphiolepis (India hawthorn)	F
	Springtime-Monme raphiolepis	F
Raphiolepis ovata	Roundleaf raphiolepis	C,F
Rhododendron calendulaceum	Cannon's double azalea	C,F
	Flame azalea	F
	Golden flare azalea	C,F
	Klondike azalea	C,F
Rhododendron campylocarpum	Butterfly rhododendron	F
Rhododendron carolinianum x daurium	PJM rhododendron	C,F
Rhododendron catawbiense	Catawba album rhododendron	C,F
	Catawba rhododendron	C,F
	Lord Roberts rhododendron	C,F

20/25

	Rocket rhododendron	C,F
Rhododendron caucasicum x ponticum	Cunningham White rhododendron	C,F
Rhododendron exbury	Cannon's double azalea	C,F
	Golden flare azalea	C,F
	Klondike azalea	C,F
Rhododendron forrestii repens	Gomer Waterer rhododendron	C,F
Rhododendron forrestii x griersonianum	Elizabeth rhododendron	C,F
Rhododendron griffithianum	Jean Marie rhododendron	C,F
Rhododendron hybrid spp.	America rhododendron	C,F
	English Roseum rhododendron	F
	Nova Zembla rhododendron	C,F
	Scintillation rhododendron	C,F
Rhododendron impeditum	Rhododendron	C,F
Rhododendron indica	Formosa azalea	C,F
	Waucabusa azalea	C,F
Rhododendron kaempferi	Blue danube azalea	C,F
Rhododendron kerume	Coral bells azalea	C,F
	Hino crimson azalea	C,F
	Hino pink azalea	C,F
	Mildred azalea	C,F
	Snow azalea	C,F
Rhododendron maximum	Rhodie max (rosebay)	C,F
Rhododendron mucronulatum	Rhododendron	F
Rhododendron obtusum	Hino-crimson azalea	C,F
Rhododendron ponticum	Chionoides rhododendron	C,F
	Daphnoides rhododendron	C,F
Rhododendron racemosum	Dwarf scarlet wonder rhododendron	C,F
	Tribly rhododendron	C,F
	Unique rhododendron	C,F
	Vulcan rhododendron	C,F
Rhododendron sassthigiatim x carolinianum	Ramapo rhododendron	C,F
Rhododendron satuski	Gumpo pink azalea	C,F
	Higasa azalea	F
	Reijn azalea	C,F
Rhododendron spp. hybrids	Carror azalea	C,F
	Fashion azalea	C,F
	Gerard Christina azalea	F
	Girard Roberta azalea	C,F
	Golden flare exbury azalea	F
	Helmut vogel azalea	C,F
	Hersey red azalea	F
	Hot shot azalea	C,F
	Hume azalea	F
	Inga azalea	F
	Irene Koster azalea	C,F
	President Clay azalea	C,F
	Tradition azalea	C,F
Rhus lancea	Sumac, African	C,F
Rosa rugosa	Ramanas rose	C,F
Rosmarinus officinalis	Rosemary	F
Skimmia japonica	Japanese skimmia	C,F
Skimmia revesiana	Reeve's skimmia	C,F
Spiraea bumalda	Anthony Waterer spiraea	C,F
Spiraea japonica	Dolchia spiraea	C,F

21/25

	Japanese alpine spiraea	C,F
	Shirobana spiraea	C,F
Spiraea vanhouttei	Bridal wreath	C,F
Syringa rothomagensis	Chinese lilac	C,F
Syringa vulgaris	Lilac, common	F
Taxus cuspidata	Yew, Japanese	F
Tecomaria capensis	Cape honeysuckle	C,F
Ternstroemia gymnanthera	Ternstroemia, Japanese	C,F
Thuja occidentalis	Emerald arborvitae	C,F
	Globosa-globe arborvitae	C,F
	Little giant-dwarf arborvitae	C,F
	Nigra-dark American arborvitae	C,F
	Pyramidalis arborvitae	C,F
	Rheingold arborvitae	C,F
	Woodwardii arborvitae	C,F
Thuja orientalis	Aureus nana-dwarf golden arborvitae	C,F
	Minima glauca-dwarf arborvitae	C,F
Veitchia merrilli	Christmas palm	F
Viburnum bodnantense	Pink dawn viburnum	C,F
Viburnum davidii	David viburnum	C,F
Viburnum japonicum	Viburnum	F
Viburnum lantana	Wayfaring tree	F
Viburnum opulus sterile	Common snowball viburnum	F
Viburnum plicatum tomentosum	Doublefile viburnum	C,F
Viburnum setigerum	Tea viburnum	F
Viburnum tinus compactum	Spring bouquet viburnum	F
Viburnum trilobum	Cranberry bush	C,F
Viburnum trilobum compactum	Dwarf cranberry bush	C,F
Viburnum x pragense	Viburnum	C,F
Weigela florida	Bristol ruby weigela	C,F
	Java red weigela	C,F
	Minuet weigela	C,F
Xylosma congestum	Xylosma	F
Yucca filamentosa	Yucca	C,F

Groundcovers

Recommended Treatment Method:

C = Container Grown

F = Field Grown

Scientific Name	Common Name	
Arctotheca calendula	Cape weed	F
Carex spp.	Variigated carex	C,F
Carpobrotus edulis	Ice plant, largeleaf (see label)	F
Cortaderia selloana	Pampas grass	F
Delosperma alba	White iceplant	F
Descampsia caespitosa	Descampsia	C,F
Drosanthemum floribundum	Trailing rosea iceplant	F
Drosantheumum hispidum	Iceplant	C,F
Gazania spp.	Gazania	C,F
Hedera canariensis	Ivy, Algerian	F
Hedera helix	Ivy, English	F
Hemerocallis spp.	Daylily	C,F
Hosta lancifolia	Albo-marginata hosta	C,F
Hypericum spp.	St. Johnswort	C,F
Jasminum nitidum	Angelwing jasmine	C,F

22/25

Lampranthus spectabilis	Trailing iceplant	F
Liriope gigantea	White lily turf	C,F
Liriope muscari	Lilac beauty lily turf	C,F
	Majestic lily turf	C,F
	Monroe white lily turf	C,F
	Silvery sunproof lily turf	C,F
	Variiegated liriope lily turf	C,F
	Big blue lily turf	C,F
Liriope spicata	Green/creeping lily turf	C,F
Lonicera japonica	Honeysuckle, Japanese	F
Ophiopogon japonicus	Dwarf mondo grass	F
	Mondo grass	F
Osteospermum fruticosum	Daisy, trailing African	F
Pachysandra terminalis	Japanese spurge	F
Trachelospermum asiaticum	Asian jasmine	C,F
Vinca major	Periwinkle, bigleaf	C,F
Vinca minor	Periwinkle, dwarf	F
Vinca spp.	Periwinkle	F

Non-bearing Fruit and Nut Trees and Non-bearing Vineyards† **Recommended Treatment Method:**
C = Container Grown
F = Field Grown

Common Name

Almond	F
Apple	F
Apricot	F
Avacado	F
Blackberry	F
Blueberry	F
Boysenberry	F
Cherry, sour	F
Cherry, sweet	F
Currant	F
Dewberry	F
Elderberry	F
Fig	F
Filbert	F
Gooseberry	F
Grape, American	F
Grape, European	F
Grapefruit	F
Kiwi	F
Lemon	F
Loganberry	F
Macadamia nut	F
Nectarine	F
Olive	F
Orange	F
Peach	F
Pear	F
Pecan	F
Pistachio	F
Plum	F

23/25

Pomegranate	F
Prune	F
Raspberry	F
Walnut, black	F
Walnut, English	F

† Non-bearing fruit and nut trees and non-bearing vineyards are defined as plants which will not bear fruit for at least one year after treatment.

Ornamental Bulbs

Do not use Snapshot 80 Dry Flowable in bulb production fields. For use in landscape settings only.

Snapshot 80 Dry Flowable may be applied for control of susceptible annual weeds in ornamental bulbs such as bulbous iris, daffodil (narcissus), gladiolus, hyacinth, and tulip. Apply Snapshot 80 Dry Flowable to the soil surface 2 to 4 weeks after planting, but prior to the emergence of annual weeds. Snapshot 80 Dry Flowable may also be applied following emergence but prior to and after flowering. For fall planted bulbs, apply Snapshot 80 Dry Flowable again in late winter or early spring to weed-free soil surfaces.

Special Use Precautions:

Do not apply to tulip plants that have emerged to a height greater than 3/4 inch.
 Do not apply to gladioli prior to emergence or to plants less than one inch in diameter.
 Do not apply to bulbous plants during flowering.

Application Rates

Time of Application	Soil Texture	Snapshot 80 Dry Flowable (oz/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (oz/1000 sq ft)
Fall	Coarse	0.46	3	0.92
Fall	Medium and Fine	0.92	3	1.38
Feb. - March	All Soil Textures	0.46	3	1.38

Shade House Areas

Snapshot 80 Dry Flowable may be applied in open shadehouse-type structures where the natural flow of air is unimpeded. Do not apply in enclosed greenhouses or in enclosed shadehouse-type structures. Do not apply within three weeks prior to enclosing greenhouse or poly-type structures.

Christmas Tree Plantations

Snapshot 80 Dry Flowable Alone

Apply Snapshot 80 Dry Flowable as a directed spray to the soil surface or as an overtop spray to established plantings of field grown Christmas tree species listed in this label. Do not apply to Douglas-fir (*Pseudotsuga menziesii*) or eastern hemlock (*Tsuga canadensis*). Do not apply to seedbeds or seedling transplant beds. Apply only to established plantings. Established plants are defined as those that have been transplanted into their final growing location for a sufficient period of time to allow the soil to be firmly settled around the roots from packing and rainfall or irrigation. Follow all instructions provided in the "General Information" section of this label.

Application Rates and Application Frequency

24/25

Snapshot 80 Dry Flowable (lb/acre)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (lb/acre)
2.5	2	13.3
3.75	3	13.3
5	3	13.3

Tank Mix Combinations

Tank mix combinations of Snapshot 80 Dry Flowable plus other labeled herbicides may be used as directed or otop sprays in established Christmas tree plantings. When applied according to use directions, these tank mixes will provide control of susceptible weed species listed on the respective product labels. Refer to tank mix product labels for specific use directions, precautions and limitations before use.

Snapshot 80 Dry Flowable Plus glyphosate: Apply tank mix combinations of Snapshot 80 Dry Flowable plus glyphosate only as directed sprays in Christmas tree plantings. When applied according to use directions, Snapshot 80 Dry Flowable plus glyphosate will provide postemergence control of susceptible weed species listed on the label for glyphosate and residual preemergence control of susceptible weed species listed on the label for Snapshot 80 Dry Flowable. Refer to the label for glyphosate for specific use directions, precautions and limitations before use.

Special Use Precautions:

Do not apply sprays containing glyphosate over the top of Christmas tree plantings.

Extreme care must be exercised to avoid contact of spray containing glyphosate with foliage and stems of Christmas trees or severe damage or death may result.

Noncropland Areas: Industrial Sites, Utility Substations, Highway Guardrails, Sign Posts, and Delineators

Snapshot 80 Dry Flowable is recommended as a preemergence treatment for control of certain annual grasses and broadleaf weeds in non-cropland areas, including industrial sites, utility substations, highway guardrails, sign posts and delineators. Apply Snapshot 80 Dry Flowable anytime prior to germination of target weeds. Areas to be treated should be free of established weeds or existing weeds should be controlled with postemergence herbicides. Refer to the general use precautions, mixing and application directions, and information on application rates and weeds controlled. Snapshot 80 Dry Flowable is compatible and may be tank mixed with other herbicides registered for use on noncropland. Applied as directed, tank mixes containing Snapshot 80 Dry Flowable will provide control of susceptible weed species listed on the respective labels. All directions, precautions and limitations on the respective product labels apply to the tank mix use.

Application Rates and Application Frequency

Snapshot 80 Dry Flowable (lb/acre)	Minimum Time Between Applications (months)	Total Amount Allowed Per Year (lb/acre)
2.5	2	10
3.75	4	13.75
5	4	15

26/25

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. **UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**

To the extent allowed by law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Snapshot is a Registered Trademark of United Phosphorus, Inc.

EPA-Last accepted 11/04/1999

Draft for RED - 11/7/05