

EPA Reg. Number:

Date of Issuance:

SEP 2 7 2007

70506-56

Term of Issuance:

Unconditional

Name of Pesticide Product:

XL 2GR

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

NOTICE OF PESTICIDE:

Registration X Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

United Phosphorus Inc.

423 Riverview Plaza

Trenton, NJ 08611

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you agree in writing to:

- 1. To the General Precautions and Restrictions section, change "When applied to turf, this product must be watered in as soon as possible after application to dissolve the granules" to "This product must be watered in immediately after application to dissolve the granules.".
- 2. Place the First Aid section on the front panel of the label after "Caution".
- 3. Remove "Wash thoroughly with soap and water after handling." from the Hazards to Humans and Domestic Animals statements.

- 4. To the Conditions of Sale and Limitation of Warranty and Liability section change "All such risks" to "To the extent consistent with applicable law, all such risks", "United Phosphorus, Inc. makes no" to "To the extent consistent with applicable law, United Phosphorus, Inc. makes no", and "the exclusive remedy" to "To the extent consistent with applicable law, the exclusive remedy".
- 5. To the Environmental Hazards section, change "This product is toxic to fish" to "This product is toxic to fish and aquatic invertebrates".

Signature of Approving Official:

Date:

SEP 2 7 2007

Joanne Miller, Team 23

Herbicide Branch, Registration Division (7505P)

EPA Form 8570-6

You will submit one copy of your final printed labeling before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). If you have any questions please call Erik Kraft at 703-308-9358.

1. Miller

with COMME. S In EPA Letter Dated:

Page 1

XL 2GR

Contains Surflan®

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

XL 2GR is an easy-to-use herbicide. XL 2GR stops unsightly weeds and weed grasses BEFORE they start to grow in your established ornamental shrubs, flower beds and warm-season lawn grasses around your home. Used according to label directions, XL 2GR provides effective weed control all season long — and gives your landscape that neat and clean look.

XL 2GR controls weeds in the following areas:

- · Landscape ornamental trees and shrubs
- Ornamental ground covers
- · Established flower beds
- Ornamental bulbs
- · Established warm season turfgrasses including:
 - bermudagrass
- buffalograss
- centipedegrass

- St. Augustinegrass
- zoysiagrass
- bahiagrass

• tall fescue

Active Ingredients:

, leave ingresions.	
benefin: N-butyl-N-ethyl- α , α , α -trifluoro-2,6-dinitro- ρ -toluidine	1.0%
oryzalin: 3,5-dinitro-N , N -dipropylsulfanilamide	1.0%
Inert Ingredients	98.0%
Total	

Contains __lb of active ingredient per __lb bag

Keep Out of Reach of Children

CAUTION

PRECAUCION

Si usted no entiende la etiqueta, busque a alquien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See back panel for additional precautionary statements.

Notice: Read the entire label. Use only according to label directions. Before using this product, read the Conditions of Sale and Limitation of Warranty and Liability elsewhere on this label. If terms are unacceptable, return at once unopened.

For emergency medical assistance, call the National Pesticide Information Center 1-800-858-7378. For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

<u>Agricultural Chemical: Do not ship or store with food, feeds, drugs, or clothing.</u>

Manufactured for: (logo) United Phosphorus, Inc. United Phosphorus, Inc. 423 Riverview Plaza Trenton, NJ 08611 1-800-247-1557 • www.upi-usa.com

EPA Reg. No. 70506-56 EPA Est. No.

Read all directions carefully before using this product. Herbicide

Net Weight Ib

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with labeling. Read all directions carefully before using this product.

GENERAL PRECAUTIONS AND RESTRICTIONS

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

- Do not apply this product by air.
- Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.
- Do not apply this product in a way that will contact any person or pets, either directly or through drift.
 Keep people and pets out of the area during application.
- When applied to turf, this product must be watered in as soon as possible after application to dissolve the granules. A single rainfall or irrigation of ½ inch or more is required after product application.
- Do not allow other people or pets to enter the treated area until dusts have settled. Except for those people involved in the watering-in, do not enter or allow other people or pets to enter the treated areas until the watering-in is complete and the grass is dry.
- Do not graze or feed forage from treated areas to livestock.

WHAT IS XL 2GR?

XL 2GR is a pre-emergence herbicide applied to the soil surface to prevent the growth of undesirable weeds and grasses. Simple for homeowner use, XL 2GR forms a protective "barrier" in the soil surface which prevent weeds from emerging.

WHAT WEEDS DOES XL 2GR CONTROL?

Used properly, XL 2GR controls the following broadleaf weeds and weed grasses:

Weed Grasses Broadleaf Weeds annual bluegrass bittercress carpetweed barnyardgrass brachiaria coast fiddleneck common chickweed browntop panicum common groundsel craborasses crowfootgrass common pursiane fall panicum desert rockpurslane Florida puslev foxtails field sandbur henbit lambsquarters goosegrass London rocket Italian ryegrass pigweeds johnsongrass (seedling only) iunalerice prostrate knotweed little barley prostrate spurge puncturevine little lovegrass redstem filaree Mexican lovegrass shepherdspurse orcutt lovegrass whitestem filaree red sprangletop

Page 3. 5/13

yellow woodsorrel (oxalis)

signalgrass

southwestern cupgrass

Texas panicum wild oats witchgrass

XL 2GR provides partial control of the following weeds, depending on soil and weather conditions:

Broadleaf Weeds annual sowthistle morningglory black mustard prickly lettuce

black mustard prickly lettuce black nightshade smartweed climbing milkweed spotted spurge

common mallow teaweed (prickly sida)

common ragweed velvetleaf volunteer wheat ladysthumb wild mustard

WHERE CAN I USE XL 2GR?

Apply XL 2GR over the top of ESTABLISHED shrubs, flowers, ground covers, bulbs, and warm season grasses. ESTABLISHED plants are those that have been transplanted into their final growing location and have the soil firmly settled around the roots from packing and rainfall or irrigation. To determine if it is permitted to apply XL 2GR over the top of any specific plant, refer to the "List of Tolerant Plants" located near the end of this label. In areas treated with XL 2GR, plant only ornamental species listed on the label for a period of one-year following application or injury may occur.

Note: If you will be transplanting plants into previously treated areas, untreated soil should be used to backfill such plants, or injury may occur.

Do not apply XL 2GR in ornamental plantings where the herbicide may runoff onto lawn areas containing dichondra or cool season turfgrasses. XL 2GR can cause severe injury or death in these plants.

Always read and follow the label directions when applying XL 2GR.

WHEN TO APPLY XL 2GR

XL 2GR stops weeds during germination. IT WILL NOT KILL ESTABLISHED WEEDS. Proper timing of application is important for best results. To be effective, XL 2GR must be applied before the weeds emerge.

In ornamental flower and shrub beds

For best results, apply XL 2GR as soon as possible after tilling, killing or pulling existing weeds from bed area. Since many weeds germinate in the spring, an application in early spring prior to the germination of annual broadleaf weeds and weed grasses is ideal (consult your extension agent or master gardener for the proper time to treat in your area). In warm climates with long growing seasons, a second application may be required 6 to 8 months after the first application to extend effective control.

Over the top of ornamental bulbs

Apply XL 2GR to the soil surface in the fail 2 to 4 weeks after planting of bulbs, but before annual weeds begin to sprout. Apply XL 2GR again in late winter or early spring. Before applying, remove any existing weeds from bed area. In the spring, do not apply to tulip plants that have grown to a height greater than 3/4". If you will be transplanting plants listed on this label into previously treated beds, use untreated soil to backfill such plants, or injury may occur. Otherwise, wait one year and cultivate deeply prior to planting other plants following use of XL 2GR.

In warm-season turfgrasses such as: tall fescue, bermudagrass, St. Augustinegrass, centipedegrass, zoysiagrass, buffalograss or bahiagrass

For maximum effectiveness, apply prior to the germination of broadleaf weeds and grass weeds (consult your extension agent for the proper time in your area). Apply in late winter to prevent emergence of spring germinating weeds, early spring to prevent summer annual weed grasses and broadleaf weeds, and late summer or early fall to prevent emergence of annual bluegrass and broadleaf winter annuals. If you miss the initial application period, partial control can still be achieved by using XL 2GR later than the optimal time since weeds and weed grasses continue to sprout throughout the growing season. Be sure to kill or pull all existing weeds prior to applying XL 2GR because XL 2GR will not control established weeds. In warm climates with long growing seasons, a second application may be required 4 to 6 months after the first application to extend control.

Turfgrass Precautions:

- XL 2GR should not be applied in the spring to turfgrass planted the previous fall.
- In bermudagrass areas that have been over-seeded with winter grasses, a spring application of XL 2GR may thin the overseeded grasses.
- XL 2GR should not be used on dichondra.
- XL 2GR should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfs are well established.
- Do not use on turf type tall fescue in spring and summer following fall seeding.
- Do not apply XL 2GR in late summer or early fall to areas where fall overseeding will occur.
- Wait at least six weeks after application of XL 2GR to reseed turfgrasses. When using XL 2GR at the highest rate, delay reseeding for 12 to 16 weeks following application.
- For satisfactory reseeding results, increase the amount of seed used and use equipment designed to
 place seed in contact with the soil.
- This product must be watered in as soon as possible after application to dissolve the granules. A single rainfall or irrigation of ½ inch or more is required after product application.

HOW TO APPLY XL 2GR

Read and follow all instructions for the application equipment. Always calibrate equipment according to the manufacturer's directions before applying.

On ornamental tree and shrub and flower beds

Before applying XL 2GR, make sure shrub beds are free of established weeds. Remove weed residues, prunings, and trash. Remove or break up large soil clods. Apply XL 2GR evenly over the shrub or bedding area using equipment designed to apply granular herbicides or insecticides (such as a rotary spreader). Read and follow all instructions for the application equipment. Always calibrate equipment according to the manufacturer's directions before applying. During application, check frequently to be sure equipment is working properly and distributing granules uniformly. For a more uniform application, spread one-half the desired amount over the area in one direction, then apply the remaining half at right angles to the first. For optimum weed control performance, thoroughly water in XL 2GR within a few hours after application to dissolve the granules.

On tall fescue, bermudagrass, St. Augustinegrass, centipedegrass, zoysiagrass, buffalograss and bahiagrass lawns

Apply XL 2GR to established turf before germination of target weeds since XL 2GR will not control established weeds. Apply XL 2GR evenly over the turfgrass area using a rotary spreader or other equipment used to apply granular herbicides or insecticides. Calibrate equipment according to the manufacturer's directions before applying. During application, check frequently to be sure equipment is working properly and distributing granules uniformly. For a more uniform application, spread one-half the desired amount over the area in one direction, then apply the remaining half at right angles to the first. For optimum weed control performance, thoroughly water in XL 2GR within a few hours after application

to dissolve the granules.

HOW MUCH XL 2GR SHOULD BE APPLIED?

Use the charts below to determine how much XL 2GR to apply. Over-application may result in plant injury or excessive soil residue.

Ornamental shrubs and beds:

Desired Length of Control	Rate of XL 2GR (lb/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed per Year (lb/1000 sq ft)
3 - 4 months	4 1/2**	2	9

^{**}Equivalent to 200 lb. XL 2GR per acre.

Ornamental bulb beds:

Time of Application	Soil Texture	Rate of XL 2GR (lb/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed per Year (lb/1000 sq ft)
Fall	Coarse	1 3/4**	3	3.4
Fall	Medium and Fine	3 1/2****	3	5.2
Feb March	All Soil Textures	1 3/4**	3	5.2

^{**}Equivalent to 75 lb. XL 2GR per acre

Turfgrass:

Type of weed	Time of Application	Amount of XL 2GR ^T
spring germinating	late winter	2 1/2 to 3 1/2 lb / 1000 sq ft
summer annual grass-weeds and broadleaf weeds	early spring	3 1/2 lb / 1000 sq ft
annual bluegrass (Poa annua) and broadleaf winter annuals	late summer/early fall	3 1/2 lb / 1000 sq ft

TXL 2GR may thin established annual bluegrass (Poa annua) turfs and fine-leafed fescues at rates above 2 1/2 lb/1000 sq ft

Rate of XL 2GR (lb/1000 sq ft)	Minimum Time Between Applications (months)	Total Amount Allowed per Year (lb/1000 sq ft)
2 1/2 - 3 1/2**	3 .	5 - 7

^{**}Equivalent to 100 - 150 lb XL 2GR per acre

Helpful Application Techniques for XL 2GR

On Ornamental Shrub Beds:

1. Handheld rotary applicators are ideal for broadcast applying XL 2GR. The rate of application depends on the

^{****}Equivalent to 150 lb XL 2GR per acre

gauge setting, walking speed of the operator and the speed of turning the crank. Walk and turn the crank at a constant rate of speed.

- 2. For hard to reach, small areas, a small container with holes punched in the bottom makes an excellent shaker type applicator.
- 3. On large areas, the push-type propelled rotary applicator does a fine job.
- Optimum weed control performance will be obtained if XL 2GR is watered in to the soil within a few hours after application.

On Turfgrass:

1. Small area application is easy using a small rotary applicator. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm is ideal.

Spreader Settings:

[Editor's note: Subregistrant may include spreader settings for recommended equipment.]

List of Tolerant Plants

Do not apply to Begonia, Coleus, Slender Deutzia, Douglas Fir, Techny Arborvitae and Eastern Hemlock as injury may occur.

The following ESTABLISHED plants are tolerant to applications of XL 2GR.

Established Flowers

Aster, Stokes
Baby's Breath
Bird Of Paradise
Blazing Star
Bleeding Heart

Caladium, Fancy Leafed

Chrysanthemum Coneflower, Purple

Coreopsis

Daisy, Gloriosa (Black-Eyed Susan)

Daisy, Painted Daisy, Shasta Geranium Geum Gladiolus

Impatiens (Busy Lizzie)

Iris, Bearded Marigold Marigold, Cape Moss, Rose Pansy Petunia

Ranunculus, Persian

Ranunculus, Peri Rose Sage Snapdragon Sweet William Yarrow Zinnia, Common

Established Ornamental Shrubs

Abelia, Glossy
Acacia, Prostrate
Agave
Andromeda
Arborvitae, American

Arborvitae, Aureus Nana-Dwarf Golden

Arborvitae, Emerald Arborvitae, Globosa-Globe

Arborvitae, Little Giant-Dwarf Arborvitae, Minima Glauca-Dwarf Arborvitae, Nigra-Dark American

Arborvitae, Oriental Arborvitae, Pyramidalis Arborvitae, Rheingold Arborvitae, Woodwardii

Azalea

Bamboo, Compacta-Dwarf Heaveniv Bamboo, Harbour Dwarf-Heaveniv

Bamboo, Heavenly

Bamboo, Nana Compacta-Heavenly Bamboo, Nana Purpurea-Heavenly Bamboo, Woods Dwarf-Heavenly Barberry, Atropurea-Redleaf Japanese Barberry, Aurea-Golden Japanese

Barberry, Crimson Pygmy Bottlebrush, Lemon

Bougainvillea, Barbara Karst Bougainvillea, California Gold Bougainvillea, Scarlet O'hara Bougainvillea, Texas Dawn

Boxwood, Common

Boxwood, Japonica-Japanese

Boxwood, Littleleaf

Bridal Wreath

Broom, Hollandia-Warminster

Broom, Lena-Scotch Cactus, Easter Cherry, Brush Cinquefoil

Clevera, Japanese Cotoneaster, Bearberry Cotoneaster, Brightbead Cotoneaster, Cranberry Cotoneaster, Himalayan Cotoneaster, Parney

Cotoneaster, Praecox-Early Cotoneaster: Pyrenees Cotoneaster, Rock Cotoneaster, Rockspray

Covotebush

Cranberry Bush, Compactum-Dwarf

Crape Myrtle

Cryptomeria, Japanese Cypress, Filifera-Thread Cypress, Kosteri

Cypress, Squarrosa Minima-Dwarf Moss

Deutzia, Nakiana-Dwarf

Dogwood, Flaviramea-Yellowtwig

Dogwood, Flowering Dogwood, Kousa

Dogwood, Sibirica-Siberian

Dwarf Burning Bush

Escallonia

Euonymus, Canadale Gold

Euonymus, Emerald'n Gold Euonymus, Evergreen Euonymus, Silver King Euonymus, Spreading Euonymus, Stringybark

Euonymus, Sunspot Euonymus, Winged

Fatshedera

Firethorn, Formosa Firethorn, Scarlet Flame Azalea Forsythia Gardenia Grape, Oregon Hawthorn, India Hibiscus, Chinese

Hibiscus, Ross Estey

Holly, Balkans Holly, Blue Boy Holly, Blue Girl Holly, Chinese

Holly, Compacta-Dwarf Japanese

Holly, Convexa Holly, Dwarf Chinese Holly, Ebony Magic Holly, English Holly, Gold Coast

Holly, Helleri-Heller's Japanese

Holly, Japanese

Holly, Nana-Dwarf Yaupon Holly, Nordica-Inkberry

Holly, Pendula-Weeping Yaupon

Holly, San Jose Holly, Yaupon Honeysuckle, Mexican Honeysuckle, Trumpet Honevsuckle, Winter Hopseed Bush

Jasmine, Chinese Star

Juniper

Leucothoe, Coast Leucothoe, Drooping

Lilac Lilac, Wild

Lily-Of-The-Valley

Lily-Of-The-Valley, Temple Bells Lily-Of-The-Valley, Valley Rose

Manzanita, Stanford

Mockorange

Myoporum, Prostrate

Myrtle, True Oleander

Oleander, Hardy Red Oleander, Ruby Lace Osmanthus, Holly-Leaf

Palm, Areca

Palm, Chinese Fountain

Palm, Christmas

Palm, Pigmy Date

Photinia, Fraser's

Pittosporum

Plant, Century

Plumbago, Blue Cape

Privet, Amur

Privet, California

Privet, Glossy

Privet, Howardi

Privet, Japanese

Privet, Vicary Golden

Privet, Wax Leaf

Protea

Pyracantha, Lolendei Monrovia

Pyracantha, Monon

Pyracantha, Rutgers Hybrid

Pyracantha, Santa Cruz

Pyracantha, Victory

Pyrcantha, Red Elf Hybrid

Rhaphiolepis, Enchantress-Moness

Rhaphiolepis, Roundleaf

Rhaphiolepis, Springtime-Monme

Rose Of Sharon, Red Bird

Rose Of Sharon, Red Heart

Rose Of Sharon, Woodbridge

Rose, Ramanas

Rosemary

Shrimp Plant

Snowdrift

Sotol, Desert Spoon

Spiraea, False

Spruce, Repens-Spreading Norway

Spruce, Repens-Weeping Norway

Spurge, Japanese

St. Johnswort

Sumac, African

Viburnum

Viburnum, Compactum-Spring Bouquet

Viburnum, David

Viburnum, Doublefile

Viburnum, Eastern Snowball

Viburnum, Laurustinus,

Viburnum, Sandankwa

Viburnum, Tea

Weigela, Bristol Ruby

Weigela, Java Red ...

Weigela, Minuet

Wintercreeper

Woadwaxen

Woodbine, Flowering

Woodbine, Serotina

Xylosma

Yew

Yew, Japanese

Yewpine

Yucca, Pendulous

Yucca, Soaptree,

Established Ground Covers

Bellflower

Blue fescue

Cape weed

Carpet bugle

Daisy, Trailing African

Gazania

Grass, Mondo

Grass, Pampas

Honeysuckle, Japanese

Ice plant

Ice plant, largeleaf

tceplant, Red Trailing

Iceplant, Trailing rosea

scepiant, White

Ivy, Algerian

Ivy, English

Jasmine, Chinese Star Periwinkle, Bigleaf

Lily Turf, Big blue

Lily Turf, Lilac Beauty

Lily Turf, Majestic

Lily Turf. Monroe white

Lily Turf, Silvery Sunproof

Lily Turf, Variegated liriope

Lily Turf, White

Lily, Day

Lily, plantain

Lily-of-the-Nile

Periwinkle, Bigleaf.

Periwinkle, Dwarf

St. Johnswort

Stonecrop

Trumpet Vine, Violet

Established Trees

Ash

Birch, Paper

Birch, River

Birch, White

Blue Palo Verde

California Pepper Tree

Carob

Carrotwood.

Cherry, Mahaleb

Cherry, Yoshino Flowering

Cottonwood

Crabapple

Cryptomeria, Japanese

Cypress, Arizona

Cypress, Filicoides-femspray

Cypress, Gracilis-slender Hinoki

Cypress, Italian

Cypress, Lawson false Cypress, Sawara-false Cypress, Squarrosa-moss

Dogwood, flowering

Elm, Chinese

Eucalyptus, Mealy

Eucalyptus, Narrow-leaved

Eucalyptus, Red gum

Eucalyptus, Red ironbark

Eucalyptus, Silver Dollar

Ficus

Fir, Alpine

Fir, Balsam

Fir, Fraser

Fir. Grand

Fir. Veitch

Fir, White

Goldenrain tree

Gum, American sweet

Laurel, California

Laurelcherry, Carolina

Laurelcherry, English

Linden, Littleleaf

Locust, Honey

Magnolia, Southern

Mahogany

Maidenhair tree

Maple, Albus Flowering

Maple, Luteus flowering

Maple, Roseus flowering

Maple, Tangerine floweringMaple, Vesuvius red

flowering

Maple

Mountain Laurel

Mulberry, White

Oak

Olive, Black

Olive, Russian

Olive

Paim, Queen

Palm, Cat

Palm

Paim, Parlor

Palm, Mexican fan

Pine

Podocarpus

Redbud

Redcedar, Eastern

Redcedar, Western

Redwood, Coast

Sequoia, Giant

Spruce, Norway

Spruce, Englemann

Spruce, Conica-dwarf Alberta

Spruce, White

Spruce, Black

Spruce, Glauca-Colorado blue Spruce, Hoopsil-Hoop's blue Spruce, Koster-Koster Blue Sycamore, American Sycamore, California Toyon Willow, Babylon weeping Willow, Corkscrew Yellow tab

Established Ornamental Bulbs

bulbous iris narcissus gladiolus daffodils tulips

Established Warm-Season Turfgrasses

tall fescue
centipedegrass
bahiagrass
bermudagrass
buffalograss
St Augustinegrass
zoysiagrass

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Causes Moderate Eye Irritation • Prolonged Or Frequently Repeated Skin Contact May Cause Allergic Reactions In Some Individuals

Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid

if in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information, contact the National Pesticide Hotline Number 1-800-858-7378.

For chemical emergency: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Storage and Disposal

Do not contaminate water, food or feed by storage and disposal.

Pesticide Storage: Store in original container out of the reach of children and pets. In case of leak or spill, contain material and dispose of waste.

Pesticide Disposal: If empty: Do not reuse this container. Place in trash or offer for recycling if

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. [Editor's Note: Supplemental distributor will insert own toll free number.]

IMPORTANT INFORMATION READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Surflan is a registered trademark of United Phosphorus, Inc.

Draft for RED 11/3/05 EPA-last accepted 7/23/03