

70506-59

06/23/2006

113



U.S. ENVIRONMENTAL PROTECTION AGENCY  
Office of Pesticide Programs  
Registration Division (7505P)  
Ariel Rios Building  
1200 Pennsylvania Ave., NW  
Washington, D.C. 20460

EPA Reg. Number:  
70506-59

Date of Issuance:  
JUN 23 2006

NOTICE OF PESTICIDE:  
 Registration  
 Reregistration  
(under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product:  
STORM HERBICIDE

Name and Address of Registrant (include ZIP Code):

UNITED PHOSPHORUS, INC.  
423 RIVERVIEW PLAZA  
TRENTON, NJ 08611

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the Reregistration Eligibility Document(s), EPA has reregistered the product listed above. Enclosed is a copy of your label stamped "Accepted". This action is taken under the authority of section 4 (g) (2) (c) of the Federal Insecticide, Fungicide, and Rodenticide Act, as amended.

Reregistration under this section does not eliminated the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

A stamped copy of your labeling is enclosed for your records.

Signature of Approving Official:

Joanne I. Miller  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

Date:

JUN 23 2006

2/13

# STORM Herbicide

**For use on peanuts, rice and soybeans**

**ACTIVE INGREDIENTS\*:**

Sodium salt of bentazon.....	29.2%
Sodium salt of acifluorfen.....	13.4%
INERT INGREDIENTS.....	57.4%
TOTAL.....	100.0%

\* Equivalent to 2.67 pounds of bentazon and 1.33 pounds of sodium acifluorfen per gallon

**KEEP OUT OF REACH OF CHILDREN  
DANGER/PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See inside booklet for complete **Precautionary Statements, Statements of Practical Treatment, Directions for Use, and Conditions of Sale and Warranty.**

**FOR CHEMICAL EMERGENCY:** Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300

**ACCEPTED**

JUN 23 2006

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

70506-59



United Phosphorus, Inc.

423 Riverview Plaza  
Trenton, NJ 08611  
1-800-247-1557  
[www.upi-usa.com](http://www.upi-usa.com)

EPA Reg. No. 70506-59  
EPA Est. No.  
Net Contents: 2.5 gal

Storm Draft Label for RED

June 19, 2006

Page 2 of 12

<b>FIRST AID</b>	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 – 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 – 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical assistance, call The National Pesticide Information Center at 1-800-858-7378.	
<b>Note to Physician:</b> Probable mucosal damage may contraindicate the use of gastric lavage. ANTIDOTE---No specific antidote is available. Treat symptomatically.	

**PRECAUTIONARY STATEMENTS****HAZARDS TO HUMANS AND DOMESTIC ANIMALS****DANGER**

Corrosive. Causes irreversible eye damage. Harmful if swallowed or absorbed through the skin. Do not get in eyes or on clothing. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**PERSONAL PROTECTION EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are made of any waterproof material. If you want more options, follow the instructions for category A on and EPA chemical-resistance category selection chart.

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves
- Shoes plus socks
- Goggles or face shield

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not re-use them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separate from other laundry.

**Engineering Controls Statement**

When handlers use closed systems, enclosed cabs, or cockpits in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark, except as specified on this label for application to rice. Do not contaminate water when disposing of equipment wash waters or rinsate. Do not apply when weather conditions favor drift from target area.

**PHYSICAL AND CHEMICAL HAZARDS**

This product is a reducing agent and should not be mixed or stored in close proximity to strong oxidizing agents.

**GROUND WATER ADVISORY**

Sodium acifluorfen and bentazon are known to leach through soil to groundwater under certain conditions a result of agricultural use. Use of this product in areas where soils are permeable (sandy or sandy/loamy soils) and water tables are shallow could result in contamination of groundwater. Use of irrigated water in such areas will increase the likelihood of groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other people, either directly or through drift. Only handlers wearing PPE may be in the treatment area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This pesticide is toxic to vascular plants and should be used strictly in accordance with the drift and run-off precautions on this label to minimize off-site exposures.

All applicable directions, restrictions, precautions and Conditions of Sale and Warranty are to be followed. This labeling must be in the user's possession during application.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 48 hours.

The following PPE is required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water.:

- Coveralls over long sleeved shirt and long pants.
- Chemical-resistant gloves made of any waterproof material
- Chemical-resistant footwear plus socks
- Chemical-resistant headgear if overhead exposure
- Protective eyewear

Notify workers of pesticide application by warning them orally and by posting warning signs at entrances to treated areas.

**I. GENERAL INFORMATION**

Storm herbicide is intended for selective postemergence control of certain broadleaf weeds in peanuts, rice, and soybeans. In addition, Storm may provide partial control of some grasses.

**Crop Tolerance**

Soybeans and peanuts are tolerant to Storm at the stages of growth listed. Leaf speckling, yellowing, bronzing, or burning may occur, but plants generally outgrow this condition with 10 days. New growth is normal and crop vigor is not reduced. Storm has no adverse effect on rice when used according to directions and may be used on first and second (ratoon) crops.

**Cleaning Spray Equipment**

Clean application equipment thoroughly by using a strong detergent or commercial spray cleaner according to the manufacturer's directions and then triple rinsing the equipment before and after applying this product.

**II. APPLICATION INSTRUCTIONS**

Apply 1.5 pints of Storm per acre as follows unless instructed differently in Section VI. Crop-Specific Information. Applications can be made to actively growing weeds as aerial or broadcast applications at the rates and growth stages listed. The most effective control will result from making postemergence applications of Storm early, when weeds are small. Early application to weeds results in improved weed control and makes thorough spray coverage easier to obtain. Delaying application permits weeds to exceed the maximum size stated and will prevent adequate control.

**Irrigation**

In irrigated areas, it may be necessary to irrigate before treatment to ensure active weed growth. Weeds growing under drought conditions usually are not adequately controlled.

**Spray Coverage**

Weeds must be thoroughly covered with spray. Always use an adequate volume of spray solution to ensure thorough coverage. Dense leaf canopies shelter smaller weeds and can prevent adequate spray coverage.

**Cultivation**

Do not cultivate within 5 days before or 7 days after applying Storm. Cultivating 7 days after treatment may help provide season-long control.

**Aerial Application Methods and Equipment**

**Water Volume:** Use 5-10 gallons of water per acre.

**Spray Pressure:** Use up to 40 psi when using flat fan nozzles and use 40-60 psi when using hollow cone nozzles.

**Application Equipment:** Use only diaphragm-type nozzles to produce cone or fan-spray spray patterns. Nozzles must be oriented to discharge straight back with the air stream (opposite the direction of travel of the aircraft) and not more than 20° downward. Nozzles must be positioned 6-10 feet above crop.

**Special Directions for Aerial Application**

- To obtain uniform coverage and to avoid drift hazards, consult the Spray Drift Management section below. Do not apply Storm by air if ornamentals or sensitive nontarget crops such as cotton, sugar beets, sunflowers, or okra are within 200 feet downwind.

**SPRAY DRIFT MANAGEMENT**

Use best practices to avoid drift to all other crops and non-target areas. Do not apply when conditions favor drift from target areas. The interaction of many equipment and weather-related factors determine the potential for spray drift. Avoiding spray drift at the application site is the responsibility of the applicator. The applicator must follow the most restrictive use precautions to avoid drift, including those found in this labeling as well as applicable state and local regulations and ordinances. A drift control agent may reduce drift, however, it may also decrease weed control.

**Requirements for ground applications:**

For ground applications, adjust nozzle height and droplet size with wind speed according to the following table:

Wind speed	Nozzle height	Droplet size for standard nozzles (ASAE standard 572)
Less than 10 mph	Up to 2 feet	medium or coarser
	2-4 feet	coarse or coarser
	4-6 feet	very coarse or coarser
10 to 15 mph	0-2 feet	coarse or coarser
	2-4 feet	very coarse or coarser
	4-6 feet	extremely coarse

Do not apply when the wind speed exceeds 15 miles per hour. Do not apply at a nozzle height of greater than 6 feet above the ground or crop canopy. Apply as a medium or coarser spray (ASAE standard 572).

**Requirements for aerial applications:**

For aerial applications, apply only when the wind speed is less than or equal to 15 miles per hour using a release height of no more than 10 feet above the ground or crop canopy. If the wind speed is less than 10 mph, apply as a medium or coarser spray (ASAE standard 572). If the wind speed is between 10 mph and 15 mph, apply as a coarse or coarser spray (ASAE standard 572). The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Do not make aerial applications into temperature inversions. When aerial applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

**Table 1 - Application Timing – Peanuts and Soybeans**

Weeds Controlled (including triazine and ALS-resistant biotypes)	Weed Growth Stages	
	Leaf Stage (up to)	Maximum Height <sup>a</sup>
Anoda, Spurred <sup>b</sup>	4	2"
Carpetweed	3" diam.	2"
Cocklebur <sup>c</sup>	6	6"
Copperleaf, Hophornbeam	4	4"
Crotalaria	6	6"

Croton, Tropic	2	<2"
,Wooly	2	<2"
Eclipta	6	<2"
Jimsonweed	6	6"
Ladysthumb	6	6"
Lambsquarters <sup>b</sup>	6	2"
Mallow, Venice	6	2"
Morningglories	4	2"
Mustard, Wild	6	4"
Nightshade, Black	6	2"
,Eastern Black	6	2"
Pigweed, Redroot	6	2"
,Smooth	6	3"
Ragweed, Common	6	3"
,Giant	4	6"
Redweed	4	3"
Sesbania, Hemp	4	6"
Sida, Prickly or Teaweed <sup>b</sup>	4	2"
Smartweed, Pennsylvania	6	6"
Starbur, Bristly	6	3"
Texasweed	3	3"
Velvetleaf <sup>b</sup>	4	2"
Waterhemp, Common	6	3"
,Tall	6	3"

<sup>a</sup> A second application of 1.5 pints of **Storm** herbicide per acre can be made for controlling subsequent weed flushes or escaped weeds before they reach the maximum weed size listed. Refer to **Table 4** for the maximum application rate per year.

<sup>b</sup> For regrowth or new germination, a follow-up application of Basagran herbicide may be necessary (refer to Basagran label).

<sup>c</sup> Do not treat earlier than the two true leaf stage. Do not count cotyledon leaves

**Ground Application Methods and Equipment (Broadcast)**

**Water Volume:** Use 10-20 gallons of spray solution per broadcast acre for optimal performance. Increase water volume up to 50 gallons if crop or weed foliage is dense.

**Spray Pressure:** Use a minimum of 40 psi (measured at the boom, not at the pump or in the line).

**Note:** When using the lower water volume (i.e. 10 gallons per acre) or when crop and weed foliage is dense, use a minimum of 60 psi for best results.

**Application Equipment**

Use standard high-pressure pesticide flat fan or hollow cone nozzles spaced up to 20" apart. Do not use flood, whirl chamber, or controlled droplet applicator (CDA) nozzles as erratic coverage can cause inconsistent weed control. Do not use selective application equipment such as recirculating sprayers or wiper applicators.

**III. ADDITIVES**

To achieve consistent weed control, one of the following additives is needed: ammonium sulfate, crop oil concentrate, nonionic surfactant, or urea ammonium nitrate. AMS (or UAN) should be used when velvetleaf is a target weed. Additives may cause some leaf burn, but new growth is normal and crop vigor is not reduced. The potential for leaf burn is increased when relative humidity and temperature are high. See **Table 2 - Additive Rate Per Acre** for additive rates and **Table 3 - Additive Options for Storm Tank Mixes**.

**Ammonium Sulfate (AMS)**

AMS is a dry, granular nitrogen-source fertilizer. Use only fine feed-grade or spray-grade AMS because inferior grades of AMS do not dissolve adequately and can plug spray nozzles. United Phosphorus, Inc. does not recommend applying AMS if applied in less than 10 gallons per acre because of potential problems with precipitation in reduced volumes. Use AMS only if it has been demonstrated to be successful in local experience.

**Nonionic Surfactant**

The standard label recommendation is 1-2 pints of an 80% active nonionic spray surfactant per 100 gallons of water.

**Oil Concentrate**

The oil concentrate must contain either a petroleum or vegetable oil base and must meet all of the following criteria:

- be nonphytotoxic,
- contain only EPA-exempt ingredients,
- provide good mixing quality in the compatibility test, and
- be successful in local experience.

The exact composition of suitable products will vary; however, vegetable and petroleum oil concentrates should contain emulsifiers to provide good mixing quality. Highly refined vegetable oils have proven more satisfactory than unrefined vegetable oils. For additional information, see **Compatibility Test for Mix Components**. Some oil concentrates cause excessive leaf burn. Refer to your supplier for information concerning successful local experience before purchasing any oil concentrate.

**Urea Ammonium Nitrate (UAN)**

Commonly referred to as 28%, 30%, or 32% nitrogen solution. UAN may be added in place of other spray additives to improve weed control. Because most nitrogen solutions are mildly corrosive to galvanized, mild steel, and brass spray equipment, rinse the entire spray system with water soon after. Do not use brass or aluminum nozzles when spraying UAN.

**Temperature and Relative Humidity Effects**

The following standard will help determine the optimum additive rate to use. If the temperature and relative humidity exceed 150 (e.g. temperature of 85°F plus 70% relative humidity = 155), use the lower additive rates.

**Table 2 – Additive Rate Per Acre**

Additive	Ground Application	Air Application
AMS	2.5 pounds	2.5 pounds
Oil Concentrate	1-2 pints	1 pint
UAN Solution	4-8 pints	4 pints
Nonionic Surfactant	1-2 pints per 100 gallons	1-2 pints per 100 gallons

**Table 3 – Additive Options for Storm Tank Mixes**

Additive Options	Nonionic Surfactant (1-2 pints per 100 gallons)	AMS (2.5 pounds) or UAN (4-8 pints per acre)	Crop Oil Concentrate (1-2 pints per acre)	Nonionic Surfactant (1-2 pints per 100 gallons) + AMS (1-2 pounds per acre) or UAN (2-4 pints per acre)	Crop Oil Concentrate (1 pint per acre) + AMS (1-2 pounds per acre) or UAN (2-4 pints per acre)
Option A	•				
Option B		•			
Option C			•		
Option D				•	
Option E					•

**IV. GENERAL MIXING INFORMATION**

Additives and/or other pesticides may be mixed in the spray with **Storm** using the information in this section.

**Tank Mix Partners/Components**

The following products may be tank mixed with **Storm** according to the specific tank mixing instructions in this label and respective product labels.

- Assure<sup>®</sup> II (quizalofop)
- Basagran<sup>®</sup> (bentazon)
- Classic<sup>®</sup> (chlorimuron ethyl)
- Concert<sup>®</sup> SP (thifensulfuron methyl+chlorimuron ethyl)
- Facet 75 DF<sup>®</sup> (quinclorac)
- FirstRate<sup>®</sup> (chloransulam-methyl)
- Frontier<sup>®</sup> 6.0 (dimethenamid)
- Fusilade<sup>®</sup> DX (fluazifop-p-butyl)
- Fusion<sup>®</sup> (fluazifop-p-butyl + fenoxaprop-p-ethyl)
- glyphosate
- Matador<sup>®</sup> (quizalofop)
- Pinnacle<sup>®</sup> (thifensulfuron methyl)
- Poast<sup>®</sup> (sethoxydim)
- Poast<sup>®</sup> HC (sethoxydim)

- Propanil
- Pursuit<sup>®</sup> (imazethapyr)
- Raptor<sup>®</sup> (imazamox)
- Reliance<sup>®</sup> STS (thifensulfuron methyl + chlorimuron ethyl)
- Resource<sup>®</sup> (flumiclorac)
- Scepter<sup>®</sup> (imazaquin)
- Select<sup>®</sup> (clethodim)
- Skirmish<sup>®</sup> (chlorimuron ethyl)
- Starfire<sup>®</sup> (paraquat)
- Synchrony<sup>®</sup> STS (thifensulfuron methyl + chlorimuron ethyl)
- 2,4-DB

See **Section VI. Crop-Specific Information** for more details. Read and follow the applicable **Restrictions and Limitations and Directions for Use** on all products involved in tank mixing. The most restrictive labeling applies to tank mixes. Separate applications should be made if all target weeds are not at the labeled growth stage for treatment at the same time.

Physical incompatibility, reduced weed control, or crop injury may result from mixing **Storm** with other pesticides (fungicides, herbicides, insecticides, or miticides), additives, or fertilizers. United Phosphorus, Inc. does not recommend using tank mixes other than those listed on United Phosphorus, Inc. labeling. Local agricultural authorities may be a source of information when using other than United Phosphorus, Inc. recommended tank mixes.

#### Compatibility Test for Mix Components

Before mixing additives and/or other pesticides, always perform a compatibility jar test. For 20 gallons per acre spray volume, use 3.3 cups (800 ml) of water. For other spray volumes, adjust rates accordingly. Only use water from the intended source at the source temperature.

Add components in the sequence indicated in the **Mixing Order** using 2 teaspoons for each pound or 1 teaspoon for each pint of recommended label rate per acre. Always cap the jar and invert 10 cycles between component additions.

When the components have all been added to the jar, let the solution stand for 15 minutes. Evaluate the solution for uniformity and stability. The spray solution should not have free oil on the surface, nor fine particles that precipitate to the bottom, nor thick (clabbered) texture. If the spray solution is not compatible, repeat the compatibility test with the addition of a suitable compatibility agent. If the solution is then compatible, use the compatibility agent as directed on its label. If the solution is still incompatible, do not mix the ingredients in the same tank.

#### Mixing Order

When mixing additives and/or other pesticides in a spray tank, add the products to be used in the following sequence.

1. **Water.** Begin by agitating a thoroughly clean sprayer tank three-quarters full of clean water.
2. **Agitation.** Maintain constant agitation throughout mixing and application.
3. **Products in PVA Bags.** Place any product contained in water-soluble PVA bags into the mixing tank. Wait until all water-soluble PVA bags have fully dissolved and the product is evenly mixed in the spray tank before continuing.
4. **Water dispersible products** (such as dry flowables, wettable powders, suspension concentrates, or suspo-emulsions). If an inductor is used, rinse it thoroughly after the component has been added.
5. **Water-soluble products** (such as **Storm** herbicide). If an inductor is used, rinse it thoroughly after the component has been added.
6. **Emulsifiable concentrates** (such as oil concentrate when applicable). If an inductor is used, rinse it thoroughly after the component has been added.
7. **Water-soluble additives** (such as AMS or UAN when applicable). If an inductor is used, rinse it thoroughly after the component has been added.
8. **Remaining quantity of water.** Maintain constant agitation during application.

#### V. RESTRICTIONS AND LIMITATIONS

- **Maximum Seasonal Use Rate:** Do not apply more than a total of **3 pints of Storm herbicide** per acre, per season for peanuts and soybeans. Do not apply more than a total of **1.5 pints of Storm** per acre, per season for rice. Refer to **Table 4** for the maximum rate per acre per application.
- Do not apply more than a total of **2.0 pounds of bentazon a.i.** (from all sources) per acre, per calendar year.
- Do not apply **sequential** applications of **Ultra Blazer** or **Storm** within **15 days** following the initial application of **Storm**.



- **Preharvest Interval (PHI):** Do not apply Storm within 50 days of soybean or rice harvest, or 75 days of peanut harvest.
- **Restricted Entry Interval (REI):** 48 hours
- **Crop Rotation Restriction:** Small grains must not be planted in fields treated with Storm for 40 days following treatment. All other rotated crops must not be planted in fields treated with Storm for 100 days following treatment. In case of crop failure, only peanuts, rice, or soybeans may be immediately replanted. Do not reapply Storm if the application will exceed the maximum rate allowed per acre per season.
- Do not allow livestock to graze on treated forage for soybeans or peanuts. Do not feed treated vines.
- **Stress:** Do not apply to weeds or crops under stress due to lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures, as unsatisfactory control may result.
- Do not apply Storm to crops that show injury (leaf phytotoxicity or plant stunting) produced by any other prior herbicide applications, because this injury may be enhanced or prolonged. In the Southeast, in-furrow treatments of insecticides/nematocides may predispose peanuts to injury from Storm.
- **Rainfast Period:** Rainfall or overhead irrigation within 4 hours after application may reduce the effectiveness of Storm.
- Do not apply through any type of irrigation system.

**Table 4 – Crop-Specific Restrictions and Limitations**

Crop	Minimum Time from Application to Harvest (PHI)	Maximum Rate Per Acre Per Application	Maximum Rate Per Acre Per Season	Livestock Grazing or Feeding	Aircraft Application
Peanuts	75 days	1.5 pints	3 pints	No	Yes
Rice	50 days	1.5 pints	1.5 pints	No	Yes
Soybeans	50 days	1.5 pints	3 pints	No	Yes

**VI. SPECIFIC CROP INFORMATION**

**PEANUTS**

Apply 1.5 pints of Storm herbicide per acre to peanuts preemergence at cracking stage (initiation of soil cracking, but before peanut emergence from the soil), or postemergence to peanuts to control susceptible weeds but no more than 75 days before harvest.

An additional 2 pints of Basagran herbicide may be applied per acre following an application of 3 pints of Storm per acre, per season, but no additional Ultra Blazer herbicide should be applied. An additional 3 pints of Basagran or 1 pint of Ultra Blazer may be applied following an application of 1.5 pints of Storm per acre per season.

**Crop-Specific Restrictions and Limitations:**

In-furrow treatments of insecticides/nematocides may predispose peanuts to injury from Storm.

**Peanut Tank Mixes**

Storm may be applied in a tank mix with one of the following herbicides:

<u>Tank Mix Partner</u>	<u>Additive Option</u>
Frontier® 6.0 .....	A or C
Starfire® .....	A
2,4-DB .....	A

Refer to Table 3 for the additive option appropriate for each tank mix.

**RICE**

Apply 1.5 pints of Storm per acre when rice is at the late tillering stage up to the early boot stage, which normally occurs in June or July. Rice must be past the 3-leaf stage.

Do not apply more than 1.5 pints of Basagran following an application of Storm.

Do not apply Ultra Blazer to rice treated with Storm.

Do not apply Storm to rice with ground equipment when field is flooded because splashing will wash Storm off weed leaf surfaces and result in ineffective control.

Do not use Storm on rice fields where the commercial cultivation of catfish or crayfish is practiced.

Do not use water containing residues of Storm from rice cultivation to irrigate crops other than soybeans or peanuts.

Do not apply more than one application of Storm per acre, per season.

**Rice Tank Mixes**

Storm may be applied in a tank mix with one of the following herbicides:

<u>Tank Mix Partner</u>	<u>Additive Option</u>
Basagran <sup>®</sup>	A
Facet <sup>®</sup> 75 DF	A
Propanil*	A

\* Do not apply this tank mix if Ultra Blazer has been previously applied. Refer to Table 3 for the additive option appropriate for each tank mix.

**Table 5 – Storm herbicide – Rice**  
**Application Rate and Timing Table for Drained or Flooded Fields**

Weeds Controlled*	Leaf Stage	1.5 Pints of Storm Per Acre	
		Maximum Weed Height in Drained Fields	Maximum Weed Height Above Water Level
Cocklebur	2-10	10"	6"
Dayflower	2-10	6"	5"
Ducksalad	2-4	2"	--
Gooseweed	4-6	4"	--
Sesbania, Hemp	**	**	4"
Morningglory species	up to 4	2"	1"
Redstem	up to 6	4"	3"
Redweed	4-6	6"	--
Smartweed	2-10	6"	5"
Spikerush	2-6	6"	--
Nutsedge, Yellow***	4-6	6"	5"

\* Add a nonionic surfactant at a rate (concentration) of 0.25% v/v (2 pints per 100 gallons of spray solution).  
 \*\* Effective control can be obtained at practically all heights provided Storm plus a nonionic surfactant is applied before the bloom (flowering).  
 \*\*\* Add oil concentrate at a rate (concentration) of 1.25% v/v (2 pints per 100 gallons of spray solution) instead of a nonionic surfactant. Partial control can be expected.

**SOYBEANS**

To ensure optimum spray coverage of weeds, apply Storm herbicide to small actively growing weeds. Refer to Section II, Application Instructions and Table 1 for more information.

**Sequential application information:** An additional 2 pints of Basagran herbicide may be applied following applications totaling 3 pints of Storm per acre, per season, but no additional Ultra Blazer herbicide should be applied. An additional 3 pints of Basagran or 1 pint of Ultra Blazer may be applied following an application of 1.5 pints of Storm per acre, per season.

**Soybean Tank Mixes**

Storm may be applied in a tank mix with one of the following herbicides:

<u>Tank Mix Partner</u>	<u>Additive Option</u>
Assure <sup>®</sup> II <sup>1</sup>	D or E
Basagran <sup>®</sup>	A, B, or C
Classic <sup>®</sup>	D
Concert <sup>®</sup> SP (up to 0.25 ounce)	D
FirstRate <sup>®</sup>	D
Frontier <sup>®</sup> 6.0	A, B, or C
Fusilade <sup>®</sup> DX <sup>1</sup>	D or E
Fusion <sup>®</sup> 1	D or E
Matador <sup>®</sup>	D or E
Pinnacle <sup>®</sup> (up to 0.25 ounce)	D
Poast <sup>®</sup> 1	E
Poast <sup>®</sup> HC <sup>1</sup>	E
Pursuit <sup>®</sup>	D
Raptor <sup>®</sup>	D
Reliance <sup>®</sup> STS SP <sup>2</sup> (up to 0.25 ounce)	D
Resource <sup>®</sup>	C
Glyphoste	8.5-17 pounds of AMS per 100 gallons
Scepter <sup>®</sup>	D

11/13

Select <sup>®</sup> 2 EC .....	E
Skirmish <sup>®</sup> .....	D
Synchrony <sup>®</sup> STS <sup>2</sup> (up to 0.5 ounce) .....	E

- <sup>1</sup> For best results if applying as part of a weed control program with **Storm**, follow these guidelines:
- If the partner is applied prior to the **Storm** application, wait 24 hours before applying **Storm**.
  - If the partner is applied following the **Storm** application, wait 7 days before applying.
- <sup>2</sup> When applying this tank mix to soybean varieties other than those designated as STS, do not add oil concentrate.

Refer to **Table 3** for the additive option appropriate for each tank mix.

<b>Crops</b> This product can be used on the following crops:
<b>Peanuts</b>
<b>Rice</b>
<b>Soybeans</b>
Read label for complete Restrictions and Limitations and Application Instructions.

Weeds Listed in this Label	
Common Name	Scientific Name
Anoda, Spurred	<i>Anoda cristata</i>
Carpetweed	<i>Mollugo verticillata</i>
Crotolaria	<i>Crotolaria spp.</i>
Cocklebur	<i>Xanthium strumarium</i>
Dayflower	<i>Commelina spp.</i>
Devilsclaw	<i>Proboscidea louisianica</i>
Ducksalad	<i>Heteranthera limosa</i>
Galinsoga	<i>Galinsoga spp.</i>
Gooseweed	<i>Sphenoclea zeylandica</i>
Jimsonweed	<i>Datura stramonium</i>
Ladysthumb	<i>Polygonum persicaria</i>
Lambsquarters, Common	<i>Chenopodium album</i>
Mallow, Venice	<i>Hibiscus trionum</i>
Morningglory, Common (tall)	<i>Ipomoea purpurea</i>
.Cypressvine	<i>Ipomoea quamoclit</i>
.Entireleaf	<i>Ipomoea hederacea</i>
.Ivy leaf	<i>Ipomoea hederacea</i>
.Palmleaf	<i>Ipomoea wrightii</i>
.Pitted	<i>Ipomoea lacunosa</i>
.Purple Moonflower	<i>Ipomoea muricata</i>
.Smallflower	<i>Jacquemontia tamnifolia</i>
Mustard, Wild	<i>Brassica kaber</i>
Nightshade, Black	<i>Solanum nigrum</i>
.Eastern Black	<i>Solanum ptycanthum</i>
Nutsedge, Yellow	<i>Cyperus esculentus</i>
Pigweed, Redroot	<i>Amaranthus retroflexus</i>
.Smooth	<i>Amaranthus hybridis</i>
Ragweed, Common	<i>Ambrosia artemisiifolia</i>
.Giant	<i>Ambrosia trifida</i>
Redweed	<i>Melochia corchorifolia</i>
Redstem	
Sesbania, Hemp	<i>Sesbania exaltata</i>
Sida, Prickly or Teaweed	<i>Sida spinosa</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Spikerush	<i>Eleocharis spp.</i>
Starbur, Bristly	<i>Acanthospermum hispidum</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Waterhemp, Common	<i>Amaranthus rudis</i>
.Tall	<i>Amaranthus tuberculatus</i>

12/13

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Do not store below 40°F or above 100°F. Store in a dry place away from heat or open flame.

**Pesticide Disposal:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mix, or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact the state agency responsible for pesticide regulation or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Disposal:**

- **Plastic Containers:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.
- **Bulk/Mini-bulk Containers:** Refillable/re-usable containers should be returned to the point of purchase for cleaning and refilling because the container must be thoroughly cleaned before refilling.

**Steps to be taken in case material is released or spilled:** Dike and contain the spill with inert materials (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal. Remove contaminated clothing, and wash affected skin areas with soap and water. Wash clothing before re-use. Keep the spill out of all sewers and open bodies of water.

**IMPORTANT INFORMATION  
READ BEFORE USING PRODUCT**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. The Directions for Use of this product reflect the opinion of experts based on field use and tests. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent allowed by law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Basagran, Facet, Frontier, Poast, Pursuit, Raptor, and Scepter are registered trademarks of BASF AG.

Ultra Blazer and Storm are trademarks of United Phosphorus, Inc.

Butoxone is a registered trademark of Nufarm Americas, Inc.

Butyrac is a registered trademark of Rhone-Poulenc Nederland B.V.

Assure, Classic, Concert, Pinnacle, Reliance, and Synchrony are registered trademarks and STS is a trademark of E.I.

Dupont de Nemours and Company.

FirstRate is a registered trademark of Dow AgroSciences LLC.

Matador and Skirmish are registered trademarks of FMC Corp.

Fusilade, Fusion and Starfire are registered trademarks of a Syngenta Group Company.

Resource and Select are registered trademarks of Valent USA Corp.

Draft per EPA review June 7, 2006