



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

July 14, 2021

Jeanette Covert  
Regional Regulatory Manager  
UPL NA Inc.  
630 Freedom Business Center, Ste. 402  
King of Prussia, PA 19406

Subject: Registration Review Label Mitigation for Bifenthrin  
Product Name: UP-STAR GC GRANULAR INSECTICIDE  
EPA Registration Number: 70506-74  
Application Date: 03/18/2021  
Decision Number: 572286

Dear Ms. Covert:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Bifenthrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2  
EPA Reg. No. 70506-74  
Decision No. 572286

If you have any questions about this letter, please contact Quinn Gavin by phone at 703-347-0325, or via email at [gavin.quinn@epa.gov](mailto:gavin.quinn@epa.gov).

Sincerely,

A handwritten signature in blue ink, appearing to read 'Linda Arrington', with a stylized flourish at the end.

Linda Arrington, Branch Chief  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

Enclosure

**RESTRICTED USE PESTICIDE****Toxic to Fish and Aquatic Organisms**

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by the certified applicator's certification.

<b>Bifenthrin</b>	<b>GROUP</b>	<b>3A</b>	<b>INSECTICIDE</b>
-------------------	--------------	-----------	--------------------

## UP-Star® GC Granular Insecticide

**For use to control ants (including imported fire ants), mole crickets, and other insect pests on lawns in landscaped areas and perimeters around residential, institutional, public, commercial, and industrial buildings, parks, recreational areas, golf courses, sod farms, and athletic fields.**

**DO NOT USE THIS PRODUCT ON GOLF COURSES AND SOD FARMS IN NASSAU COUNTY OR SUFFOLK COUNTY, NEW YORK.**

**For outdoor use only.**

Active Ingredient:	By Wt.
<b>Bifenthrin*</b> .....	0.2%
<b>Other Ingredients:</b> .....	99.8%
<b>TOTAL</b> .....	100.0%

\*Cis isomers 97% minimum, trans isomers 3% maximum

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See other panels for additional precautionary information or see back panel pamphlet for additional precautionary information and complete directions for use.

<b>FIRST AID</b>	
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>NOTE TO PHYSICIAN</b>	
This product is a pyrethroid. If large amounts have been ingested, milk, cream, and other digestible fats and oils may increase absorption and so should be avoided.	
<b>HOTLINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical assistance, call the Rocky Mountain Poison Control Center at 1-866-673-6671.	

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident,  
call CHEMTREC 1-800-424-9300.**

UPL NA Inc.  
630 Freedom Business Center  
King of Prussia, PA 19406  
1-800-438-6071

EPA Reg. No. 70506-74  
EPA Est. No.

Net Contents:

**ACCEPTED**

Jul 14, 2021

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 70506-74

**PRECAUTIONARY STATEMENTS**  
**Hazards to Humans and Domestic Animals**

**CAUTION**

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

**The following Personal Protective Equipment requirements apply to sod farm use only:**

**Personal Protective Equipment (PPE):**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations:**

**Users Should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to fish and aquatic invertebrates. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street back onto the treated area of the lawn or garden will help to prevent runoff to water bodies or drainage systems.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply more than 0.2 lb. active ingredient (a.i.) per acre per application on residential use sites (i.e., around private homes, apartment buildings, condominiums, non-agricultural outbuildings, non-commercial greenhouses, pre-schools or day care facilities). May be applied at up to 0.4 lb. a.i. per acre per application on non-residential use sites (i.e., around institutional, public, commercial or industrial buildings; parks; recreational areas or athletic fields).

Aerial application is prohibited.

Do not broadcast apply more than 0.4 lbs. a.i. (200 lbs. UP-Star GC Granular) per acre per year.

For applications at rates above 0.34 lbs. ai bifenthrin/A (170 lbs. product/A, users must water the treated area immediately after application.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

Apply this product directly to the lawn or garden area. Water treated area as directed on this label.

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours).

Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

For soil or foliar applications, do not apply by ground within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply or irrigate to the point of runoff.

Do not apply when the wind speed is greater than 15 mph.

Not for use on nurseries, in commercial greenhouses or on grass grown for seed.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### **Agricultural Use Requirements**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry to treated areas during the restricted entry interval (REI) of 12 hours.

PPI required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

Coveralls

Waterproof gloves

Shoes plus socks

### **Non-Agricultural Use Requirements**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

If watering-in of the granules is necessary, do not allow people or pets on treated areas until the grass or

soil is dry.

Do not touch treated surface until dry.

## STORAGE AND DISPOSAL

**Prohibitions:** Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use close tightly.

In case of spill, avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

**To Confine Spill:** Cover to prevent dispersal. Place damaged package in a holding container. Identify contents.

**Pesticide Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal: Nonrefillable container. Do not reuse or refill this container.**

Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## RESISTANCE

For resistance management, Up-Star GC Granular Insecticide contains a Group 3A insecticide. Any insect population may contain individuals naturally resistant to Up-Star GC Granular Insecticide and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of Up-Star GC Granular Insecticide or other Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pest.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
  - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
  - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
  - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
  - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they still may provide pest management benefits.
  - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.

- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for that specific site and pest problems in your area.

For further information or to report suspected resistance, contact UPL at 1-800-438-6071.

#### VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing (name of pyrethroid) onto fields where a maintained vegetative filter strip of at **least 25 feet** exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
  - For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
  - The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
  - Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
  - A functional terrace system is maintained on the area of application.
  - Water and sediment control basins for the area of application are functional and maintained.
  - The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. <https://www.regulations.gov/document?D=EPA-HQ-OPP-2008-0331-0175>

#### Buffer Zones - Ground Application

- Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

#### GRASS AND PERIMETER APPLICATIONS (Including Golf Courses and Sod Farms)

Grass Application: Broadcast UP-Star GC Granular Insecticide with application equipment which will ensure uniform coverage of the treatment area.

#### APPLICATION RATES

The application rates listed below are expected to provide excellent control of the target pests under normal conditions. However, based on the experience of the applicator, UP-Star GC Granular may be applied at up to 0.4 lb a.i./A to control each of the pests listed in this Table. For residential lawn use, do not apply more than 0.2 lbs. a.i. per acre per application.

Insect	Rate per acre	Instructions
--------	---------------	--------------

<p>Armyworms Cutworms Sod Webworm</p>	<p>50 lbs. product/A;  1.15 lbs product/1000 sq. ft.  (0.1 lbs a.i./A)</p>	<p><b>Armyworms, Cutworms, and Sod Webworms:</b> For optimum control, irrigate the treated area immediately following treatment with up to 0.1 inches of water to activate (release from the granule) the insecticide.</p>
<p>Annual Bluegrass Weevil (<i>Hyperodes</i>) (Adult) Billbugs (Adult) Black Turfgrass Ataenius (Adult) Leafhoppers Mealybugs</p>	<p>50-100 lbs. product/A;  1.15-2.3 lbs. product/1000 sq. ft.  (0.1-0.2 lbs a.i./A)</p>	<p><b>Annual Bluegrass Weevil (<i>Hyperodes</i>) Adults:</b> Time application to control adult weevils as they leave their over-wintering sites and move into grass areas. This movement usually begins when Forsythia is in full bloom and ends when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more specific information regarding application timing. <b>Billbug Adults:</b> Apply when adult billbugs are first seen during April and May. Degree day models have been developed to optimize application timing. Consult your State Cooperative Extension Service for information specific to your region. In temperate areas, spring applications made to control billbug adults will also control over-wintered chinch bugs. <b>Black Turfgrass Ataenius Adults:</b> Apply in May and July to control the first and second generation of black turfgrass ataenius adults, respectively. Time the May application so that it coincides with the full bloom stage of Vanhoutte spiraea (<i>Spiraea vanhouttei</i>) and horse chestnut (<i>Aesulus hippocastanum</i>). Time the July application so that it coincides with the blooming of Rose of Sharon (<i>Hibiscus syriacus</i>).</p>
<p>Ants Centipedes Chinch Bugs Fleas (Larvae) Fleas (Adult) Imported Fire Ants (Adult) Imported Fire Ants (Mounds) Millipedes Mole Cricket (Adult) Mole Cricket (Nymph) Ticks</p>	<p>100-200 lbs. product/A;  2.3-4.6 lbs. product/1000 sq. ft.  (0.2-0.4 lbs a.i./A)</p>	<p><b>Chinch Bugs:</b> Chinch bugs are found at the base of grass plants and also in the thatch layer. Immediately following application, irrigate the treated area with as much as 0.25 inches of water to activate (release from the granule) the insecticide. Since Chinch bugs can be one of the most difficult pests to control in grasses, the higher application rates may be necessary to control populations that contain both nymphs and adults during the summer. <b>Flea Larvae:</b> Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Immediately after application, irrigate the treated area with as much as 0.5 inches of water in order to activate (release from the granule) the insecticide. <b>Imported Fire Ants:</b> For best control combine broadcast applications (for control of foraging workers and newly mated fly-in queens) with mound treatments (for control of existing colonies). If the soil is dry, it should be irrigated before application. Use up to 0.4 lb a.i./A for broadcast treatments. Treat mounds by sprinkling 1/2 cup of UP-Star GC Granular on each mound and then drenching each mound with 1 to 2 gallons of water. Treat the mounds with enough force to break their apex and allow the insecticide solution to flow inside. Treat a four foot diameter circle around the mound. For optimum results, make application in cool weather (65-80° F) or in early morning or late evening hours. Treatment as described above will kill the queen within 24 hours of</p>



		<p>application. Note: a spray rig that is calibrated to apply 0.1 lb a.i./A (20 ounces) of BIFENTHRIN Flowable in 2.5 gallons per 1,000 sq. ft. contains the approximate dilution (1 teaspoon per gallon) that is required for fire ant mound drenches in the spray tank. Application of 0.25 lb/acre of UP-Star GC Granular Insecticide will provide six months residual activity for control of both foraging imported Fire Ants and newly mated fly-in queens.</p> <p><b>Mole Cricket Adults:</b> It is difficult to achieve acceptable control of adult mole crickets because they continuously invade preferred grass areas during the early spring. Apply as late in the day as possible and water in with up to 0.5 inches of water immediately following treatment. If the soil is dry, it is important to irrigate before application in order to bring the mole crickets closer to the soil surface where they will come in contact with insecticide. Grass areas infested by adult mole crickets should be treated at peak egg hatch to ensure optimum control of later nymph populations (see below).</p> <p><b>Mole Cricket Nymphs:</b> If grass areas received intense adult mole cricket pressure in the spring, they should be treated immediately before peak egg hatch. This is the time of best control because young nymphs are more susceptible to insecticides and because they are near the soil surface. Control of larger, more damaging, nymphs later in the year is difficult and may require both higher application rates and more frequent applications to maintain acceptable control. Apply as late in the days as possible and water in with as much as 0.5 inches of water immediately following treatment. If the soil is dry, it is important to irrigate before application to bring the mole crickets closer to the soil surface.</p> <p><b>Ticks (including ticks that may transmit Lyme Disease):</b> Do not make spot applications, but treat the entire area where exposure to ticks may occur. When treating areas of dense ground cover or heavy leaf litter, use higher application rates. Note that ticks may be reintroduced from surrounding areas by means of host animals. Do not allow public use of treated areas during application.</p> <p><b>Deer ticks (<i>Ixodes sp.</i>)</b> have a complicated life cycle that ranges over a two year period and involves four life stages. Apply in mid to late spring to control larvae and nymphs that live in the soil and leaf litter.</p> <p><b>American dog ticks</b> may be a problem in suburban settings, particularly where homes are built on land that was previously field or forest. Apply as necessary in areas where these ticks are found (paths, roadways) from mid spring to early fall to control American dog tick larvae, nymphs and adults.</p>
<p>Pillbugs Sowbugs European Crane Flies</p>	<p>200 lbs. product/A;  4.6 lbs. product/sq. ft.</p>	

	(0.4 lbs a.i./A)	
--	------------------	--

**In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).**

**In New York State, do make a single repeat application of UP-Star GC Granular if there are signs of renewed insect activity, but not sooner than two weeks after the first application.**

### Perimeter Application

Apply to the ground in an area 5 to 10 feet wide around and next to the structure.

Application Rate: Pounds Per 1,000 sq. ft.	Band Width	Pounds of UP-Star GC Granular for Perimeter Applications Linear Distances (Feet) Around Structure							
		100	150	200	250	300	400	500	1000
2.3	5 feet	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7
2.3	10 feet	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7
3.0	5 feet	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2
3.0	10 feet	3.9	5.4	6.9	8.5	9.9	12.9	15.9	30.9
3.5	5 feet	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5
3.5	10 feet	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5
4.6	5 feet	2.6	3.8	5.0	6.0	7.2	9.6	11.8	23.4
4.6	10 feet	6.0	8.4	10.6	13.0	15.2	19.8	24.4	47.4

Note: Use the following formula to determine the area to be treated: Multiply the linear distance by the band width and then add to their value the area of a circle whose radius is equal to the band width (to account for the area treated at the four corners of a rectangular structure).

### INDUSTRIAL NON-AGRICULTURAL APPLICATIONS

(Including Industrial Sites, Parks, Ball Fields and Cemeteries)

**Broadcast Spot Applications For the Control of Actively Foraging Ants (Including Imported Fire Ants):** Make a uniform broadcast application of 2.3 to 4.6 lbs UP-Star GC Granular Insecticide per 1000 sq. ft. (0.2 to 0.4 lbs a.i./Acre). See comments section under Grass Applications for additional rate recommendations.

**Application to Pad Mounted Electrical Sites to Reduce or Prevent Ant Mound Building Activity:** uniformly distribute 1.0 to 1.5 tablespoons (0.06 to 0.09 lbs.) of UP-Star GC Granular Insecticide per 4 sq. ft. of mounting pad. If existing transformers are completely or partially filled with soil which has been excavated by resident ants, first remove any excess fill and level the remaining soil before application is made. If food utensils such as tablespoons are used to measure pesticide, they must not be used for food purposes afterwards.

Note: Ants are more attracted to electrical transformers in the Fall or early Spring. For best results, make application in these cooler months.

### UP-STAR GC GRANULAR INSECTICIDE

#### Recommended Spreader Settings for Broadcast Applications

Equipment	Spreader Setting Ground Speed (MPH)	100 Pounds Per Acre		200 Pounds Per Acre	
		Once Over	Twice Over	Once Over	Twice Over
<b>Hand Pushed</b>					
LESCO Rotary	3	B + 1/4	A + 1/2	C + 1/2	B + 1/4
Scotts R-8A-1 Rotary	3	G	E	Not Rec.	G
Gandy Models (42, 36, 30, 24")	3	12	10.5	16	12
Spyker (All Models)	3	2.6	2	3.1	2.6

**Equipment Drawn (PTO=540 RPM)**

Grandy Models (42, 36, 30, 24")	4.5	14	11.5	19	14
Gandy 6500 Series	4.5	15	11	20	15
Gandy 10T Series	5.0	16.5	12	24	16.5
Herd GT-77	4.0	--	--	8-H	--
Herd GT-77	8.0	8-H	--	--	8-H
Lely 2.3202.000.1	4.5	2	1	3.5	2
Lely Model L-400	4.5	Not Rec.	Not Rec.	2	Not Rec.
Vicon PS402	4.5	10	7	16	10
Vicon PS1002T	4.5	13	9	21	13

**Note:** Use these recommended settings as starting points when calibrating a spreader to achieve the needed application rate of UP-Star GC Granular Insecticide. Once the equipment is calibrated, the technician should make further setting adjustments, as necessary, to maintain the proper application rate. Spreaders should be calibrated regularly to confirm that equipment wear or other factors have not changed the flow rate of UP-Star GC Granular Insecticide. Calibration instructions are included in the Owner's Manual that was provided by the equipment manufacturer at the time of purchase.

**Recommended Spreader Settings (Optional Text)**

*Note: Following will be a list of application spreaders and appropriate settings. This list will vary depending on the alternate carrier used. (This note will not appear on final printed label.)*

Note: These spreader settings were established using standard equipment available from the spreader manufacturer at swath widths and speeds typically used within the industry. UPI recommends that all spreader equipment be calibrated at the time of application to achieve the desired application rate.

**Dealers Should Sell In Original Packages Only**

**IMPORTANT INFORMATION  
READ BEFORE USING PRODUCT**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

Rev. 07/06/2021