

70506-152

7/14/2009

1 of 30



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUL 14 2009

Mr. Timothy M. Formella
United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

Subject: Revision of Storage and Disposal Statements and the Addition of Group Code

Dear Mr. Formella:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (s) (PRN) 2007-4 and 2001-5 dated June 19, 2009 for:

EPA Registration Number 70506-150	EPA Registration Number 70506-152
EPA Registration Number 70506-153	EPA Registration Number 70506-154

ImiGold 2F Turf, Ornamental and Termiticide/Insecticide	Aura 2F Insecticide
Imidacloprid 70 DF Agricultural Insecticide	Imidacloprid 70 DF Agricultural Insecticide
Fist 1.6 F	

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and 2001-5 find that the label change(s) requested falls within the scope of PRN 2007-4 and 2001-5. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, at 703 305-5409 or electronically at daniel.dani@epa.gov

Sincerely,

Dani Daniel
Registration Division (7505P)
Insecticide/Rodenticide Branch



United Phosphorus, Inc.

630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
(610) 491-2813 (phone)
(610) 491-2810 (fax)

June 19, 2009

VIA: United Parcel Service

Document Processing Desk (NOTIF)
Linda Arrington
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Room S-4900, One Potomac Yard
2777 South Crystal Drive
Arlington, VA 22202-4501

RE: Notification Regarding PR Notices 2007-4 and 2001-5

UPI Imidacloprid Technical Insecticide	EPA Reg. No. 70506-122
ImiGold 0.5 G Turf, Ornamental, and Termiticide/Insecticide	EPA Reg. No. 70506-142
ImiBloc 0.5 G Termiticide/Insecticide	EPA Reg. No. 70506-143
ImiGold 2 F Turf, Ornamental, and Termiticide/Insecticide	EPA Reg. No. 70506-150
Imidacloprid 2 F MUP Insecticide	EPA Reg. No. 70506-151
Aura 2 F Insecticide	EPA Reg. No. 70506-152
Imidacloprid 70 DF Agricultural Insecticide	EPA Reg. No. 70506-153
Fist 1.6 F Insecticide	EPA Reg. No. 70506-154
ImidaMax 4 F Insecticide	EPA Reg. No. 70506-155
ImiGold 70 DF Turf and Ornamental Insecticide	EPA Reg. No. 70506-156
ImiGold 70 DF Turf, Ornamental, and Greenhouse Insecticide	EPA Reg. No. 70506-157
ImiBloc 70 DF Termiticide/Insecticide	EPA Reg. No. 70506-158
ImiBloc 4 F Termiticide/Insecticide	EPA Reg. No. 70506-168

Dear Ms. Arrington:

United Phosphorus, Inc. hereby notifies the Agency of our intent to revise the labels for our registered pesticide products containing the active ingredient imidacloprid.

These label notifications include revision of the Storage and Disposal text in order to comply with the requirements of PR Notice 2007-4 regarding Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment." In addition, with the exception of the Technical and MUP product labels (EPA Reg. Nos. 70506-122 and 70506-151, respectively), the Insecticide Resistance Action Committee (IRAC) mode of action group code has been added to the first page of the labels per the Agency guidance in PR Notice 2001-5. The UPI warranty statements on these labels have been updated as well.


These changes are consistent with the notification provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46. Enclosed please find the following materials for each of the thirteen products to support these registration notifications:

FILE COPY

1. Completed Application for Pesticide Registration (EPA Form 8570-1)
2. A "red-lined" copy of the amended label showing the changes being made
3. A clean copy version of the revised label
4. CD with the labels in electronic format (file names: 070506-00122.20090619.pdf, 070506-00142.20090619.pdf, 070506-00143.20090619.pdf, 070506-00150.20090619.pdf, 070506-00151.20090619.pdf, 070506-00152.20090619.pdf, 070506-00153.20090619.pdf, 070506-00154.20090619.pdf, 070506-00155.20090619.pdf, 070506-00156.20090619.pdf, 070506-00157.20090619.pdf, 070506-00158.20090619.pdf, and 070506-00168.20090619.pdf)
5. Completed Certification with Respect to Label Integrity form

If you have any questions on this submission please call me at 610-491-2813 or email at tim.formella@uniphos.com.

Sincerely,


Timothy M. Formella
Senior Registration Manager

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirts and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

- When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.


ENVIRONMENTAL HAZARDS

Do not apply directly to water, areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This product is toxic to wildlife and highly toxic to aquatic invertebrates.

This chemical demonstrates the properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES; RESERVOIRS; RIVERS; PERMENENT STREAMS; MARSHES OR NATURAL PONDS; ESTUARIES, AND COMMERCIAL FISH FARM PONDS.



Resistance Management

Some insects are known to develop resistance to insecticides after repeated use. As with any insecticide, the use of this product should conform to resistance management strategies established for the use area.

Aura 2 F Insecticide contains a Group 4A Insecticide. Insecticide biotypes with acquired or inherent resistance to Group 4A may eventually dominate the insect population if Group 4A insecticides are used repeatedly as the predominant method of control for targeted species.

The active ingredient in Aura 2 F Insecticide belongs to the neonicotinoid chemical class. Insect pests resistant to other chemical classes have not shown cross-resistance to Aura 2 F Insecticide. In order to maintain susceptibility to this class of chemistry in insect species with high resistance development potential, it is recommended that for each crop season: 1) only a single soil application of Aura 2 F Insecticide be made; 2) foliar applications of products from this same class not be made following a long residual soil application of Aura 2 F Insecticide or other neonicotinoid products.

Other Group 4A neonicotinoid products used as foliar treatments include: Actara, Assail, Calypso, Centric, Gaucho, Intruder, Leverage, Provado, and Trimax.

Other Group 4A neonicotinoid products used as soil treatments include: Platinum.

Contact your local extension specialist, certified crop advisor and/or product manufacturer for additional insect resistance management recommendations. Also, for more information on Insect Resistance Management (IRM), visit the Insecticide Resistance Action Committee (IRAC) on the web at <http://www.irac-online.org/>.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.


Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treatment areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:



- Coveralls
- Chemical-resistant gloves made of any waterproof material such as, barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material is spilled for any reason or cause, absorb with sand or other inert material and dispose of absorbent in accordance with the Pesticide Disposal instructions listed below. Refer to Precautionary Statements on label for hazards associated with the handling of this material. In spill or leak incidents, keep unauthorized people away.

Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of the smoke.

Deleted: or

Deleted: To Confine Spills:

Deleted: carefully sweep material into a pile and dispose of as directed for pesticides below.

Deleted: may

Deleted: Disposal

Deleted: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of the smoke.


APPLICATION INFORMATION

For best control, direct applications of Aura 2 F Insecticide into the seed or root-zone of crop. Aura 2 F Insecticide may be applied with ground or chemigation application. Do not apply by air. Use only by broadcast or foliar applications on seedling flats or trays, or where product is meant to be washed from foliage to soil before drying on foliage.

For best results, apply Aura 2 F Insecticide to the root-zone of plants. Since Aura 2 F Insecticide is continuously taken into the roots over a long period of time, the earlier Aura 2 F Insecticide is available to a developing plant, the earlier the protection begins. The systemic nature of Aura 2 F Insecticide allows movement from roots through the xylem tissue to all vegetative parts of the plant, which results in extended residual activity of Aura 2 F Insecticide, the control of insects and the prevention and/or reduction of virus transmission or symptom expression, and plant health benefits. The rate of Aura 2 F Insecticide applied affects the length of the plant protection period; higher rates should be used when infestations occur later in crop development, or where pest pressure is continuous. Aura 2 F Insecticide will generally not control insects infesting flowers, blooms or fruit, and additional crop protection may be required for insects feeding in, or on these plant parts and for insects not listed in the crop-specific, pests controlled sections of this label.

Suppression (less than complete control of certain diseases and insect pests including reduced feeding) may also be caused by an Aura 2 F Insecticide application. In such cases, pests/diseases may require supplemental control measures.

Unless allowed by State specific supplemental labeling, the use of Aura 2 F Insecticide on crops grown for production of true seed intended for private or commercial planting is generally not



recommended. Always take care to minimize exposure of Aura 2 F Insecticide to honey bees and other pollinators. Additional information on Aura 2 F Insecticide uses for these crops and other questions may be obtained from the Cooperative Extension Service, PCAs, consultants or local United Phosphorus, Inc. representatives.

Mix Aura 2 F Insecticide with water or other appropriate diluent before application. Agitate the Aura 2 F Insecticide and water suspension to avoid settling.

Do not apply more than 0.50 lbs. active ingredient per acre, per crop season, regardless of formulation or method of application, unless specified within a crop-specific applications section for a given crop.

Mixing Instructions

To prepare the application mixture, first add a portion of the required amount of water to the tank and then add Aura 2 F Insecticide while agitation is underway. Add the remainder of water needed. Agitate during both mixing and application. Aura 2 F Insecticide may also be used with other pesticides and/or fertilizer solutions. Please see compatibility note below. When tank mixtures of Aura 2 F Insecticide and other pesticides are involved, prepare the tank mixture as recommended above and follow suggested Mixing Order below.

Mixing Order

When making pesticide mixtures, add Aura 2 F Insecticide and other wettable powders or wettable granules first, followed by flowable (suspension concentrate) products, then emulsifiable concentrates last. Agitate as each ingredient is added. Do not add an ingredient until the previous one is thoroughly mixed. If a fertilizer solution is added, a fertilizer/pesticide compatibility agent may be needed. To ensure a uniform spray mixture, continuous agitation is necessary during both mixing and application.

Compatibility Note

Test the compatibility of any intended mixture before adding Aura 2 F Insecticide to the spray or mix tank. To do this, add proportionate amounts of each ingredient in the appropriate order, to a suitable size jar, cap, shake the mixture for 5 minutes, and let set for 5 minutes. If the mixing is poor, or there is formation of precipitates that do not readily redisperse, then the blend is incompatible and should not be used. For further information, contact your local United Phosphorus, Inc. representative.

CHEMIGATION – DIRECTIONS FOR USE

Types of Irrigation Systems

Chemigation applications of Aura 2 F Insecticide may only be made to crops through chemigation systems as specified in crop-specific application sections and only through low-pressure systems unless specifically recommended for a given crop. Do not apply Aura 2 F Insecticide through any other type of irrigation system.

Uniform Water Distribution and System Calibration

The irrigation system must provide uniform distribution of treated water. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. The system must be calibrated to uniformly apply the rates specified. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Chemigation Monitoring

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Drift

Do not apply when wind speed favors drift beyond the area intended for treatment.

Required System Safety Devices

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Using Water From Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and to the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The pesticide injection pipeline must contain a functional automatic quick-closing check valve to prevent the flow of fluid back toward the injection. The pesticide injection pipeline must contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

ROTATIONAL CROPS*

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. For crops not listed on an imidacloprid label, or for crops with no established tolerances for the active ingredient, observe a 12 month plant-back interval.

Immediate Plant-Back All crops on this label plus the following crops not on this label:
barley, canola, corn (field, pop & sweet), rapeseed, sorghum,
soybean, sugarbeet and wheat

- 30-Day Plant-Back** Cereals (including buckwheat, millet, oats, rice, rye, and triticale), safflower
 - 10-Month Plant-Back** Onion and bulb vegetables
 - 12-Month Plant-Back** All other crops
- * Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.

FIELD CROPS

COTTON

PESTS	RATE	INSTRUCTIONS
Cotton Aphid Plant Bugs Thrips Whiteflies	1.3 fl oz/1000 row feet 17.0 – 21.1 fl oz/A (depending on row spacing)	Use one of the following methods to apply the specified dosage of Aura 2 F Insecticide: <ol style="list-style-type: none"> 1. In-furrow spray during planting directed on or below seed; 2. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; 3. Chemigation into root-zone through low-pressure drip or trickle irrigation. <p>Do not apply more than 21.1 fl oz (0.33 lb ai) Aura 2 F Insecticide per acre per crop season. No matter which formulation or method of application is used, do not apply more than 0.5 lb active ingredient of Aura 2 F Insecticide, Provado, Trimax or Leverage per acre per season, including seed treatment such as Gaucho, soil and foliar uses. Do not make more than 6 applications of the active ingredient per season regardless of method of application.</p> <p>Do not graze treated fields after any application of Aura 2 F Insecticide. Please note the Resistance Management section of this label.</p>

POTATOES

PESTS	RATE	INSTRUCTIONS
Controlled Aphids Colorado Potato Beetle Flea Beetles Leafhoppers Potato Psyllid	0.9 – 1.3 fl oz/1000 row feet 13.0 – 20.0 fl oz/A	Use one of the following methods to apply the specified dosage of Aura 2 F Insecticide : <ol style="list-style-type: none"> 1. In-furrow spray during planting, directed onto seed pieces or seed potatoes; 2. Subsurface side-dress on both sides of the row covered with 3 or more inches of soil; 3. Narrow band spray at ground cracking directly over the row during hilling covered with 3 or more inches of soil; 4. Narrow band directly below the eventual seed row in a bedding operation no more than 7 days before planting. Place Aura 2 F Insecticide applications below soil-surface and in contact with seed piece or within root-zone. For potatoes grown on highly permeable soils with a shallow water table, at-plant applications of Aura 2 F Insecticide may be made in 2 to 4 inch band (width of planter shoe opening) and completely covered.
Suppressed Symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis Wireworms (with in-furrow spray at planting)		

		Do not apply more than 20.0 fl oz (0.31 lb ai) Aura 2 F Insecticide per acre per crop season.
--	--	---

POTATOES

(Seed Piece Treatment)

This use not allowed in California unless otherwise directed by supplemental labeling.

PESTS	RATE	INSTRUCTIONS
Controlled Aphids Colorado potato beetle Flea beetles Leafhoppers Potato psyllid Wireworms (seed-piece protection)	0.4 – 0.8 fl oz/100 lbs seed 8.0 – 16.0 fl oz/A (based on a seeding rate of 2000 lbs/A)	Apply recommended rate as a diluted spray onto seed-pieces using a shielded spray system. Dilute with 3 parts water or less to 1 part Aura 2 F Insecticide. Agitate or stir spray solution as necessary. Apply fungicidal or inert absorbent dusts after Aura 2 F Insecticide is applied. Make applications only in areas with adequate ventilation or in areas that are equipped to remove spray mist or dust. Avoid extended exposure of seed-pieces to sunlight by planting as soon as possible after treating and in accordance with the recommendation of your local Extension specialist.
Suppressed Symptoms of: Potato leaf roll virus (PLRV) Potato yellows Net necrosis	0.8 fl oz/100 lbs seed 16.0 Fl oz/A (based on a seeding rate of 2000 lbs/A)	Consult your local United Phosphorus, Inc. representative or crop protection product dealer for information relevant to your area. Do not apply more than 20.0 fl oz (0.31 lb ai) Aura 2 F Insecticide per acre per crop season. Do not use treated seed-pieces for food, feed, or fodder. Do not apply any additional application of Aura 2 F Insecticide (in-furrow), Gaucho, Leverage, or Provado following an Aura 2 F Insecticide seed-piece treatment.

TOBACCO

PESTS	RATE	INSTRUCTIONS
Aphids Flea beetles	1.0 fl oz/1000 plants as seedling tray drench Or 1.4 fl oz/1000 plants (in-furrow transplant water)	Use one of the following methods to apply specified dosage of Aura 2 F Insecticide : 1. Uniform broadcast foliar spray to seedlings in trays (tray drench) no more than 7 days before transplanting, followed immediately by overhead irrigation to wash Aura 2 F Insecticide from foliage into potting media. Failure to wash the product from foliage may reduce pest control. 2. Handle transplants carefully during setting to avoid dislodging treated potting media from roots. 3. In-furrow spray or transplant-water drench during setting. 4. Chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
Mole crickets Whiteflies Wireworms	1.4 – 2.8 oz/1000 plants as seedling tray drench Or 1.8 – 2.8 fl oz/1000 plants (in-	Important Note: Proper tray drench applications of Aura 2 F Insecticide have been shown to be the most effective method of application. However, the specified rate of Aura 2 F Insecticide may be applied as combination of the tray drench in the planthouse and/or transplant-water drench in field. Adverse growing conditions may cause a delay in uptake of Aura 2 F Insecticide into the plant and a delay in control.

	furrow transplant water)	Pre-Harvest Interval (PHI): 14 days Do not apply more than 32.0 fl oz (0.50 lb ai) Aura 2 F Insecticide per acre per crop season.
Suppressed Cutworms Symptoms of: Tomato spotted wilt virus (TSWV)	1.4 – 2.8 oz/1000 plants as seedling tray drench Or 1.8 – 2.8 fl oz/1000 plants (in-furrow transplant water)	

VEGETABLE AND SMALL FRUIT CROPS

CUCURBIT VEGETABLES

Including: Chayote(fruit), Chinese waxgourd (Chinese preserving melon), Citron melon, Cuban pumpkin, Cucumber, Gherkin, Gourd (edible, includes hyotan, cucuzza, hechima, Chinese okra), *Momordica* (spp.), (includes balsam apple, balsam pear, bitter melon, Chinese cucumber), Muskmelon (hybrids and/or cultivars of *Cucumis melon* including true cantaloupe, cantaloupe, casaba, Crenshaw melon, golden pershaw melon, honeydew melon, honey balls, mango melon, Persian melon, pineapple melon, Santa Claus melon, snake melon, and Winter melon), Pumpkin, Squash, (includes summer squash types such as: butternut squash calabaza, crookneck squash, Hubbard squash, scallop squash, straightneck squash, vegetable marrow and zucchini, and winter squash types such as acorn squash, and spaghetti squash), Watermelon (includes hybrids and/or varieties of *Citrullus lanatus*).

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling.

Not for use in California unless specifically allowed by Supplemental labeling.

Field Applications

PESTS	RATE	INSTRUCTIONS
Control: Aphids Cucumber beetles Leafhoppers Thrips (foliage-feeding thrips only) Whiteflies Suppression: Bacterial wilt (as vectored by various cucumber beetles) Leaf silvering from whitefly feeding	16.0 – 24.0 Fl oz/A	Use one of the following methods to apply the specified dosage of Aura 2 F Insecticide: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. Incorporate product into

		<p>the root zone. Do not apply more than 24.0 fl oz (0.38 lb ai) Aura 2 F Insecticide per acre. Pre-Harvest Interval (PHI): 21 days</p>
--	--	--

Planthouse Applications

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	0.1 Fl oz/1000 plants	<p>Use one of the following methods to apply the specified dosage to seedlings in trays in the planthouse. Target soil media (tray drench), not more than 7 days prior to transplanting.</p> <ol style="list-style-type: none"> 1. Uniform, broadcast high volume foliar spray, followed by sufficient overhead irrigation to wash Aura 2 F Insecticide from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash product from foliage may reduce pest control; 2. Injection into overhead irrigation system, using sufficient volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray. <p>Applications made in the planthouse will only provide short-term protection and should not substitute for a field application. Make an additional field application within 2 weeks after transplanting to provide continuous protection. Applications of higher rates or increased number of application in planthouse may result in significant plant injury. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.</p> <p>Important Note: Not all varieties of cucurbit vegetables have been tested for tolerance to Aura 2 F Insecticide applied to seedling flats. It is therefore recommended to treat a small number of plants and confirm tolerance for 7 days before treating entire planthouse. Do not apply more than 0.1 fl oz (0.0056 lb ai)/1000 plants Aura 2 F Insecticide for planthouse applications. Do not apply more than once as a planthouse application.</p>

GREENHOUSE VEGETABLES

(Mature plants in production greenhouses)

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling.

Cucumber, Tomato, only

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	1.4 Fl oz/1000 plants	Apply specified dosage in at least 16 gallons of water for tomatoes and 21 gallons of water for cucumbers. Use soil drenches, micro-irrigation, drip irrigation, or hand-held or motorized calibrated irrigation equipment. Do

		<p>not apply to immature plants since phytotoxicity may occur.</p> <p>Apply when infestation pressure surpasses threshold and beneficials are not able to maintain pest populations below damage thresholds. Repellency of bumble bee pollinators and negative effects on some beneficials (<i>Orius</i> sp.) can occur when Aura 2 F Insecticide is applied.</p> <p>Many varieties of vegetables have been tested for tolerance to Aura 2 F Insecticide and show good safety. However, certain varieties may show more sensitivity to Aura 2 F Insecticide. Treatment of a few plants is recommended before treating a whole greenhouse.</p> <p>Pre-Harvest Interval (PHI): 0 days</p> <p>Do not apply Aura 2 F Insecticide more than once per season.</p>
--	--	---

FRUITING VEGETABLES

Including: Eggplant, Ground cherry, Okra, Pepper (including bell, chili, cooking, pimento and sweet), Tomato, Pepinos, Tomatillo

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling.

Field application

PESTS	RATE	INSTRUCTIONS
<p>Controlled:</p> <p>Aphids</p> <p>Colorado potato beetle</p> <p>Flea beetles</p> <p>Leafhoppers</p> <p>Thrips (foliage-feeding thrips, only)</p> <p>Whiteflies</p> <p>Suppressed:</p> <p>Symptoms of:</p> <p>Tomato mottle virus</p> <p>Tomato spotted wilt virus</p> <p>Tomato yellow leaf curl virus</p>	<p>Okra and Pepper</p> <p>16.0 – 32.0</p> <p>Fl oz/A</p> <p>Other Crops</p> <p>16.0 – 24.0</p> <p>Fl oz/A</p>	<p>Use one of the following methods to apply specified dosage of Aura 2 F Insecticide:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. Aura 2 F Insecticide must be incorporated into root-zone. <p>Pre-Harvest Interval (PHI): 21 days</p> <p>Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per application to pepper and okra;</p> <p>Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per application to other fruiting vegetables.</p>

Planthouse Application

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	0.1 Fl oz/1000 plants	<p>Use one of the following methods to apply the specified dosage to seedlings in trays in the planthouse. Target soil media (tray drench), not more than 7 days prior to transplanting.</p> <ol style="list-style-type: none"> 1. Uniform, broadcast high-volume foliar spray, followed immediately by sufficient overhead irrigation to wash Aura 2 F Insecticide from foliage into potting media without loss of gravitational liquid from the bottom of the tray. Failure to wash the product from foliage may result in reduced pest control. 2. Injection into overhead irrigation system, using adequate volume to thoroughly saturate soil media without loss of gravitational solution from the bottom of the tray. <p>Applications made in the planthouse will only provide short-term protection and should not substitute for a field application. Make an additional field application within 2 weeks after transplanting to provide continuous protection. Applications of higher rates or increased number of application in planthouse may result in significant plant injury. Handle transplants carefully during setting to avoid dislodging treated potting media from roots.</p> <p>Important Note: Not all varieties of fruiting vegetables have been tested for tolerance to Aura 2 F Insecticide applied to seedling flats. It is therefore recommended to treat a small number of plants and confirm tolerance for 7 days prior to treating entire planthouse. Do not apply more than 0.1 fl oz (0.0056 lb ai)/1000 plants Aura 2 F Insecticide for planthouse applications. Do not apply more than once as a planthouse application.</p>

HEAD AND STEM BRASSICA VEGETABLES

Including: Broccoli, Broccoli raab (rapini), Brussel sprouts, Cabbage, Cauliflower, Cavalo broccoli, Chinese (gai lon) broccoli, Chinese (bok choy) cabbage, Chinese (napa) cabbage, Chinese mustard (gai choy) cabbage, Collards, Kale, Kohlrabi, Mizuna, Mustard greens, Mustard Spinach, Rape greens, Turnip tops (leaves)

LEAFY VEGETABLES

Including: Amaranth (leafy amaranth, Chinese spinach, tampala), Arugula (Roquette), Chervil, Chrysanthemum (edible leaved and garland), Cilantro, Corn salad, Cress (garden), Cress (upland, yellow rocket, winter cress), Dandelion, dock (sorrel), Endive (escarole), Lettuce (head and leaf), Orach, Parsley, Purslane (garden and winter), Radicchio (red chicory), Spinach (including New Zealand and vine (Malabar spinach, Indian spinach)), Watercress (commercial production only, application must not be made to native cress growing in streams or other bodies of water), Watercress (upland).

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	10.0 – 24.0 Fl oz/36 inch rows	Use one of the following methods to apply the specified amount of Aura 2 F Insecticide: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting. 5. Post-seeding drench, transplant-water drench, or hill drench; 6. Subsurface side-dress on both sides of each row. Aura 2 F Insecticide must be incorporated into root-zone. Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per application.

LEAFY PETIOLE VEGETABLES

Including: Cardoon, Celery, Celtuce, Chinese celery (fresh leaves and stalk only), Florence fennel (including sweet anise, sweet fennel, Finocchio), Rhubarb, Swiss Chard.

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling

PESTS	RATE	INSTRUCTIONS
Aphids Leafhoppers Whiteflies	10.0 – 24.0 Fl oz/A	Apply specified dosage of Aura 2 F Insecticide using one of the following methods: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler, or equivalent equipment; 2. In-furrow spray directed on or below seed; 3. Narrow (2" or less) surface band spray over seed-line during planting incorporated to a depth of 1 to 1 ½" with sufficient irrigation within 24 hours of application; 4. Narrow band spray directly below eventual seed row in bedding operation 14 or fewer days before planting; 5. Post-seeding drench, transplant-water drench, or hill drench. 6. Subsurface side-dress on both sides of each row. Aura 2 F Insecticide must be incorporated into root-zone. Pre-Harvest Interval (PHI): 45 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per application.

LEGUME VEGETABLES except soybean, dry

Edible Podded and Succulent Shelled Pea and Bean and Dried Shelled Pea and Bean

Bean (*Lupinus* spp., includes grain lupin, sweet lupin, white lupin, and white sweet lupin)

Bean (*Phaseolus* spp., includes field bean, kidney bean, lima bean, navy bean, pinto bean, runner bean, snap bean, tepary bean, wax bean)

Bean (*Vigna* spp., includes adzuki bean, asparagus bean, blackeyed pea, catjang, Chinese longbean, cowpeas, Crowder pea, moth bean, mung bean, rice bean, southern pea, urd bean, yardlong bean)

Pea (*Pisum* spp., includes dwarf pea, edible-pod pea, English pea, field pea, garden pea, green pea, snow pea, sugar snap pea)

Other Beans and Peas (Broad Bean (fava), Chickpea (garbonzo bean), Guar, Jackbean, Lablab bean (hyacinth bean), Lentil, Pigeon pea, Soybean (immature seed), Sword bean)

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Leafhoppers Thrips (foliage-feeding thrips, only) Whiteflies Suppressed: Symptoms of: Bean common mosaic virus (BCMV) Bean golden mosaic virus (BGMV) Beet curly top hybrigeminivirus (BCTV)	16.0 – 24.0 Fl oz/A	Apply specified dosage of Aura 2 F Insecticide using one of the following methods: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray at planting directed on or below seed; 3. In a narrow (2" or less) surface band over seed-line during planting incorporated to a depth of 1 to 1 1/2" with sufficient irrigation with 24 hours following application; 4. In a narrow band directly below the eventual seed row in a bedding operation 7 or fewer days before planting; 5. As a post-seeding drench, transplant drench, or hill drench. Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per crop season.

ROOT VEGETABLES

Including: (Beet (garden*), Burdock (edible*), Carrot*, Celeriac*, Chervil (turnip-rooted*), Chicory*, Ginseng, Horseradish, Parsley (turnip-rooted), Parsnip*, Radish*, Oriental Radish (diakon*), Rutabaga*), Salsify (oyster plant), Salsify (black*), Salsify (Spanish*), Skirret and Turnip*.

* Tops and greens from these crops may be used for food or feed.

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling

PESTS	RATE	INSTRUCTIONS
Aphids Flea beetles Leafhoppers Whiteflies	0.7 – 1.7 Fl oz/1000 row feet 10.0 – 24.0 Fl oz/A	Apply specified dosage of Aura 2 F Insecticide using one of the following methods: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. In-furrow spray (rate specified per 1000 row-feet) or, shanked-in 1 to 2 inches below seed depth during planting; 3. In a narrow (2 inches or less) band directly (1 to 2 inches) below the eventual seed row in a

		<p>bedding operation 14 or fewer days before planting.</p> <p>Important Note: The rate used affects the length of control. Higher rates should be used where infestations occur later in crop development, or where pest pressure is continuous. Rates less than 0.7 fluid ounces/1000 row-feet will not provide adequate residual pest control. Aura 2 F Insecticide treated crops grown on very high organic matter soils (muck) may also require additional pest management control. Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per crop season. Do not apply more than once per crop season.</p>
--	--	---

TUBEROUS AND CORM VEGETABLES

Including: Arracacha, Arrowroot, Artichoke (Chinese and Jerusalem), Canna (edible, Queensland arrowroot), Cassava (bitter and sweet*), Chayote (root), Chufa, Dasheen (taro*), Ginger, Leren, Sweetpotato, Tanier (cocoyam*), Tumeric, Yam bean (jicama, manioc pea), Yam (true*)
(For recommended applications on potato see Field Crops section)

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling

* Tops and greens from these crops may be used for food or feed.

PESTS	RATE	INSTRUCTIONS
<p>Aphids Flea beetles Leafhoppers Whiteflies</p>	<p>0.7 – 1.7 Fl oz/1000 row feet 10.0 – 24.0 Fl oz/A</p>	<p>Use one of the following methods to apply specified dosage of Aura 2 F Insecticide:</p> <ol style="list-style-type: none"> 1. In-furrow spray (rate specified per 1000 row-feet) over planting material (hulis) or shanked-in 1 to 2 inches below hulis depth at planting; 2. Side-dress not more than 0.6 fluid ounces/1000 row-feet no later than 45 days after planting. Observe same PHI as above. <p>Important note: The rate applied affects the length of control. Use higher rates where infestations occur later in crop development, or where pest pressure is continuous. Rates less than 0.7 fluid ounces/1000 row-feet may not provide adequate residual pest control. Aura 2 F Insecticide treated crops grown on very high organic matter soils (muck) may also require additional pest management control. Pre-Harvest Interval (PHI) from planting application: 3 days (leaves); 125 days (corms). Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per crop season. Do not apply Aura 2 F Insecticide more than once per crop season.</p>

STRAWBERRIES

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling.

Do not use both application methods on the same crop in the same season.

Annual and Perennial Strawberries

PESTS	RATE	INSTRUCTIONS
Aphids Whiteflies	24.0 – 32.0 Fl oz/A	<p>Use one of the following methods to apply specified dosage of Aura 2 F Insecticide:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler, or equivalent equipment after plants are established or on perennial crops in early spring prior to bud opening; 2. As a plant material or plant hole treatment just prior to, or during transplanting. <p>The application rate affects the length of control. Use higher rates where infestations may occur later in crop development or where pest pressure is continuous. Pre-Harvest Interval (PHI): 14 days. Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per crop season.</p>

Post-Harvest Use on Perennial Strawberries

PESTS	RATE	INSTRUCTIONS
White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle, Oriental beetle)	16.0 – 24.0 Fl oz/A	<p>Make a single application post harvest to coincide with renovation of strawberry fields and during active egg-laying period of beetles. Apply recommended rate of Aura 2 F Insecticide using one of the following methods:</p> <ol style="list-style-type: none"> 1. As a ground spray by means of boom or backpack sprayer in a minimum of 20 gallons of water per acre; 2. As a row-band spray using an adjusted amount of product based on the treated row band area in proportion to the amount required per full acre. The bandwidth should be equivalent to the width of the anticipated fruiting bed; 3. As a chemigation application with 600 to 1000 gallons of water followed by 0.10 to 0.25 inches irrigation. <p>Important note: All soil-surface applications must be followed by either 0.25 inches of rainfall or overhead irrigation per acre within 2 hours of application in order to adequately incorporate Aura 2 F Insecticide into egg-deposition zone. Pre-Harvest Interval (PHI): 14 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per season.</p>

SUGARBEETS

(for use only in California)

Not for use on crops grown for seed unless specifically allowed by Supplemental labeling.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Leafhoppers Whiteflies Flea beetles Suppressed: Symptoms of: Western yellows/Beet curly top hybridgeminiavirus (BCTV)	6.0-12.0 Fl oz/A	Applied recommended rate of Aura 2 F Insecticide in sufficient carrier for uniform application, directly below each seed furrow. Apply either during the bedding operation immediately prior to planting or at the time of planting. The low rate may be used to aid establishment of stands in whitefly areas, or for early season control of the other pests listed. Do not apply more than 12.0 fl oz (0.18 lb ai)/A Aura 2 F Insecticide per crop season. Do not apply more than a total of 0.18 lb active ingredient (any formulation) on any row spacing per season.

AURA 2 F INSECTICIDE CONVERSION CHART FOR LINEAR APPLICATION								
RATE Fluid Ounces/Acre	RATE fluid ounces/1000 row-feet Based on average row spacing (in inches)							
	10	15	20	25	30	35	40	45
10	0.19	0.29	0.38	0.48	0.57	0.67	0.76	0.86
12	0.23	0.34	0.46	0.57	0.69	0.80	0.92	1.03
14	0.27	0.40	0.54	0.67	0.80	0.94	1.07	1.21
16	0.31	0.46	0.61	0.77	0.92	1.07	1.22	1.38
18	0.34	0.52	0.69	0.86	1.03	1.21	1.38	1.55
20	0.38	0.57	0.76	0.96	1.15	1.34	1.53	1.72
22	0.42	0.63	0.84	1.05	1.26	1.47	1.68	1.89
24	0.46	0.69	0.92	1.15	1.38	1.61	1.84	2.07
26	0.50	0.75	0.99	1.24	1.49	1.74	1.99	2.24
28	0.54	0.80	1.07	1.34	1.61	1.87	2.14	2.41
30	0.57	0.86	1.15	1.43	1.72	2.01	2.29	2.58
32	0.61	0.92	1.22	1.52	1.84	2.14	2.45	2.75

Important Note: The application rate affects the length of control and to a considerable extent, the degree of control or effect. Row-spacing X Aura 2 F Insecticide rate combinations in shaded blocks may not proved adequate residual pest control and are not recommended for long-term, residual control. Use higher labeled rates where infestations may occur later in crop development or where pest pressure is continuous. United Phosphorus, Inc. offers no warranty for use of Aura 2 F Insecticide at rates below .07 fluid ounces/1000 row-feet.

TREE, BUSH, AND VINE CROPS

BUSHBERRIES

Including: Blueberry, Currant, Elderberry, Gooseberry, Huckleberry, Juneberry, Lingonberry, Salal

PESTS	RATE	INSTRUCTIONS
<p>Japanese beetle (adults, feeding on foliage) White grub complex (grubs of Asiatic garden beetle, European and Masked chafer, Japanese beetle and Oriental beetle)</p>	<p>16.0 – 32.0 Fl oz/A</p>	<p>Use one of the following methods to apply specified dosage of Aura 2 F Insecticide:</p> <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. 18-inch band on each side of the row followed with 0.25 inches of irrigation immediately after application. <p>Apply Aura 2 F Insecticide to moist soil. If necessary, apply one hour of irrigation water immediately before application of Aura 2 F Insecticide. For best results, ½ to 1 inch of irrigation water or rainfall should be applied or received within 24 hours of application to facilitate movement into the soil and into the root-zone.</p> <p><u>Grubs:</u> For best grub control, apply Aura 2 F Insecticide to control 1st or 2nd instar larvae. Make application post-bloom up to 7 days before harvest, or post-harvest until October 1st.</p> <p><u>Japanese beetle larvae:</u> apply from June 1 to July 15. Do not apply during bloom.</p> <p>Apply to grass covered rows, row middles, drive lanes, headlands, and other grassy areas in and around the berry field to control resident grub populations. Applications directed to the root-zone will help protect berry plant roots from grub feeding. Pre-Harvest Interval (PHI): 7 days Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per season.</p>

CITRUS (Containerized)

Including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote spp. (*Casimiroa* spp), and other cultivars and/or hybrids of these.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Whiteflies	0.75 mL/ft ³ container media	Determine volume of container and calculate dosage necessary to treat container. Apply calculated dosage of Aura 2 F Insecticide per container as a soil drench or through low-pressure drip or trickle irrigation water. Use sufficient carrier volume to ensure thorough uniform distribution throughout the media without loss of gravitational water from the container. For best results, make treatment at planting before insect infestation occurs.
Citrus Root weevil (larval complex)	1.25 – 2.50 mL/ft ³ container media	Retreat if necessary. <u>Citrus root weevil complex (larvae):</u> apply before neonate larvae entering potting media.
Suppressed: Citrus thrips	2.50 mL/ft ³ container media	Use the higher rate for heavy infestations.

CITRUS (Field)

Including: Calamondin, Citrus citron, Citrus hybrids (includes chironja, tangelo, and tangor), Grapefruit, Kumquat, Lemon, Lime, Mandarin (tangerine), Pummelo, Orange (sweet and sour), Tangelo, Satsuma mandarin, White sapote(*Casimiroa* spp), and other cultivar and/or hybrids of these.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Sales Termites (FL only) Whiteflies	16.0 – 32.0 Fl oz/A	Apply recommended rate of Aura 2 F Insecticide using one of the following methods: 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment. For best results, apply to newly planted trees or those previously trained to drip, trickle, or micro-sprinkler irrigation. Lightly pre-wet the soil to break soil surface tension before applying Aura 2 F Insecticide. Chemigation application can be made separate to normal irrigation but should be followed by 10 to 20 minutes of additional watering to move Aura 2 F Insecticide into the root-zone. Allow 24 hours before initiating subsequent irrigations; 2. Soil surface band spray on both sides of the tree. Bands should overlap at the tree base to create a continuous band within the drip-line area of the tree, to be followed immediately with light sprinkler irrigation sufficient to move the product into the upper portion of the root-zone. This method is suitable for very coarse soils with 0.75% organic matter or less;
Suppressed Symptoms of: Citrus tristeza virus (CTV) through vector control Citrus yellows Thrips (foliage feeding thrips only)	32.0 Fl oz/A	

		<p>3. Drench to base of tree not exceeding one-quart total solution per tree immediately around trunk of tree and extending outward covering the entire fibrous root system of the tree. Only recommended for trees up to 8 feet tall;</p> <p>For control of existing termite infestations, apply specified dosage in 1 to 4 quarts of total solution volume, depending on size of tree, as a drench application to the basal portion of the tree trunk and surrounding soil in the immediate vicinity of the tree trunk.</p> <p>Pre-Harvest Interval (PHI): 0 day Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per season.</p>
--	--	--

CRANBERRIES

PESTS	RATE	INSTRUCTIONS
<p>Rootgrubs (Scarabaeidae) Rootworms (Chrysomelidae)</p>	<p>16.0 – 32.0 Fl oz/A</p>	<p>Apply Aura 2 F Insecticide to moist soil. Apply recommended rate of Aura 2 F Insecticide using one of the following methods:</p> <ol style="list-style-type: none"> 1. As a soil spray (ground application) directed to the root and crown area using at least 20 gal of water per acre; 2. As a chemigation application with 600 to 1000 gal water. <p>Immediately upon application, incorporate Aura 2 F Insecticide into root-zone by 0.1 – 0.3 inches water/Acre, either with the chemigation application or through irrigation/rainfall if not applied through chemigation. Inadequate incorporation within 24 hours of application may result in reduced control.</p> <p>Do not apply during bloom.</p> <p><u>Rootgrubs and Rootworms:</u> Best control may be achieved when application is made post-bloom immediately after bees are removed. Applications should target early instar larvae.</p> <p>Aura 2 F Insecticide has not been tested for crop response in tank mixes with other registered fungicides or insecticides. If tank mixing is desired, premix a sample of Aura 2 F Insecticide and the desired fungicide or insecticide partner at labeled rates and apply to a small area. Evaluate crop response within 48 hours and for at least two weeks prior to utilizing the tank mix on larger acreage. If crop injury results from the premix test, do not apply the tank mix to larger acreage.</p> <p>Pre-Harvest Interval (PHI): 30 days Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per season.</p>

--	--	--

GRAPES, Including: American bunch grape, Muscadine grape and Vinifera grape.

PESTS	RATE	INSTRUCTIONS
Controlled: Mealybugs Leafhoppers/Sharpshooters <i>Phylloxera</i> * spp.	16.0 – 32.0 Fl oz/A	Apply specified dosage of Aura 2 F Insecticide using one of the following methods: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation; 3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation. For best results, apply between bud-break and the pea-berry stage. * Repeated and regular use of Aura 2 F Insecticide over several, consecutive growing seasons controls existing <i>Phylloxera</i> infestations over time or prevents <i>Phylloxera</i> from becoming established. Pre-Harvest Interval (PHI): 30 days Do not apply more than 32.0 fl oz (0.50 lb)/A Aura 2 F Insecticide per season.
Suppressed: Pierce's disease	24.0 – 32.0 Fl oz/A	

HOPS

Not for use in California unless directed by supplemental labeling.

PESTS	RATE	INSTRUCTIONS
Aphids	19.2 Fl oz/A	Apply specified dosage of Aura 2 F Insecticide using one of the following methods: <ol style="list-style-type: none"> 1. Chemigation into root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Subsurface side-dress shanked into the root-zone on both sides of the plants followed by irrigation; 3. Hill drench in sufficient water to insure incorporation into the root-zone followed by irrigation. Pre-Harvest Interval (PHI): 60 days Do not apply more than 19.2 fl oz (0.3 lb ai)/A Aura 2 F Insecticide per season.

PECANS

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Twolined spittlebug Suppressed: Pecan scab (from reduction in honeydew deposition)	16.0 – 32.0 Fl oz/A	Applications may be made from May 15 until July 15. Applications made later in the season may be less effective. Apply product to slightly moist soil and allow soil to dry before additional irrigation. Apply recommended rate of Aura 2 F Insecticide using one of the following methods: <ol style="list-style-type: none"> 1. Chemigation to root-zone by means of low-pressure drip, trickle, micro-sprinkler or equivalent equipment; 2. Emitter or spot application in a minimum of 4 fluid ounces of mixture per emitter site; 3. Subsurface side-dress shanked into the root-zone near emitter line. Treat distance, wetted by the emitter set of each tree Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per season.

POME FRUITS

Including Apple, Crabapple, Loquat, Mayhaw, Pear (including Oriental pear), Quince

Not for this use in California except when allowed by Supplemental labeling.

PESTS	RATE	INSTRUCTIONS
Aphids (including wooly apple aphid) Leafhoppers	16.0 – 24.0 Fl oz/A	Apply specified dosage of Aura 2 F Insecticide through chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide season.

STONE FRUITS

Including: Apricot, Cherry (including sweet and tart), Nectarine, Peach, Plum (including Chickasaw, Damson and Japanese), Plumcot, Prune (fresh and dried)

Not for this use in California except when allowed by Supplemental labeling.

In-field Soil Application

PESTS	RATE	INSTRUCTIONS
Aphids (including wooly apple aphid) Leafhoppers	16.0 – 24.0 Fl oz/A	Applications Apply specified dosage of Aura 2 F Insecticide by chemigation into root-zone through low-pressure drip, trickle, micro-sprinkler, or equivalent equipment. Pre-Harvest Interval (PHI): 21 days Do not apply more than 24.0 fl oz (0.38 lb ai)/A Aura 2 F Insecticide per season.

Pre-Plant, Root Dip Application

PESTS	RATE	INSTRUCTIONS
Black peach aphid	2.0	Mix Aura 2 F Insecticide at recommended rate

(infesting roots)	Fl oz/10 gal root-dip solution	per 10 gallons of water. Completely wet bare-root transplant to slightly above the graft union by soaking them in the Aura 2 F Insecticide solution for up to 5 minutes. Allow solution to dry on roots then transplant trees as soon as possible following treatment.
-------------------	--------------------------------	--

TROPICAL FRUIT

Including: Acerola, Avocado, Black sapote, Canistel, Feijoa, Jaboticaba, Guava, Longan, Lychee, Marney sapote, Mango, Papaya, Passionfruit, Persimmon, Pulasan Rambutan, Sapodilla, Spanish lime, Star apple, Starfruit, Wax jambu

Not for this use in California except when allowed by Supplemental labeling.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Leafhoppers Whiteflies	24.0 – 32.0 Fl oz/A	Apply recommended rate of Aura 2 F Insecticide by chemigation through low-pressure drip, trickle, micro-sprinkler or equivalent equipment.
Suppressed: Scales	32.0 Fl oz/A	Pre-Harvest Interval (PHI): 6 days Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per application.

POPLAR/COTTONWOOD

Includes members of the genus *Populus* grown for pulp or timber

Not for this use in California except when allowed by Supplemental labeling.

PESTS	RATE	INSTRUCTIONS
Controlled: Aphids Cottonwood leaf beetle	16.0 – 32.0 Fl oz/A	Apply recommended rate of Aura 2 F Insecticide by chemigation through low-pressure drip irrigation.
Suppressed: <i>Phylloxera</i> <i>popularia</i>		Cottonwood leaf beetle: make application early, when the beetles first begin feeding, for best protection. Larger trees may need earlier treatment as a result of slower uptake. Phylloxera: apply early in the year, from break of dormancy through May. Do not apply more than 32.0 fl oz (0.50 lb ai)/A Aura 2 F Insecticide per crop season.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of

this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Deleted: To the extent consistent with applicable law, United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.]