

70506-168

09/06/2007

1/18



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs  
Registration Division (7505C)  
1200 Pennsylvania Ave., N.W.  
Washington, D.C. 20460

EPA Reg. Number:

70506-168

Date of Issuance:

SEP 6 2007

NOTICE OF PESTICIDE:

Registration  
 Reregistration

(under FIFRA, as amended)

Term of Issuance:

Conditional

Name of Pesticide Product:

Imidacloprid 4 F  
Termiticide/Insecticide

Name and Address of Registrant (include ZIP Code):

United Phosphorus, Inc.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
  - a. Revise the EPA Registration Number to read, "EPA Reg. No. 70506-168".
  - b. Within the CONCRETE SLAB-ON-GROUND OR BASEMENTS section of the label, revise the sentence "When rodding, rod holes should be spaced..." to read "When rodding, rod holes must be spaced..."

Signature of Approving Official:

Kable Bo Davis, Entomologist  
Insecticide-Rodenticide Branch, Registration Division (7505P)

Date:

SEP 6 2007

- c. Within the CRAWL SPACES: section of the label, revise the sentence "*Apply by trenching and rodding downward along...*" to read "*Apply by trenching or trenching and rodding downward along...*"
  - d. Within the BASEMENTS – OUTSIDE PERIMETER section of the label, revise the sentence "*Along the outside of the exterior walls, an apply by trenching or rodding within the trench.*" to read "*Along the outside of the exterior walls, an application must be made by trenching or rodding within the trench.*"
  - e. Within the CONTROL OF WOOD INFESTING PESTS section of the label, revise the statements "*For small trees (less than 8" in diameter, measured as the circumference at the root flare), apply 1 gallon of solution. For trees larger than 8", apply 4 gallons per 10 linear feet.*" to read "*For small trees (less than 6" in diameter, measured as the circumference at the root flare), apply 1 gallon of solution. For trees larger than 6", apply 4 gallons per 10 linear feet.*"
3. The data requirements for storage stability (830-6317) and corrosion characteristics (830- 6320) have not been satisfied, and must be submitted within eighteen months of the date of this letter.
  4. Submit one copy of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Kable Bo Davis  
Entomologist  
Insecticide-Rodenticide Branch  
Registration Division (7505P)

Enclosure

## IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE

**For use by individuals/firms licensed or registered by the state to apply termiticide products. States may have more restrictive requirements regarding qualifications of persons using this product. Consult the structural pest control regulatory agency of your state prior to use of this product.**

**For prevention or control of subterranean termites, drywood termites, dampwood termites, carpenter ants, and other wood-infesting insects.**

**ACTIVE INGREDIENT**

Imidacloprid - 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....	40.7%
OTHER INGREDIENTS.....	59.3%
<b>Total</b>	<b>100%</b>

Contains 4 pounds of active ingredient per gallon

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

<b>FIRST AID</b>	
If Swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If on Skin or Clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If Inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call an ambulance, then give artificial respiration.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If In Eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. Contact the Rocky Mountain Poison Center at 1-866-767-5089 for emergency medical treatment information.</p>	
<p><b>NOTE TO PHYSICIAN:</b> No specific antidote is available. Treat symptomatically.</p>	

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEM RECEPTED  
1-800-424-9300**

United Phosphorus, Inc.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406  
1-800-438-6071 • [www.upi-usa.com](http://www.upi-usa.com)

Contents:  
EPA Reg. No. 70506-  
EPA Est. No.

**RECEIVED  
with COMMENTS  
In EPA Letter Dated:**

SEP 6 2007

**Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under EPA Reg. No.  
70506-168**

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed, absorbed through skin, or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

When treating adjacent to an existing structure, the applicator must check the area to be treated, and immediately adjacent areas of the structure, for visible and accessible cracks and holes to prevent any leaks or significant exposures to persons occupying the structure. People present or residing in the structure during application must be advised to remove their pets and themselves from the structure if they see any signs of leakage. After application, the applicator is required to check for leaks. All leaks resulting in the disposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site. Do not allow people or pets to contact contaminated areas or to reoccupy contaminated areas of the structure until the clean up is completed.

**Personal Protective Equipment (PPE)**

Some materials that are chemical resistant to this product are listed below. More options can be obtained by following the instruction for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirts and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

In addition, all pesticide handlers must wear protective eyewear when working in a non-ventilated space or when applying termiticide by rodding or sub-slab injection.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Extreme care must be taken to avoid runoff. Apply only to soil or other fill substrate that will accept the solution at the specified rate. Do not treat soil that is water-saturated or frozen or in any condition where run-off or movement from the treatment area (site) is likely to occur.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Structures that contain wells or cisterns within the foundation of the structure can only be treated using the treated backfill method described in the treatment around wells and cisterns section of this label. Consult state and local specifications for recommended distances of wells from treated area, or if such regulations do not exist, refer to Federal Housing Administration Specifications (H.U.D.) for guidance.

**Do not formulate this product into other end-use products.**

**MIXING:** Refer to Mixing Table for proper amount of Imidacloprid 4 F Termiticide/Insecticide Termiticide/Insecticide to be used.

Mix the termiticide use dilution in the following manner. Fill tank  $\frac{1}{4}$  to  $\frac{1}{3}$  full. If using large sprayer, start pump to begin bypass agitation and place end of treating tool in tank to allow circulation through hose. Add appropriate amount of Imidacloprid 4 F Termiticide/Insecticide. Add remaining amount of water. Let pump run and allow recirculation through the hose for 2 to 3 minutes.

<b>MIXING TABLE FOR IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE</b>			
<b>GALLONS WATER</b>	<b>PLUS</b>	<b>0.05%</b>	<b>0.10%</b>
50		6.9 fl oz	13.8 fl oz
100		13.8 fl oz	27.6 fl oz
200		27.6 fl oz	55.2 fl oz
1,000		135.7 fl oz	271.4 fl oz

### APPLICATION VOLUME

The application volumes described in the directions for use should be used whenever possible. When soil conditions will not accept the application of 4 gallons per 10 linear feet, twice the amount of IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE may be applied in 2 gallons of solution per 10 linear feet.

### GENERAL CONTROL

Treatment standards for subterranean termite control may vary due to regulations, procedures used, soil types, construction practices and other factors. The purpose of chemical soil treatment for termite control is to establish a continuous chemical treated zone (horizontal or vertical as needed) between the wood and other cellulose material in the structure and the termite colonies in the soil. Follow all federal, state, and local regulations and treatment standards for protection of a structure from termites. In some cases where an aerial or above ground colony is established, supplemental treatments to control the termites, landscape modifications, and/or structural repairs may be needed to deprive termites of a moisture source. Use a 0.05% to 0.1% dilution based on local recommendations. Generally a 0.05% dilution is used for typical control situations, while a 0.1% dilution is normally used for severe or persistent infestations.

### PRE-CONSTRUCTION TREATMENT

Do not apply at a lower dosage and/or concentration than that specified on this label when applying before installation of the finished grade.

Prior to each application, applicators must notify the general contractor, construction superintendent, or similar responsible party, of the intended termiticide application and intended sites of application and instruct the responsible person to notify construction workers and other

individuals to leave the area to be treated during application and until the termiticide is absorbed into the soil.

**CONCRETE SLAB-ON-GROUND OR BASEMENTS:**

Apply an overall treatment to the entire surface of soil or other substrate to be covered by the slab including areas to be under carports, porches, basement floor and entrance platforms. Apply at the rate of 1 gallon of solution to accurately and uniformly cover 10 square feet. If fill under slab is gravel or other coarse aggregate, apply at the rate of 1.5 gallons or sufficient volume of solutions, to accurately and uniformly cover 10 square feet. In addition, apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet to provide a uniform treated zone in soil at critical areas such as along the inside of foundation walls, and around plumbing, bath traps, utility services, and other features that will penetrate the slab.

After completion of grading, make an application by trenching or trenching and rodding around the slab or foundation perimeter. Rodding may be done from the bottom of a shallow trench. When rodding, rod holes should be spaced in a manner that will allow for a continuous chemical treated zone, no greater than 12 inches, to be deposited along the treated area. Rod holes should not extend below the footing. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet, per foot of depth to provide a uniform treated zone. When trenching, the trench along the outside foundation should be about 6 inches in depth. Use a low pressure spray (not to exceed 25 psi at the treatment tool when the valve is open) to treat soil which will be placed in the trench after rodding. Mix the spray solution with soil as it is being placed in the trench. When treating voids in hollow masonry units, use 2 gallons of solution per 10 linear feet of wall. Apply solution so it will reach the footing by injecting into the lower areas of the wall, just above the floor or footing.

When treating foundations deeper than 4 feet, apply the termiticide as the backfill is being replaced, or if the construction contractor fails to notify the applicator to permit this, treat the foundation to a minimum depth of 4 feet after the backfill has been installed. The applicator must trench and rod into the trench or trench along the foundation walls and around pillars and other foundation elements, at the rate prescribed from grade to a minimum depth of 4 feet. When the top of the footing is exposed, the applicator must treat the soil adjacent to the footing to a depth not to exceed the bottom of the footing. However, in no case should a structure be treated below the footing.

Rodding in trench followed by flooding of trench and treatment of backfill may create a better continuous chemical treated zone than using soil rodding alone to establish a vertical termiticide treated zone.

**CRAWL SPACES:** Apply by trenching and rodding downward along the inside and outside of foundation walls, around piers, interior supports in contact with the soil, plumbing, and utility services. Apply four gallons of solution (see APPLICATION VOLUME) per 10 linear feet, per foot of depth to create a uniform treated zone. Rodding may be done from the bottom of a shallow trench to top of the footing or a minimum of 4 feet. When rodding, space rod holes to create a continuous chemical treated zone along the treated area. Rod holes should not extend below the footing. When trenching, the trench should be about 6 inches wide and 6 inches deep. Use a low pressure spray to treat soil which will be placed in the trench, mixing the spray solution and soil together as it is being placed in the trench.

**HOLLOW BLOCK FOUNDATIONS OR VOIDS:** Hollow block foundations or voids in masonry resting on the footing may be treated to provide a continuous chemical treated zone in the voids at the footing. Apply 2 gallons of solution per 10 linear feet to the lower part of the void so that it reaches the top of the footing or soil.

Examine treatment of voids in block or rubble foundation walls closely. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site (refer to Precautionary Statements). Do not allow people or pets to contact or to reoccupy the contaminated areas of the structure until the clean up is completed.

## POST-CONSTRUCTION TREATMENT

**CONCRETE-SLAB-ON-GROUND:** To apply under the slab in areas such as attached porches, carports, entrance platforms, garages and similar slab structures, it may be necessary to drill through the slab or exterior foundation. Space drill holes should be spaced so that a continuous chemical treated zone is created. Treat all existing cracks and cold, construction, or expansion joints. Also, treat around bath traps, plumbing and utility services which penetrate the slab. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet per foot of depth to provide a uniform treated zone. DO NOT TREAT UNTIL LOCATION OF HEAT OR AIR CONDITIONING DUCTS AND VENTS IS KNOWN AND EACH IS LOCATED. USE EXTREME CAUTION TO AVOID CONTAMINATION OF DUCTS AND VENTS. Plug and fill all drilled holes in commonly occupied areas with a suitable sealant. If plugs are not constructed of non-cellulose material, they must be covered by an impervious, non-cellulose material.

An application should be made by trenching or trenching and rodding around the outside of the foundation wall. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet per foot of depth to create a uniform treated zone. When trenching, make the trench along the outside foundation about 6 inches wide and 6 inches deep. Use a low pressure spray to treat soil as it is being placed in the trench.

Rodding can be done from the bottom of a shallow trench. When rodding, space rod holes to create a continuous chemical treated zone, not greater than 12 inches, to be deposited along the treated area. Rod hole depth should not extend below the footing.

**BATH TRAPS:** Exposed soil or soil covered with tar or a similar type sealant under and around plumbing and/or drain pipe entry areas should be treated with 3 gallons of solution per square foot. Cut/install an access door or inspection vent if one is not already present. Inspect and remove any wood or cellulose debris, then treat soil by rodding or drenching the soil.

**CRAWL SPACES:** When there is not enough clearance between floor joists and ground surfaces to allow applicator access, excavate the area, if possible, and treat according to crawl spaces (refer to Pre-Construction Treatment). If excavation is not possible, treat crawl space soil and wood to prevent surface access by termites. Apply 1 gallon of solution (see APPLICATION VOLUME) per 10 square feet to create a uniform chemical treated zone. Use a very coarse spray at a pressure not greater than 25 psi at the treatment tool when the valve is open.

If a crawl space cannot be reached with the application wand, use extension wands or other suitable equipment to apply a coarse spray on the soil, wood and structural members which contact the soil, at the rates above. Do not apply to inaccessible crawl space areas using pressures greater than 25 psi at the treatment tool when the valve is open.

Treatment may also be made by drilling through the foundation wall or through the floor above and treating the soil perimeter using a rate of 1 gallon of solution per 10 square feet. Drill

spacing must be at intervals not to exceed 16 inches. Many states have smaller intervals so check state regulations which may apply.

To prevent subterranean termites from constructing mud tubes between soil and crawl space wood members above, an overall soil treatment of this product may be applied. Remove all cellulose debris before application. Apply 1 gallon of solution (see APPLICATION VOLUME) per 10 square feet to provide a uniform chemical treated zone.

**SHALLOW FOUNDATIONS:** For shallow foundations, one foot or less in depth, dig a narrow trench about 6 inches wide and deep along both the outside and inside of the foundation walls, being careful not to dig below the bottom of the footings. If the foundation has an exposed footing, dig a trench alongside the footing taking care not to undermine the footing. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet to the top of footer to create a uniform treated zone. Apply the dilution to the trench and mix it with the soil as it is placed in the trench.

**BASEMENTS – OUTSIDE PERIMETER:** Along the outside of the exterior walls, an apply by trenching or rodding within the trench. Rodding depth should be to the top of the footer, or to a minimum of 4 feet or according to state or local regulations. When rodding through a trench, dig a narrow trench about 6 inches wide and 6 inches deep, then apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet, per foot of depth to create a uniform treated zone by rodding through the trench. Use a low pressure spray to treat soil which will be placed into the trench after rodding. Mix spray solution with the soil as it is being placed in the trench.

**BASEMENTS – INSIDE PERIMETER:** If necessary, treat by drilling along the perimeter of the interior walls. Applications also may be necessary around sewer pipes, floor drains, conduits, expansion joints or any cracks or holes in the basement floor. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet to create a uniform treated zone.

Space drill holes to allow for creation of a continuous chemical treated zone. Plug and fill all drill holes in commonly occupied areas of the building with a suitable sealant. If plugs are not made of non-cellulose material, they must be covered by an impervious, non-cellulose material.

**HOLLOW BLOCK FOUNDATION OR VOIDS:** Hollow block foundations or voids in masonry resting on the footing may be treated to create a continuous chemical treated zone in the voids at the footing. Apply 2 gallons of solution per 10 linear feet to the lower part of the void so that it reaches the top of the footing or soil. Space drill holes so that intervals do not exceed 16 inches. Many states have smaller intervals so check state regulations which may apply.

Examine treatment of voids in block or rubble foundation walls closely as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment.

All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site (refer to Precautionary Statements). Do not allow people or pets to contact or to reoccupy the contaminated areas of the structure until the cleanup is completed.

**PLENUMS:** For plenum-type structures which use a sealed underfloor space to circulate heated and/or cooled air throughout the structure, apply the dilution at the rate of 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet, per foot of depth of soil to create a uniform treated zone adjacent to both sides of foundation walls, supporting piers, plumbing and conduits. Treat the soil by trenching to a depth of 6 inches or trenching and rodding (where conditions permit) or to the top of the footing. Where trenching or rodding are not possible, a surface



application next to interior foundation walls may be made, but the treated strip shall not exceed a width of 18 inches, horizontally, from the foundation walls, piers or pipes. Make the surface application at a rate of 1.5 gallons of solution per 10 square feet as very coarse spray under low pressure (not greater than 25 psi when measured at the treating tool when valve is on).

When treating plenums, turn off air circulation system of the structure until application is finished and all termiticide has been absorbed by the soil.

**TREATMENT AROUND WELLS OR CISTERNS:** Do not contaminate wells or cisterns.

**Structures with Wells/Cisterns Inside Foundations:** Structures that contain wells or cisterns within the foundation of a structure may only be treated using the techniques below.

1. Do not apply within 5 feet of any well or cistern by rodding and/or trenching or by the backfill method. Treat soil between 5 and 10 feet from the well or cistern by the backfill method only. Treatment of soil adjacent to water pipes within 3 feet of grade should only be done by the backfill method.
  - a) Trench and remove soil to be treated onto heavy plastic sheeting or similar material or into a wheelbarrow.
  - b) Treat the soil at the rate of 4 gallons of solution per 10 linear feet per foot of the trench, or 1 gallon per 1.0 cubic feet of soil. Mix thoroughly into the soil taking care to contain the liquid and prevent runoff or spillage.
  - c) After the treated soil has absorbed the solution, replace the soil into the trench.
2. Treat infested and/or damaged wood in place using an injection technique such as described in the "Control of Wood Infesting Pests" section of this label.

**Structures with Adjacent Wells/Cisterns and/or Other Water Bodies:** Applicators must inspect all structures with nearby water sources such as wells, cisterns, surface ponds, streams, and other bodies of water and evaluate, at a minimum, the treatment recommendations listed below before making an application.

1. Before treatment, if feasible, expose the water pipe(s) coming from the well to the structure, if the pipe(s) enter the structure within 3 feet of grade.
2. Before treatment applicators are advised to take precautions to limit the risk of applying the termiticide into subsurface drains that could empty into any bodies of water. These precautions include evaluating whether application of the termiticide to the top of the footer may result in contamination of the subsurface drain. Consideration should be given to factors such as depth to the drain system and soil type and degree of compaction when determining the depth of treatment.
3. When appropriate (i.e., on the water side of the structure), the treated backfill technique (described above) can also be used to minimize off-site movement of termiticide.

## **EXTERIOR PERIMETER/INTERIOR SPOT TREATMENT**

### **General Information**

Exterior Perimeter/Interior Spot Treatment is an optional method of termite treatment only for use in post-construction applications after the final grade is established. Structural protection when using the Exterior Perimeter/Interior Spot Treatment is accomplished by: 1) establishing a continuous treated zone around the entire exterior foundation wall of the building; and 2) spot-treating infested areas on the building interior. Soil adjacent to the exterior foundation wall must be treated in the same manner as conventional (full) application. It is required that a complete and continuous treated zone be achieved around the entire exterior perimeter, including under any attached slabs such as garages, porches, patios, driveways and pavement adjoining the foundation. Interior spot treatments must then be made to any indoor areas where termite activity is present. Optional interior spot treatments may also be made to high risk areas including, but not limited to, plumbing and utility penetrations (including bath traps), along settlement cracks and expansion joints, and dirt-filled porches.

Exterior Perimeter/Interior Spot Treatment can be used either as a preventative treatment (before structural infestation occurs) or as a curative treatment (after structural infestation occurs) in existing structures. Preventative treatment does not include pre-construction applications made to protect new construction. It is required that a thorough structural inspection be completed, before treatment, to locate all areas of active infestation. Spot treatment of all known sites of termite activity is required with this optional method of treatment. If no termite activity is observed inside the structure, interior spot treatments are not required. Do not apply at a lower dosage and/or concentration than specified on this label.

#### **EXTERIOR PERIMETER TREATMENT**

It is required that all structures, regardless of the type of construction, be protected by establishing a vertical treated zone along the outer perimeter of the foundation wall. Consult the OUTER FOUNDATION WALLS section of this label for detailed directions on this treatment procedure.

**1. OUTER FOUNDATION WALLS:** Application must be made by trenching, or where appropriate (see below) by trenching, or trenching and rodding from the bottom of the trench, around the outside of the foundation walls. When trenching, excavate a trench along the outside foundation that is about 6 inches wide and 6 inches deep. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet, per foot of depth to provide a uniform vertical treated zone.

- For shallow foundations, one foot or less depth, dig a narrow trench that does not exceed 6 inches wide and 6 inches deep along the outside of the foundation walls, being careful not to dig below the bottom of the footings. For foundations with exposed footings, dig a trench alongside the footing taking care not to undermine the footing.
- For basements and other foundations deeper than one foot, the application must be made by trenching and rodding from the bottom of a shallow trench. When rodding, rod holes must be spaced in a manner that will allow for a continuous treated zone, not exceeding 12 inches, to be deposited along the treated area. Rod holes must not extend below the footing. Rodding depth should be to the top of the footer, or to a maximum depth of 4 feet, or according to state or local regulations.

For all applications, apply the solution into the trench and mix with the excavated soil as it is replaced into the trench. Use a low-pressure spray to treat soil that will be replaced into the trench after rodding. Mix spray solution with the soil as it is being replaced in the trench.

**Note:** Where direct access to soil on the outer foundation wall is impossible due to attached porches, entrance platforms, garages and similar slab structures, consult the CONCRETE SLAB-ON-GROUND section of this label for directions on treatment of soil beneath these structures. However, where obstructions (e.g., concrete walkways) adjacent but not attached to foundation, or where soil type and/or conditions, prevent trenching the exterior perimeter treatment may be performed at the obstructed location by rodding alone. When rodding, rod holes must be spaced in a manner that will allow for a continuous treated zone, not exceeding 12 inches, to be deposited along the treated area.

**2. CONCRETE SLAB-ON-GROUND:** To treat soil beneath a slab, including attached porches, carports, entrance platforms, garages, and similar slab structures abutting the foundation wall, it is necessary to drill through the slab. If an infestation is associated with an expansion joint crack, utility penetration, or similar access point in the slab, treat by drilling and injecting through the slab. Drill holes must be spaced in a manner that will allow for application of a continuous chemical treated zone, but must extend a minimum of 3 feet on both sides of the infested site. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 liter feet.

DO NOT MAKE TREATMENT UNTIL LOCATION OF HEAT OR AIR CONDITIONING DUCTS AND VENTS ARE KNOWN AND IDENTIFIED. USE EXTREME CAUTION TO AVOID CONTAMINATION OF DUCTS AND VENTS. Plug and fill all drilled holes in commonly occupied areas with suitable sealant. Plugs must be of non-cellulose material or covered by an impervious, non-cellulose material.

**3. INACCESSIBLE CRAWL SPACES:** If termite activity is found along the perimeter wall or on a pier within an inaccessible crawl space, areas with termite activity must be treated. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet to create a vertical treated zone, which must extend a minimum of 3 feet on both sides of the infested site.

Optional directions for horizontal rodding: Treatment may also be made by drilling through the foundation wall (or through the floor above) to treat the soil along the perimeter wall at a rate of 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet. Drill spacing must be at intervals not to exceed 16 inches. Many states have shorter intervals so check state regulations which may apply. If termite activity is neither along the perimeter wall nor on a pier within the inaccessible crawl space, to prevent subterranean termites from constructing mud tubes between soil in the crawl space and wooden elements in the structure, an overall soil treatment of this product may be applied. Remove all cellulose debris before application. Apply 1 gallon of solution (see APPLICATION VOLUME) per 10 square feet to provide a uniform chemical treated zone.

**4. ACCESSIBLE CRAWL SPACES:** If termite activity is found within an accessible crawl space, the area(s) where termite activity exist must be treated by trenching, or trenching and rodding from the bottom of the trench, along the interior foundation walls, around piers, interior supports in contact with the soil, plumbing, or utility services. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet, per foot of depth, to create a vertical treated zone, which must extend a minimum of 3 feet on both sides of the infested site. Rodding may be done from the bottom of a shallow trench to the top of the footing or to a minimum of depth of 4 feet. When rodding, rod holes must be spaced in a manner that will allow for a continuous treated zone, not to exceed 12 inches, to be deposited along the treated area. Rod holes must not extend below the footing. When trenching, dig a narrow trench about 6 inches wide and 6 inches deep. Use a low-pressure spray to treat soil which will be placed in the trench, mixing the spray solution with soil as it is being placed in the trench.

#### INTERIOR SPOT TREATMENT

Targeted applications must be made to all known infested sites inside the structure. One or more of the following application methods must be used to make interior spot treatments:

- Sub-slab injections made through the slab at or near areas where termites are known to be penetrating the slab to reach wood in the structure and/or at or near sites of active infestations. Apply 4 gallons per 10 linear feet per foot depth. Sub-slab injections must extend to a minimum of 3 feet on both sides of every known infested site at expansion joints or cracks in slabs.
- Void treatments using injection of sprays, mists or foams into above ground structural voids, termite carton nests, and other infested locations.
- Wood treatments using injection techniques and/or surface applications, to treat active infestations in structural timbers.

To maximize dispersion of treatment solution in soil and in above ground locations, the use of foam and directional dispersion tips is encouraged for all interior spot treatments. Consult section(s) of this label appropriate to the element of construction, FOAM APPLICATIONS or CONTROL OF WOOD INFESTING PESTS for detailed directions on any of these treatment procedures.

**1. INTERIOR SLABS:** When termite activity is located within an interior wall or structural member, the soil beneath the slab and the wall void at this site of activity must be treated. The

source of infestation at an expansion joint, crack, through a utility penetration, or similar access point in the slab, must be treated by drilling and injecting through the slab. Drill holes in the slab must be spaced in a manner that will allow for application of a continuous chemical treated zone, which must extend a minimum of 3 feet on both sides of the infested site. Apply 4 gallons of solution (see APPLICATION VOLUME) per 10 linear feet. To maximize dispersion of treatment solution in soil, the use of foam and directional dispersion tips is encouraged. To treat the wall void, consult section(s) of this label appropriate to the element of construction, FOAM APPLICATIONS or CONTROL OF WOOD INFESTING PESTS for detailed directions on any of these treatment procedures.

**DO NOT MAKE TREATMENT UNTIL LOCATION OF HEAT OR AIR CONDITIONING DUCTS AND VENTS ARE KNOWN AND IDENTIFIED. USE EXTREME CAUTION TO AVOID CONTAMINATION OF DUCTS AND VENTS.** Plug and fill all drilled holes in commonly occupied areas with suitable sealant. Plugs must be of non-cellulose material or covered by an impervious, non-cellulose material.

**2. HOLLOW BLOCK FOUNDATION OR MASONRY VOIDS:** Termite activity located within hollow-block foundations or masonry voids must be treated. Spot treatment at the site(s) of termite activity must extend a minimum of 3 feet on both sides. Treat masonry voids by applying 2 gallons of solution per 10 linear feet to the lower part of the void so that it reaches the top of the footing or soil. Drill spacing in masonry voids must be at intervals not to exceed 16 inches; states may have shorter intervals so check state regulations which may apply. To maximize dispersion of treatment solution in voids, the use of foam and directional dispersion tips is encouraged. To treat structural voids above sites of termite activity in masonry, consult section(s) of this label appropriate to the element of construction, FOAM APPLICATIONS or CONTROL OF WOOD INFESTING PESTS for detailed directions on any of these treatment procedures. Treatment of voids in block or rubble foundation walls must be closely examined. Applicators must inspect areas of possible runoff as a precaution against application leakage in the treated areas. Some areas may not be treatable or may require mechanical alteration prior to treatment. All leaks resulting in the deposition of termiticide in locations other than those prescribed on this label must be cleaned up prior to leaving the application site (refer to Precautionary Statements). Do not allow people or pets to contact or to reoccupy the contaminated areas of the structure until the cleanup is completed.

**3. BATH TRAPS:** If termite activity is observed within 2 feet of the bath trap, then exposed soil or soil covered with tar or a similar type of sealant around plumbing and/or drain pipe entry areas must be treated. Tar or sealant may have to be removed to allow for adequate soil treatment. An access door or inspection portal should be installed if one is not present. After inspection and removal of any wood or cellulose debris, the soil can be treated by rodding or drenching the soil at a volume of no less than 3 gallons of solution per square foot.

**4. SHOWER OR FLOOR DRAINS:** If termite activity is observed within 2 feet of a shower or floor drain in the slab, then soil beneath the drain must be treated. Drill through the slab adjacent to the drain and use sub-slab injection to apply solution to the soil. Multiple access points may be drilled adjacent to the drain. Treat soil at a volume of 1 gallon of solution per square foot.

#### **FOAM APPLICATIONS**

In some cases construction practices, soil subsidence and other factors may make it difficult to create a continuous chemical treatment zone using conventional treatment alone. In situations where necessary, supplement conventional application methods by use of foam generating equipment, or similar devices, to create a continuous treated zone.

Foam application may be made alone or combined with conventional application methods, provided that the labeled amount of active ingredient per unit area is used.

**Foam Application Use Directions:** Mix solution of IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE with manufacturer's recommended volume of foaming agent (see table for foaming recommendations). Apply a sufficient volume of IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE foam alone or in combination with liquid solution to create a continuous treated zone at the recommended rate for specific application sites. Use appropriate dispersion tips and application method for site.

<b>MIXING TABLE FOR IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE FOAM</b>				
<b>FL. OZ. OF IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE</b>	<b>Gallons of Water</b>	<b>Foam Expansion Ratio</b>	<b>Finished Foam</b>	
			<b>(gallons)</b>	<b>(ai%)</b>
3.5 ounces	1	25:1	25	0.05
	2.5	10:1		
	5	5:1		
7.0 ounces	1	50:1	50	
	2.5	20:1		
	5	10:1		
* Add the manufacturer's recommended quantity of foam agent to the IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE solution.				

Depending on the circumstances, foam applications may be used alone or combined with liquid solution applications. Applications may be made behind veneers, piers, chimney bases, into rubble foundations, into block voids or structural voids, wall voids, under slabs, stoops, porches, or to the soil in crawl spaces, and other similar voids.

Foam and liquid applications must be consistent with volume and active ingredient instructions in order to ensure proper application has been made. The volume and amount of active ingredient are essential to an effective treatment. At least 75% of the gallons of IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE must be applied as a typical liquid treatment. The remaining 25% or less gallons are delivered to appropriate locations using a foam application.

NOTE: When foam is used solely to kill subterranean termites in above ground locations (such as feeding galleries in wooden framing, or in voids with framed walls), and whenever the target pest is other than subterranean termites (drywood termites, beetles, ants, etc.), dilute solutions of imidacloprid may be expanded by foaming without concentrating the imidacloprid solution as previously described for soil applications. Add the manufacturer's recommended volume of foaming agent to produce foam of the desired expansion ratio. Use application tips and methods suitable to the site and pest.

### **CONTROL OF WOOD INFESTING PESTS**

For control of **above ground termites and carpenter ants** in localized areas, apply a 0.05 to 0.1% solution or sufficient volume of IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE foam to voids and galleries in damaged wood, and in spaces between wooden structural members and between the sill plate and foundation where wood is vulnerable. Apply to inaccessible areas by first drilling, then injecting the suspension or foam with a suitable directional injector into the damaged wood or wall voids. Termite carton nests in building voids may be injected with a 0.05 to 0.1% suspension or foam. It may be necessary to inject at multiple points to varying depths. When such nests are found, it is best to physically remove carton nest material. Apply to attics, crawl spaces, unfinished basements, or man-made voids using a coarse fan spray of 0.05 to 0.1%

solution or foam to control exposed workers and winged reproductive forms of termites or carpenter ants. This type of application is intended to be a supplemental treatment for control of above ground subterranean termites and carpenter ants.

It is recommended to remove or prune away any shrubbery, bushes, and tree branches touching the structure, because vegetation touching the structure may offer entry to ants. This may allow ants to inhabit the structure without coming in contact with the treatment. If nests are found, apply IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE Insecticide directly to these nests.

To control existing infestations of or to prevent infestation by termites or carpenter ants in trees, utility poles, fencing and decking materials, landscape timbers and similar non-structural wood-to-soil contacts, use a 0.05% to 0.1% solution. If possible, locate the interior infested cavity and inject a 0.05 to 0.1% solution or sufficient volume of IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE foam using an appropriate treatment tool with splashback guard. These non-structured wood-to-soil contacts may also be treated by applying a solution to the soil as a spot application or continuous treated zone applied as a drench or by rodding around the base of the point(s) of soil contact(s). Place the rod holes about 3 inches away from the soil contact point(s) and no more than 12 inches along the perimeter of the soil contact(s). For small poles or posts (<6 inches in diameter), apply 1 gallon per foot of depth. For larger constructions, apply 4 gallons per 10 linear feet per foot of depth. Retreat as needed to ensure protection.

Termite carton nests in trees may be injected with a 0.05 to 0.1% solution or sufficient volume of foam using a pointed injection tool. It may be necessary to use multiple injection points to varying depths. Removal of carton material from trees is preferable but may not be necessary when foam application is used. In some cases, a perimeter application of a 0.05% to 0.1% solution applied to soil around the root flare of the tree may be needed to prevent reinfestation by termites in the soil. For small trees (less than 8" in diameter, measured as the circumference at the root flare), apply 1 gallon of solution. For trees larger than 8", apply 4 gallons per 10 linear feet.

For protection of firewood or other wood products stored in contact with soil from carpenter ants and termites, before stacking treat soil with a 0.05 to 0.1% solution at 1 gallon per 10 square feet to prevent infestation. Curative application to the soil around firewood or other wood products stored in contact with soil may be made as described for non-structural wood-to-wood contacts (above).

**Drywood termites and wood-infesting beetles or borers** (such as, but not limited to, powder post beetles, anobiid or deathwatch beetles, false powder post beetles, old house borers, wharf borers, or ambrosia oak bark beetles). **Galleries and structure voids** can be treated with sprays, mists, or foams of a 0.05% to 0.1% IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE solution. Locate galleries by using visual signs (frass or pellets, blistered wood, emergence or clean cut holes), the presence of live insects, mechanical sounding techniques, or listening devices (e.g., stethoscopes, acoustic emission detectors). Drill holes to receive the injector tip or treatment tool. Distribute drill holes to adequately cover the gallery system. (NOTE: Avoid drilling where electrical wiring, plumbing lines, etc. are located). Apply IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE solutions using low pressure (about 20 psi) spray or by misting, or where appropriate, by foaming. It is not necessary to treat to the point where runoff is detected from adjacent holes. (NOTE: Do not apply where electrical shock hazards exist). Seal drill holes after treatment. Also, **wood surfaces** can be sprayed or misted with a 0.05% to 0.1% solution, or where appropriate, use a sufficient volume of foam. For inaccessible surfaces, drill and treat the interior of structural voids. Surfaces treated may include exposed wooden surfaces in crawl spaces, basements, or attics, wooden exterior surfaces such as decks, fencing, or siding, structural voids, channels in damaged wood, in spaces between members of a structure, and in junctions between wood and foundations. Apply by brushing or as a coarse, low pressure (about

20 psi) spray to the wood surface; apply sufficient volume to cover the surface to the point of wetness, but avoid applying to the point of runoff. When spraying overhead in living areas, cover surfaces below the treated area with plastic sheeting or similar material. Avoid contact with treated surfaces until spray deposits have dried. Retreat as needed to maintain protection.

**Localized treatment for carpenter bees:** Apply a 0.05% to 0.1% solution as a spray or mist, or sufficient volume of foam, directly into gallery entrance holes. After treatment, plug entrance holes with small pieces of steel wool or similar material.

#### **RETREATMENT**

Retreatment for subterranean termites may only be performed if there is clear evidence of reinfestation or if there has been disruption of the treated zone due to construction, excavation, or landscaping and/or evidence of the breakdown of the termiticide treated zone in the soil. Vulnerable or reinfested areas may be retreated in accordance with application techniques described in this label. The timing and type of retreatments will depend on factors such as termite pressure, soil types, soil conditions, and other factors which may reduce the effectiveness of the treated zone. Retreatment may be made as either a spot or complete treatment.

When a structure is not known to be reinfested and the treated zone is not disturbed, but where the structure was last treated five or more years ago, retreatment may be performed, if, in the judgment of the applicator, it is necessary to ensure the adequate protection of the structure. In determining the timing of any retreatment, the applicator should consider efficacy and/or degradation data and/or site-specific conditions and previous experience that indicate a vulnerability of the structure to termite attack.

**Annual treatment of the structure is prohibited unless there is clear evidence that reinfestation or treated zone disruption has occurred.**

**When another registered termite control product/system is used as the primary treatment for prevention or control of subterranean termites and is applied to all label-specified areas, IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE may be applied as a spot application in a secondary treatment to critical areas of the structure including plumbing and utility entry sites, bath traps, expansion joints, foundation cracks, the outside foundation wall and areas of known or suspected activity at either a pre-construction or post-construction timing. Such secondary treatments must be applied in amounts and concentration in accordance with label directions relevant to the treatment area(s) to receive the secondary treatment.**

**For control of ants in houses and other structures,** apply a 0.05% to 0.1% solution as a general surface, spot crack and crevice or wall void application. Apply to surfaces of buildings, porches, patios, and other structures, around doors and windows, eaves and attic vents, utility entry points, soffit areas and other exterior openings (including foundation cracks or drilled holes) where ants enter the structures or where they crawl or hide. Spray into cracks and crevices. Spray, mist or foam into voids where these ants or their nests are present. Apply s volume of spray mist or foam sufficient to cover the area, but do not allow excessive dripping or runoff to occur from vertical or overhead surfaces.

Treat soil, turf or ground cover next to the structure where ants are trailing or may find food or harborage. Apply to flower, shrub or ornamental plant beds next to the structure where ants may find food or forage. To control ants tunneling in soil apply a 0.05% to 0.1% solution as a drench or soil injection at intervals to establish a continuous treated zone. Treat along the edge of walls, driveways or other hard surfaces where ants are tunneling beneath the surface.

**Aerial Nests:** If any nests are located in tree hollows or non-structural wooden construction (e.g., posts, fences, decks) treat the interior cavity and/or the nest site by injecting a 0.05% to 0.01% solution as a spray mist, or sufficient volume of foam.

Apply in sufficient water to cover the foliage and soil area being treated. Maximum application is once per month to maintain control.

Do not allow residents or pets into the immediate area during the application or contact with treated areas until spray has dried.

Interior applications for ant control are limited to spot, crack and crevice, or wall void applications only.

**Do not use this product against native or imported fire ants, pharaoh or harvester ants.**

**NOTE:** Where severe pest pressures may exist and when rapid knockdown or exclusion at pest entry points is desired, make supplemental treatments using IMIDACLOPRID 4 F TERMITICIDE/INSECTICIDE with targeted applications of a pyrethroid such as TEMPO® SC ULTRA, or SUSPEND® SC to doors and windows, utility entry points, and other places where these pests enter the structure. Read and follow all label directions for use of this companion product.

#### **GENERAL PRECAUTIONS FOR APPLICATIONS**

After treatment, plug and fill all holes in concrete slab areas of the building with a suitable sealant.

Do not apply solution until location of heat pipes, ducts, water and sewer lines and electrical conduits are known and identified. Take care to avoid puncturing and injection into these structural elements.

Do not plant for the purpose of consumption, edible plants into the treated areas of soil.

Avoid contamination of public and private water supplies.

Use anti-backflow equipment or an air gap on filling hoses.

Consult State, Federal or local authorities for information regarding the approved treatment practices for areas in close proximity to potable water supplies.

#### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place, out of direct sunlight, and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Handle and open container in a manner as to prevent spillage. If the container is leaking, invert to prevent leakage. If the container is leaking or material spilled for any reason or cause, carefully dam up spilled material to prevent runoff. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Absorb spilled material with absorbing type compounds and dispose of as directed for pesticides below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**Container Disposal:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



**IMPORTANT INFORMATION  
READ BEFORE USING PRODUCT**

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

ADMIRE, CALYPSO, LEVERAGE, AND PROVADO are registered trademarks of Bayer.  
Actara, Centric, Cruiser and Platinum are trademarks of the Syngenta Group Company.  
Assail and Intruder are registered trademarks of Nippon Soda Company, LTD.  
Bidrin is a registered trademark of AMVAC Chemical Corporation.

**LABEL CHANGE NOTES**

**Rev. 8/2/07** – separate off termite uses for separate registration submission

**Rev. 5/2/07** – further revisions requested by EPA in telephone call of 5/2/07, including use 1<sup>st</sup> Storage & Disposal statement in place of second Storage & Disposal statement then delete 1<sup>st</sup>, remove second Rotational Crop section

**Rev. 4/30/07** – revision during initial registration process to remove uses on termites from this label as per EPA. Termite use to be submitted as a separate label as required by EPA.