

70506-172

5/14/2009

10614

5849757

332/1300



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

MAY 14 2009

Mr. Timothy M. Formella  
Senior Registration Manager  
United Phosphorus, Inc.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406

Subject: Kryocide Insecticide  
EPA Reg. No. 70506-172  
Notification Application dated April 21, 2009

Dear Mr. Formella:

The Agency has received your Application for Pesticide Notifications under Pesticide Registration Notice (PRN) 2007-4 dated April 21, 2009, for the product Kryocide Insecticide, EPA Reg. No. 70506-172. The Registration Division (RD) has reviewed this request for applicability under PRN 2007-4 and finds that it falls within the scope of PRN 2007-4. The label submitted with the application has been stamped "Notification" and will be placed in our records. You informed me via telephone yesterday that this product is sold only in bags. The revised "**STORAGE AND DISPOSAL**" is appropriate for such containers.

If you have any questions, please call me directly at 703-305-6406.

Sincerely,

A handwritten signature in black ink, appearing to read "William W. Jacobs".

William W. Jacobs  
Insecticide-Rodenticide Branch  
Registration Division (7505P)  
Office of Pesticide Programs

2 of 14

**EPA**

United States  
**Environmental Protection Agency**  
 Washington, DC 20460

☐ **Registration**  
☐ **Amendment**  
☒ **Other:**

OPP Identifier Number

**Application for Pesticide - Section I**

1. Company/Product Number  
 70506-172

2. EPA Product Manager  
 Autumn Metzger

3. Proposed Classification

4. Company/Product (Name)  
 KRYOCIDE Insecticide

PM#  
 1

☒ None ☐ Restricted

5. Name and Address of Applicant (Include ZIP Code)  
 United Phosphorus, Inc.  
 630 Freedom Business Center, Suite 402  
 King of Prussia, PA 19406

6. **Expedited Review.** In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to:

EPA Reg. No. \_\_\_\_\_

**NOTIFICATION**

Product Name \_\_\_\_\_

**MAY 14 2009**

☐ Check if this is a new address

**Section - II**

☐ Amendment - Explain below.

☐ Final printed labels in response to Agency letter dated \_\_\_\_\_

☐ Resubmission in response to Agency letter dated \_\_\_\_\_

☐ "Me Too" Application

☒ Notification - Explain below.

☐ Other - Explain below

**Explanation:** Use additional page(s) if necessary. (For Section I and Section II.)

Notification of label change per PR Notice 2007-4. This notification is consistent with the guidance in PR Notice 2007-4 and the requirements of EPA's regulations at 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156. No other changes have been made to the labeling or the Confidential Statement of Formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if the amended label is not consistent with the requirements of 40 CFR §§ 156.10, 156.140, 156.144, 156.146, and 156.156, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

**Section - III**

1. Material This Product Will Be Packaged In:

Child-Resistant Packaging

☐ Yes\*  
☐ No

Unit Packaging

☐ Yes  
☐ No

Water Soluble Packaging

☐ Yes  
☐ No

2. Type of Container

☐ Metal  
☐ Plastic  
☐ Glass  
☐ Paper  
☐ Other (laminated bag)

**\*Certification must  
 be submitted**

If "Yes"  
 Unit Packaging wgt.

No. per  
 container

If "Yes"  
 Package wgt.

No. per  
 container

3. Location of Net Contents Information

☐ Label ☐ Container

4. Size(s) Retail Container

Including but not limited to 2.0 oz gel per tube

5. Location of Label Directions

☐ On Label  
☐ On labeling accompanying product

6. Manner in Which Label is Affixed to Product

☐ Lithograph  
☐ Paper glued  
☐ Stenciled

☐ Other \_\_\_\_\_

**Section - IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)

Name

Timothy M. Formella

Title

Senior Registration Manager

Telephone No. (Include Area Code)

610-491-2813

**Certification**

I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete, and I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

2. Signature

3. Title

Senior Registration Manager

4. Typed Name

Timothy M. Formella

5. Date

April 21, 2009

6. Date Application  
 Received  
 (Stamped)



**United Phosphorus, Inc.**

630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406  
(610) 491-2813 (phone)  
(610) 491-2810 (fax)

April 21, 2009

VIA: UPS

Document Processing Desk (NOTIF)  
Linda Arrington  
Office of Pesticide Programs (7504P)  
U.S. Environmental Protection Agency  
Room S-4900, One Potomac Yard  
2777 South Crystal Drive  
Arlington, VA 22202-4501

**RE: Notification Regarding PR Notice 2007-4  
Kryocide Insecticide (EPA Reg. No. 70506-172)**

Dear Ms. Arrington:

United Phosphorus, Inc. hereby notifies the Agency of our intent to revise the label for Kryocide Insecticide (EPA Reg. No. 70506-172) containing the active ingredient cryolite.

This label notification has to do with revision of the Storage and Disposal text in order to comply with the requirements of PR Notice 2007-4 regarding Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment."

The changes are consistent with the notification provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46. Enclosed please find the following materials to support this registration notification:


1. Completed Application for Pesticide Registration (EPA Form 8570-1)
2. A "red-lined" copy of the amended label showing the container disposal language revisions
3. A clean copy version of the revised label
4. CD with the label in electronic format (file name: 070506-00172.20090421.pdf)
5. Completed Certification with Respect to Label Integrity form

If you have any questions on this submission please call me at 610-491-2813 or email at [tim.formella@uniphos.com](mailto:tim.formella@uniphos.com).

Sincerely,

A handwritten signature in black ink that reads "Timothy M. Formella".

Timothy M. Formella  
Senior Registration Manager



# KRYOCIDE®

## INSECTICIDE

**ACTIVE INGREDIENT**

Cryolite: sodium aluminofluoride (Fluorine-not less than 50%) ... 96.0%

OTHER INGREDIENTS ..... 4.0%

TOTAL ..... 100.0%

**KEEP OUT OF REACH OF CHILDREN**

**NOTIFICATION**

**CAUTION**

**MAY 14 2009**

**HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN OR INHALED. CAUSES MODERATE EYE IRRITATION.**

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If swallowed	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	

EPA Registration No. 70506-172

EPA Establishment No. 33770-JP-01

Net Weight 50 Lbs. (22.68 kg)

Batch/Lot # \_\_\_\_\_

UNITED PHOSPHORUS, INC.  
630 Freedom Business Center, Suite 402  
King of Prussia, PA 19406

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS (AND DOMESTIC ANIMALS)**

## CAUTION

HARMFUL IF SWALLOWED, ABSORBED THROUGH SKIN, OR INHALED. CAUSES MODERATE EYE IRRITATION. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist.

### Personal Protective Equipment:

Some materials that are chemical-resistant to this product are butyl rubber, nitrile rubber or neoprene. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- Long sleeved shirt and long pants
- Chemical resistant gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash Hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

## GENERAL INFORMATION

KRYOCIDE is a fine powder suitable for spray or dust application by ground equipment or air. A thorough application should be made, covering the underside of the leaves, if possible, as well as the top surfaces. Some processors may have additional use restrictions. Check with your processor. This product is compatible with Integrated Pest Management (IPM) Programs.

KRYOCIDE is recommended for control of the insects on the crops listed. KRYOCIDE may be used in combination with other commonly used pesticides and approved adjuvants. Thorough coverage of all foliage with Kryocide is necessary for best efficacy. It is highly recommended that every row be treated.

NOTE: Do not use KRYOCIDE in combination with lime or compounds containing free lime.

Remove visible residues on edible portions of fruit and vegetables by washing, brushing, field trimming, or other effective means.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Apply only when wind speed is 10 mph or less at the application site.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants
- Chemical resistant gloves
- Socks and chemical resistant footwear

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**Pesticide Storage:** Store in the original container in a dry area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling sweep up spillage and dispose of in accordance with the Pesticide Disposal instructions listed below.

**Pesticide Disposal:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container.

Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## DIRECTIONS FOR USE ON VEGETABLE AND FRUIT CROPS

**When used as a spray:** Apply in a minimum of 5 gallons of water by air and in a minimum of 10 gallons of water by ground or in sufficient spray volume to obtain thorough coverage.

**When used as a dust:** May be applied alone, with dusting sulfur, or other dust products. When applied in combination with sulfur or other dusting products, the label directions, and restrictions appearing on the other products labels must be adhered to such that the most restrictive of label limitations and restrictions is followed.

**When used as a spray or dust:** Apply when insects are small (early instar). Use the higher rates for severe insect infestations.

## SPRAY DRIFT LABELING

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses, or to applications using dry formulations.

1. The distance of the outer most nozzles on the boom must not exceed  $\frac{1}{4}$  the length of the wingspan or rotor.
2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they should be observed.

The applicator should be familiar with and take into account the information covered in the following Aerial Drift Reduction Advisory Information.

### Information on Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversions).

### Controlling Droplet Size

- Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.
- Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.
- Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

#### Boom Length

For some use patterns, reducing the effective boom length to less than  $\frac{1}{4}$  of the wingspan or rotor length may further reduce drift without reducing swath width.

#### Application Height

Applications should not be made at a height greater than 10 feet above the top of the largest plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

#### Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downward. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

#### Wind

Drift potential is lowest between wind speeds of 2 to 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

#### Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

#### Temperature Inversions

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

#### Sensitive Areas

The pesticide should only be applied when the potential for drift to adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g. when wind is blowing away from sensitive areas).

CROP	INSECTS	SPRAY or DUST RATE LBS/A	REMARKS
Broccoli	Cabbage Looper	8-16	Apply as needed with a minimum

CROP	INSECTS	SPRAY or DUST RATE LBS/A	REMARKS
Cauliflower Brussels sprouts Kohlrabi	Corn Earworm Cutworms Diabrotica Beetles Diamondback Moth Caterpillar Flea Beetles Imported Cabbageworm Yellowstriped Armyworm		of 7 days between applications. Do not exceed 96 lbs. per acre per season. Do not apply within 7 days of harvest.
Cabbage	Cabbage Looper Cutworms Corn Earworm Diabrotica Beetles Diamondback Moth Caterpillar Flea Beetles Imported Cabbageworm Yellowstriped Armyworm	8-16	Apply as needed with a minimum of 7 days between applications. Do not exceed 96 lbs. per acre per season. Do not apply within 14 days of harvest.
Cucumber	Cabbage Looper Cucumber Beetles Diabrotica Beetles Flea Beetles Melonworm Pickleworm	8-12	Apply as needed with a minimum of 10 days between applications. Do not exceed 48 lbs. per acre per season. Do not apply within 14 days of harvest.
Winter Squash Melons Pumpkins	Cabbage Looper Cucumber Beetles Diabrotica Beetles Flea Beetles Melonworm Pickleworm	8-16	Apply as needed with a minimum of 7 days between applications. Do not exceed 64 lbs. per acre per season. Do not apply within 14 days of harvest.
Summer Squash	Cabbage Looper Diabrotica Beetles Cucumber Beetles Flea Beetles Melonworm Pickleworm	8-16	Apply as needed with a minimum of 7 days between applications. Do not exceed 64 lbs. per acre per season. Do not apply within 7 days of harvest.
Collards	Cabbage Looper Cutworms Corn Earworm Diabrotica Beetles Diamondback Moth Caterpillar Flea Beetles Imported Cabbageworm Yellowstriped Armyworm	8-16	Apply as needed with a minimum of 10 days between applications. Do not exceed 96 lbs. per acre per season. Do not apply within 14 days of harvest.
Citrus	Citrus Cutworm Fruittree Leafroller Fuller Rose Beetle Garden Tortrix Halcocera	8-30	Apply as needed with a minimum of 30 days between applications. Do not exceed 90 lbs. per acre per season.

CROP	INSECTS	SPRAY or DUST RATE LBS/A	REMARKS
	Katydid Orange Tortrix Orangedog Variegated Cutworm Blue Green Citrus Root Weevil*		Do not apply within 15 days of harvest.
Grapes	Flea Beetles Climbing Cutworms	4-10	Apply as needed with a minimum of 14 days between applications. When needed to control later broods, make additional applications after the fruit is 1/4 inch or larger in diameter. Do not apply more than 20 lbs. per acre for preharvest use per crop year. Do not apply within 30 days of harvest. For application to grapes grown for table use after the fruit is 1/4 inch or larger in diameter, use of concentrate sprays will reduce visible spots which may result from the use of dilute applications at that stage. If used on wine grapes or grapes that may be sold to a winery for export, please observe their restrictions on post-bloom applications of KRYOCIDE.
	Omnivorous Leafroller Yellowstriped Armyworm	6-10	
	Grape Leafroller Orange Tortrix Western Grapeleaf Skeletonizer Grape Berry Moth *	5-8	
Grapevines (after harvest)	Cutworms Grape Leafroller Omnivorous Leafroller Orange Tortrix Western Grapeleaf Skeletonizer Grape Berry Moth *	5-8	For control of leaf feeding insects after all harvestable fruit or raisins have been removed from the vineyard. Apply before leaf drop while insects are actively feeding. Do not exceed 8 lbs. per acre per season for post harvest use.
Kiwi FOR USE IN CALIFORNIA ONLY	Omnivorous Leafroller	10	Apply with ground equipment only. When used as a spray use sufficient water to obtain thorough coverage, up to 200 gallons spray per acre. Apply as needed with a minimum of 15 days between applications. Do not exceed 40 lbs. per acre per season. Do not apply within 30 days of harvest.
Lettuce (leaf and head varieties)	Armyworm Cabbage Looper Corn Earworm Tobacco Budworm	8-20	Apply as needed with a minimum of 7 days between applications. Do not exceed 160 lbs. per acre per season. Do not apply within 14 days of harvest.
Peppers	Armyworm	8-12	Apply as needed with a minimum

CROP	INSECTS	SPRAY or DUST RATE LBS/A	REMARKS
	Cabbage Looper Hornworms Pepper Weevil Omnivorous Leafroller		of 7 days between applications. Do not exceed 24 lbs. per acre per crop. Do not apply within 14 days of harvest.
Potatoes	Colorado Potato Beetle  Potato Tuberworm	10-12	For spray applications: Apply by air in 5-15 gallons of water per acre or by ground in 15-100 gallons of water. Apply with a minimum of 7 days between applications. Do not apply more than 96 lbs. per acre per season. Application to exposed tubers may result in excess residues.
Tomatoes Eggplant	Armyworm Colorado Potato Beetle Larvae* Blister Beetles Cabbage Looper Flea Beetles Fruitworms Hornworms Tomato Pinworms	8-16	Apply as needed with a minimum of 7 days between applications. Do not exceed 64 lbs. per acre per season. Do not apply within 14 days of harvest.

\* Not registered for this use in California.

#### DIRECTIONS FOR USE ON ORNAMENTALS

CROP	INSECTS	SPRAY or DUST RATE LBS/A	REMARKS
Ornamentals Shade Trees	Codling Moth Flea Beetles Fuller Rose Beetle Gypsy Moth Caterpillars Katydid Plum Curculio Leafrollers	8-24	For spray applications: Apply in a minimum of 5 gallons of water by air and in a minimum of 10 gallons of water by ground. Use sufficient water volume to obtain thorough coverage of the plants. For spray or dust applications: Use the higher rate for severe insect infestations.

#### DIRECTIONS FOR USE THROUGH CHEMIGATION SYSTEMS

#### NOT FOR THIS USE IN CALIFORNIA

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### GENERAL INSTRUCTIONS

Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, or hand move irrigation systems. Do not apply this product through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Do not connect chemigation system to any public water system. Public water system means a system for the provision of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

## SYSTEM REQUIREMENTS

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

The system must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch that will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

## APPLICATION INSTRUCTIONS

Observe the requirements in the System Requirements section above.

Apply KRYOCIDE only through systems containing anti-siphon and check valves designed to prevent water source contamination or overflow of the mix tank and containing interlocking controls between the metering device and the water pump to insure simultaneous shutoff.

Maintain a gentle continuous agitation in mix tank during mixing and application to assure a uniform suspension.

Greater accuracy in calibration and distribution will be achieved by injecting a larger volume of a more dilute suspension per unit time.

Application of more than recommended quantities of irrigation water per acre may result in decreased product performance.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product cannot be flushed and must be dismantled and drained. In a center pivot system, block

the nozzle set nearest the well/pivot/injection unit to prevent spray being applied to this area.

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water.

### SPRAY PREPARATION

Remove scale, pesticide residues, and other foreign matter from the chemical tank and entire injector system. Flush with clean water. Prepare a suspension of KRYOCIDE in a mix tank. Fill the tank with 1/2 or 3/4 the desired amount of water. Start mechanical or hydraulic agitation. Slowly add the required amount of KRYOCIDE and then the remaining volume of water.

### Sprinkler Irrigation - Notes

Observe all System Requirements and Application Instructions above.

Set sprinkler system to deliver 0.1 to 1.25 inches of water per acre. Volumes of water higher than this may reduce efficacy. Start sprinkler and then uniformly inject the suspension of KRYOCIDE into the irrigation water line so as to deliver the desired rate per acre. The suspension of KRYOCIDE should be injected with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. When treatment with KRYOCIDE has been completed, do not irrigate the treated area for 24 to 48 hours to prevent washing the chemical off the crop.

For overhead chemigation, apply only when wind speed is 10 mph or less.

Where sprinkler distributed patterns do not overlap sufficiently, unacceptable pest control may result.

Check local restrictions and requirements regarding sprinkler irrigation applications, as they may vary from state to state.

### EMERGENCY TELEPHONE NUMBERS:

**CHEMTREC: (800) 424-9300**

**MEDICAL: (866) 673-6671**  
Rocky Mountain Poison Control Center

#### IMPORTANT INFORMATION READ BEFORE USING PRODUCT

#### CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

To the extent consistent with applicable law, United Phosphorus, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably

foreseeable to or beyond the control of Seller or United Phosphorus, Inc., and Buyer and User assume the risk of any such use. To the extent consistent with applicable law, UNITED PHOSPHORUS, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Rev 04/21/2009