



1/12



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY WASHINGTON, D.C. 20460

OFFICE OF PREVENTION, PESTICIDES AND TOXIC SUBSTANCES

SEP 2 5 2009

Mr. David L. Olson United Phosphorus, Inc. 630 Freedom Business Center, Suite 402 King of Prussia, PA 19406

Subject: Label Revision of the Storage and Disposal Statement

Dear Mr. Olson:

The Agency is in receipt of your Application(s) for Pesticide Notification under Pesticide Registration Notice (PRN) 2007-4 dated August 26, 2009 for:

Vendex Technical Miticide Vendex 50 WP EPA Reg. No. 70506-210 EPA Reg. No. 70506-211

The Registration Division (RD) has conducted a review of this request for applicability under PRN 2007-4 and finds that the label change(s) requested falls within the scope of PRN-2007-4. The label has been date-stamped "Notification" and will be placed in our records.

If you have any questions, at 703 305-5409 or electronically at daniel.dani@epa.gov

Sincerely,

Dani Daniel

Registration Division (7505P)

Insecticide/Rodenticide Branch

| Please read instructions on reverse before completi | na form. |)TIFIC/ | ATIO | N (m Approved | . OMB No. 2070- | 0060. Approvai | 4/12 I expires 05-31-98 |
|---|--------------------|---------------------|------------|-------------------|--------------------|------------------------|----------------------------|
| | Inited States | SEP 25 | | | | | dentifier Number |
| EPA Environmen | tal Protection | | | Amend | | | |
| | ington, DC 20460 | | | Other: | iiGiit | | |
| | Application | for Pest | | | | | |
| 1. Company/Product Number | | 2. EPA Pro | | | | 3 Propos | ed Classification |
| 70506-211 | | John Heber | | lagei | | 3. F10p0s | ed Classification |
| Company/Product (Name) United Phosphorus Inc./Vendex 50WP Mitic | | PM# 7 | | | | None | Restricted |
| 5. Name and Address of Applicant (Include ZIP (| Code) | | | | accordance w | | |
| United Phosphorus, Inc. 630 Freedom Business Center, Suite 402 | , | | produc | t is simila | r or identical ir | n compositio | on and labeling |
| King of Prussia, PA 19406 | - | to: EPA Reg. No. | | | | | |
| 3 | | El / (Nog. / | | | | | |
| Check if this is a new address | | Product Na | me | | | | |
| | | Section | - 11 | | | | |
| Amendment – Explain below. | | | Final | printed labe | els in response to | Agency letter | dated |
| Resubmission in response to Agency letter | dated | | | | | | |
| Notification - Explain below. | | [| Othe | r - Explain b | elow | | |
| Explanation: Use additional page(s) in | | | | | | | 1 |
| Notification of label change per PR Notice 20 | | | | | | | |
| requirements of EPA's regulations at 40 CFR the labeling or the Confidential Statement of | | | | | | | |
| make any false statement to EPA. I further u | inderstand that if | the amende | ed label | is not cons | sistent with the | requirements | of 40 CFR §§ |
| 156.10, 156.140, 156.144, 156.146, and 156 | | t may be in | violation | of FIFRA | and I may be s | ubject to enfo | rcement action |
| and penalties under sections 12 and 14 of FI | FRA. | | | | | | |
| | | Section | - | | | | |
| 1. Material This Product Will Be Packaged In: | | | | | | | |
| | ckaging | | | oluble Pack | aging | 2. Type of C | ontainer |
| ☐ Yes* | | | Yes | | | Metal | |
| No Sir Yes" | | o. per | If "Yes" | | No nor | ☐ Plastic | |
| Unit Do | | ontainer | Package | e wgt. | No. per container | Glass | |
| "Certification must | | | | • | | Paper | - V.J |
| be submitted 3. Location of Net Contents Information | 4. Size(s) Ret | ail Containes | | | 6 Location of | Other É | |
| Label Container | 4. 0126(3) 1(0) | an Containe | | ļ | On Label | ι. (| . (|
| | 12 x 1 lb water | soluble bags | 3 | Ì | = | ocompany | ina product |
| 6. Manner in Which Label is Affixed to Product | Lithograp | h | Ø | Other: Pa | per glued to hox | ((((| (|
| | Paper glu | ed | | | · · | ٤٠٠ (| C r |
| | Stenciled | | | | | <u>(6) (11</u> | |
| | | Section | - IV | | * (i (c i f | C. (| |
| Contact Point (Complete items directly below | | individual to | be conta | cted, if nece | | | |
| Name Devid I Olsen | Title | rector Dos | ulatore | Affaire | | Telephone No: Code) | ીંતૃહોµde Area ા ૮ ૯ |
| David L. Olson | ווט | rector, Reg | uiaiOiy | Allalis | Š | | -d01-2814 |
| | Certification | <u>"</u> | | | c r c | cc 6. Dáte | Application |
| I certify that the statements I have made on this | | | | | | .1. | èived |
| acknowledge that any knowingly false or mislea- under applicable law. | ung statement ma | y be punisnat | ne by tine | or imprisor | ment or both | 4 | Gtamped) |

3. Title

5. Date

Director, Regulatory Affairs

David L. Olson

2. Signature

4. Typed Name

Certification with Respect to Label Integrity

version: 9/11/02

I certify that the information (including, but not limited to, text, tables, and graphics) contained in the electronic files identified below by file name and submitted with this certification is the same information as that on the paper copies of these documents included with this submission.

| PROPOSED LABEL | | | | |
|----------------------|---|--------------------------|--|--|
| EPA Registration # | Date Submitted to EPA | Electronic file name | | |
| 70506-198 | 08/26/2009 | 070506-00198.20090826.pd | | |
| 70506-210 | 08/26/2009 | 070506-00210.20090826.pd | | |
| 70506-211 | 08/26/2009 070506-00211.20090826 | | | |
| 70506-212 08/26/2009 | | 070506-00212.20090826.pd | | |
| 70506-214 | 08/26/2009 | 070506-00214.20090826.pd | | |

I certify that the statements that I have made on this form are true, accurate, and complete. I acknowledge that any knowingly false or misleading statements may be punishable by fine or imprisonment or both under applicable law.

8/26/09

DAVID L. OLSON Name (typed)

DIRECTOR REGULITORY AFFAIRS



United Phosphorus, Inc.

VIA: United Parcel Service

630 Freedom Business Center, Suite 402 King of Prussia, PA 19406 (610) 491-2814 (phone) (610) 491-2810 (fax)

August 26, 2009

Document Processing Desk (NOTIF) Linda Arrington Office of Pesticide Programs (7504P) U.S. Environmental Protection Agency Room S-4900, One Potomac Yard 2777 South Crystal Drive Arlington, VA 22202-4501

RE: Notification Regarding PR Notices 2007-4

TPTH Technical EPA Reg. No. 70506-198
Vendex Technical Miticide EPA Reg. No. 70506-210
Vendex 50 WP EPA Reg. No. 70506-211
Super Tin 4L EPA Reg. No. 70506-212
Super Tin 80 WP EPA Reg. No. 70506-214

Dear Ms. Arrington:

United Phosphorus, Inc. hereby notifies the Agency of our intent to revise the subject labels for our registered pesticide products containing the active ingredients TPTH and Fenbutatin-oxide.

These label notifications have to do with revision of the Storage and Disposal text in order to comply with the requirements of PR Notice 2007-4 regarding Labeling Revisions Required by the Final Rule "Pesticide Management and Disposal; Standards for Pesticide Containers and Containment."

The changes are consistent with the notification provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46. Enclosed please find the following materials to support these registration notifications:

- 1. Completed Applications for Pesticide Registration (EPA Form 8570-1)
- 2. Copies of the revised labels
- 3. CD with the labels in electronic format (file names: 070506-00198.20090826.pdf, 070506-00210.20090826.pdf, 070506-00211.20090826.pdf, 070506-00212.20090826.pdf, and 070506-00214.20090826.pdf)
- 4. Completed Certification with Respect to Label Integrity form

If you have any questions on this submission please call me at 610-491-2814 or email at a dave.olson@uniphos.com.

Sincerely,

David L. Olson

Director, Regulatory Affairs

L Z ()

RESTRICTED USE PESTICIDE

SEP 2 5 2009

DUE TO ACUTE HUMAN TOXICITY AND VERY HIGH TOXICITY TO AQUATIC ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification. Direct supervision for this product requires the certified applicator to review federal and supplemental label instructions with all personnel prior to application, mixing, loading, or repair or cleaning of application equipment.

GROUP

12B

ACARICIDE

Vendex® 50VP

Water Soluble Pack

Miticide

| ACTIVE INGREDIENT By | Weight |
|---|--------|
| Fenbutatin-oxide[Hexakis (2-methyl-2-phenylpropyl)distannoxane] | 50% |
| OTHER INGREDIENTS | 50% |
| TOTAL | 100% |

KEEP OUT OF REACH OF CHILDREN DANGER PELIGRO



Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

| | FIRST AID |
|----------------------------|--|
| If Swallowed: | Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person. |
| If On Skin or Clothing: | Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice. |
| If Inhaled: | Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. |
| If In Eyes: | Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first five minutes, then continue rinsing. Call a poison control center or doctor for further treatment advice. |
| | uct container or label with you when calling a poison control center or doctor, or going for (reatment. Confact the in Poison Center at 1-866-673-6671 for emergency medical treatment information. |
| | rSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures agairest circulatory tory depression, and convulsion may be needed. |

See Inside for additional Precautionary Statements, and Directions For Use. [FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC at 1-800-424-9300.



Manufactured for:

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071 • www.upi-usa.com

EPA Reg. No. 70506-211 EPA Est. No. 67545-AZ-1

Net Contents: 1 lb.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER! Fatal if inhaled. Corrosive. Causes irreversible eye damage. Causes skin irritation. Harmful if swallowed or absorbed through skin. Do not get in eyes, on skin or on clothing. Do not breathe dust or spray mist. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse. Avoid handling the inner bag as moisture will cause breakage.

Do not graze or feed animals on cover crops grown in treated areas. Avoid contamination of food, feedstuffs, and domestic water supplies.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- · Coveralls over short sleeve shirt and short pants
- · Socks and chemical-resistant shoes
- Chemical-resistant gloves (category A)
- Protective eyewear (goggles, face shield, or safety glasses)
- · Chemical resistant headgear
- NIOSH approved respirator with any R, P or HE filter.
- For cleaning equipment, add a chemical-resistant apron.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROL STATEMENTS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

Water soluble packets when used correctly qualify as a closed loading system under the WPS. Handlers handling this product while it is enclosed in intact water-soluble packets are permitted to wear long-sleeved shirt, long pants, shoes plus socks, chemical-resistant gloves, and chemical-resistant apron, provided the other required PPE is immediately available in case the bag is opened.

USER SAFETY RECOMMENDATIONS

USERS SHOULD: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to birds, mammals, fish and aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift from runoff from treated areas maybe hazardous to aquatic organisms in neighboring areas.

Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves (category A)
- Shoes plus socks
- · Protective eyewear (goggles, face shield, or safety glasses)

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within a scope of the Worker Protection Standard for agricultural pesticides 40 CFR part 170. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

Do not apply this product through any type of irrigation system. Not for use in residential orchard settings.

INFORMATION

Vendex 50WP Miticide is a wettable powder contained in a water soluble packet that is used to control a wide range of herbivorous mites, including strains that are resistant to other miticides. Vendex 50WP should be dispersed in water and applied using conventional dilute sprayers or concentrate sprayers. Agitation is required during mixing and spraying.

PACKAGING

Vendex 50WP is premeasured in 1 pound Soluble Packets, which readily dissolve in water. Each Soluble Packet is contained in an individual waterproof, foil bag, with 12 bags enclosed in a cardboard box. DO NOT attempt to contained in Soluble Packets. Take the following precautions when handling vender: 50WP Soluble Packets:

- Do not handle the Soluble Packets excessively.
- Do not handle the Soluble Packets with wet hands. ' ເ
- Do not expose the Soluble Packets to water or moisture, as this will cause breakage.
- Do not place the Soluble Packets on wet surfaces.

- Do not place the foil bag in the spray tank. It is NOT soluble in water.
- · Do not sell individual water Soluble Packets.

APPLICATION RECOMMENDATIONS

To achieve the best results, apply Vendex 50WP when mites first appear or when populations reach an average of 1-2 mites per lens field. Thorough and complete coverage of infested foliage and fruit is necessary for optimum control. Vendex 50WP performs best when the daily temperature at application averages above 70° F. When the daily temperature at application averages below 70° F, performance is reduced. Vendex 50WP may be applied when honeybees and beneficial mites are present.

For best results and to improve thorough coverage, use ground or airblast application equipment to make applications of Vendex 50WP.

Apply this product only as recommended on this label.

DETERMINING GALLONAGE

To apply the correct amount of Vendex 50WP to your orchard, determine the number of gallons of water needed to spray 1 acre of your trees to the point of drip. If you have not already determined this gallonage, conduct a test to determine it. If you need assistance in calculating the proper gallonage, contact your equipment dealer or State Extension specialist.

SPRAY PREPARATION

CAREFULLY OPEN ENVELOPE AND IMMEDIATELY DROP THE INNER BAG INTO THE SPRAY TANK. DO NOT OPEN OR HANDLE THE INNER BAG. MOISTURE WILL CAUSE BREAKAGE.

Before mixing Vendex 50WP, thoroughly clean the spray tank if the spray tank has been used previously to mix boron containing or free chlorine releasing pesticide products. Even minimal carryover concentration of boron or chlorine containing products in the spray tank may cause the dissolved water soluble bag material to precipitate in the spray tank. Do not put water soluble bags close to the recirculating inlet and outlet, as they might block the line before completely dissolved.

To prepare an application of Vendex 50WP, follow these directions:

- 1. Fill a clean spray tank 1/4 to 1/2 full with water and agitate.
- 2. Add the required number of Vendex 50WP Soluble Packets (see Specific Uses for specific rate recommendations).
 - **NOTE:** To add Vendex 50WP packets, open the foil bag and drop the packet directly into water. DO NOT place the foil bag in the spray tank, as it is insoluble.
- 3. Allow the packets to dissolve completely. This should take about 5 minutes. Continue agitating the mixture to ensure that the Vendex 50WP is thoroughly mixed with the water.
- 4. Add the remaining water.

Tank Mixing

This product can be tank mixed with other pesticides provided that label directions are followed for the most restrictive of label precautions and limitations. Tank mixtures are permitted only in those states where the tank mix partner is registered.

When tank mixing Vendex 50WP with other products, introduce the products into the tank in the following order:

(1) water soluble packets (such as Vendex 50WP), (2) water dispersible granules, (3) wettable powders, (4) water based suspension concentrates, (5) water soluble concentrates, (6) oil based suspension concentrates, (7) emulsifiable concentrates, (8) adjuvants, surfactants and oils, (9) soluble fertilizers, and (10) drift retardants. Always allow each product to fully disperse before adding the next product.

Products containing boron will interfere with film solubility of the water soluble packets. If boron products are added to the spray tank, add the Vendex 50WP soluble packets first, making sure they are completely dissolved before adding any boron products.

Compatibility

Since formulations may be changed and new ones introduced, it is recommended that users check the desired tank mix and observe for possible adverse changes (settling out, flocculation, etc.) before use. Avoid mixtures of several materials and very concentrated spray mixtures.

RESISTANCE MANAGEMENT

For resistance management, Vendex 50WP is a group 12B acaricide. Repeated exclusive use of Vendex 50WP, or other group 12B acaricides, may lead to the buildup of resistant strains of mites in some crops. If more than one application of an acaricide is needed to control heavy and prolonged populations of mites, consider applying a product with a different mode of action.

Some mites are known to develop resistance to products used repeatedly for control. Because development of resistance cannot be predicted, this product may be used as part of resistance management strategies established for the use area. These strategies may include incorporation of cultural and biological control practices, alternation of mode of action classes of miticides on succeeding generations and targeting the most susceptible life stage. Consult your local or state agricultural authorities for details. If resistance to this product develops in your area, this product or other products with a similar mode of action, may not provide adequate control. If poor performance cannot be attributed to improper application or extreme weather conditions, a resistant strain of mite may be present. If you experience difficulty with control and resistance is a reasonable cause, immediately consult your local company representative or agricultural advisor for the best alternate method of control in your area. For additional information on mite and insect resistance monitoring, visit the Insecticide Resistance Action Committee (IRAC) on the web at http://www.irac-online.org.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all these factors when making application decisions. AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150 to 200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage. APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS! See Wind, Temperature and Humidity and Temperature Inversions sections of this label.

Controlling Droplet Size - General Gethniques

- Volume Use high flow rate nozzlอล to apply the highest practical spray volume. Nozzles with higher rated flows.produce larger droplets.
- Pressure Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES

ARE NEEDED, USE A HIGHER-CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using lowdrift nozzles.

Controlling Droplet Size - Aircraft

- Number of Nozzles Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- Nozzle Type Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.
- Boom Length The boom length should not exceed 3/4 of the wing or rotor length - longer booms increase drift potential.
- Application Height Application more than 10 feet above the canopy increases the potential for spray drift.

BOOM HEIGHT

Setting the boom at the lowest labeled height (if specified) which provides uniform coverage reduces the exposure of droplets to evaporation and wind. For ground equipment, the boom should remain level with the crop and have minimal bounce.

WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud.

Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

AIR ASSISTED (AIR BLAST) FIELD CROP SPRAYERS

Air assisted field crop sprayers carry droplets to the target via a downward directed air stream. Some may reduce the potential for drift, but if a sprayer is unsuitable for the application and/or set up improperly, high drift potential can result. It is the responsibility of the applicator to determine that a sprayer is suitable for

the intended application, is configured properly, and that drift is not occurring.

NOTE: Air assisted field sprayers can affect product performance by affecting spray coverage and canopy penetration. Consult the application equipment section of this label to determine if use of an air assisted sprayer is recommended.

AIR ASSISTED (AIR BLAST) TREE AND VINE SPRAYERS

Air assisted tree and vine sprayers carry droplets into the canopy of trees and vines via a radially or laterally directed air stream.

In addition to the general drift management principles already described, the following specific practices will further reduce the potential for drift:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy.
- Block off upward pointed nozzles when there is no overhanging canopy.
- Use only enough air volume to penetrate the canopy and provide good coverage.
- Do not allow spray to go beyond the edge of the cultivated area.
 Spray the outside row only from outside the planting.

SPECIFIC USE RATES

| CROP | MITES CONTROLLED | POUNDS PRODUCT PER ACRE† | MINIMUM-MAXIMUM GALLONS WATER PER ACRE | PRE-HARVEST INTERVAL (Days) | | | |
|-------------------------------------|--|--|--|--------------------------------|--|--|--|
| Apple | Apple Rust, Carpini Spider, European Red, McDaniel Spider, Two-spotted Spider | 1-2 | Max. 400 | 14 | | | |
| | | Apply when mites first appear. Do not apply more than two times per season. Apply no more than 4 pounds per acre per year. | | | | | |
| Pear | European Red, McDaniel Spider, Pear Rust, Two-spotted Spider | 1-2 | Max. 400 | 14 | | | |
| | Apply when mites first appear. Do not apply more than two times Apply no more than 4 pounds per | | | | | | |
| Grape | Pacific spider, European Red, Two-spotted Spider | 1-2.5 | Max. 250 | 28 | | | |
| | Apply when mites first appear. Make no more than two applications per season. Apply no more than 4 pounds per acre per year. Do not spray in less than 21 day intervals. | | | | | | |
| Citrus in California and Arizona | Citrus Red, Citrus Rust, Texas Citrus, Two-spotted Spider, Yuma Spider | 2-4 | Max. 1600 | 7 | | | |
| | | | | | | | |
| Citrus in Texas | Citrus Red, Citrus Rust, Texas Citrus, Two-spotted Spider, Yuma Spider | 2-3 | Max. 800 | 7 | | | |
| | Apply when mites first appear. Make no more than two applications per year. Apply no more than 6 pounds per acre per year. Do not apply in less than a 60-day interval. | | | | | | |
| Citrus in Florida | Citrus Red, Citrus Rust, Texas Citrus, Two-spotted Spider, Yuma Spider | 2-3 | Max. 800 | 7 | | | |
| | and low-metallic copper fungicides, begin applications when mite populations are less than 1 mite per lens field. Low metallic copper fungicides are defined as fungicides that contain less than 50% equivalent of metallic copper. Make no more than 1 wo applications per year. Apply no more than 4 pounds per acre per year. Apply no more than 4 pounds per acre per year. Vendex 50WP, when tank-mixed with low-metallic copper fungicides and applied when mite populations are below one mite per lens field, will provide effective residual control. Residual control with Vendex 50WP may be reduced when applied with high-metallic copper fungicides or oil. High metallic copper fungicides are defined as fungicides that contain 50% or greater equivalent of metallic copper. Do not apply in less than a 60 day interval. Ground Application Citrus groves may be planted close to bodies of water. Do not apply within 25 feet of lakes, reservoirs, rivers, permanent streams, natural ponds, marshes, or estuaries that border citrus groves. For all trees within 75 feet of bodies of water (described above), apply Vendex 50WP so that the spray is directed away from the body of water only. Shut off the sprayer when turning at row ends. Do not apply when gusts or sustained winds exceed 12 mph. Aerial Application Do not apply within 125 feet of water (described above). Do not apply winen gusts or sustained winds exceed 8 mph. Do not apply in less than 10 gallons of final spray per acre. Do not apply East of U.S. Highway #1, South and East of state road #846, or South of West Palm Beach canal. The boom length must not exceed 3/4 of the wing or rotor length (i.e. the distance of the outermost nozzles on the boom must not exceed 3/4 of the length of the wingspan or rotor). Do not apply at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Nozzles must always point backward and never be pointed downward more than 45 degrees. For best results by air: Add an ad | | | | | | |

SPECIFIC USE RATES (continued)

| CROP | MITES CONTROLLED | POUNDS PRODUCT PER ACRE† | MINIMUM-MAXIMUM GALLONS WATER PER ACRE | PRE-HARVEST INTERVAL (Days) | | | |
|---|--|--|--|--------------------------------|--|--|--|
| Peach, Plum, Prune, Nectarine | European Red McDaniel Spider, Pacific Spider, Two-spotted Spider, Big-beaked Plum, Plum Nursery | 1-2 . | Max. 400 | 14 | | | |
| | | Apply when mites first appear. Make no more than two applications per season. Apply no more than 3 pounds per acre per year. | | | | | |
| Cherry (including Sweet and Sour) | European Red, McDaniel Spider, Cherry Rust | 1.5-3 | Max. 600 | 14 | | | |
| | Apply when mites first appear. Make no more than two applications per season. Apply no more than 4.5 pounds per acre per year. | | | | | | |
| Papaya U.S. (Except California) | Carmine, Citrus Red, Texas Citrus, Broad, Red, Black Flat | 2 | Min. 100 Max. 400 | 7 | | | |
| | Apply when mites first appear. Repeat as necessary to maintain mite control, making no more than four applications per season. Apply no more than 8 pounds per acre per year. Do not apply more than once per month to bearing trees. | | | | | | |
| Almond and Pistachio | Pacific Spider, European Red, Two-spotted Spider, Hickory Spider | 1-2.5 | Max. 500 | 14 | | | |
| Pecan | Pecan, Hickory | , | | | | | |
| Walnut | Two-spotted Spider, European Red, Pacific Spider | | | | | | |
| | Apply when mites first appear. Make no more than two applications per season. Apply no more than 4 pounds per acre per year. | | | | | | |
| Strawberry | Two-spotted Spider | 1.5-3 | See below | 1 | | | |
| (California only) | Apply when mites first appear. Make no more than three applications per season. Apply no more than 9 pounds per acre per season. For good mite control, adjust spray volume and nozzle placement to assure coverage of top and underside of leaves. Apply in sufficient water to fully cover the crop; 50 to 100 gallons per acre for small plants, 150 to 300 gallons per acre for larger plant | | | | | | |
| Strawberry | Two-spotted Spider | 1.5-2 | See below | 1 | | | |
| U.S., except California | Apply when mites first appear. Make no more than two applications per season. Apply no more than 4 pounds per acre per season. For good mite control, adjust spray volume and nozzle placement to assure coverage of top and underside of leaves. Apply in sufficient water to fully cover the crop; 50 to 100 gallons per acre for small plants, 150 to 300 gallons per acre for larger | | | | | | |
| Eggplant | Clover, Two-spotted Spider | 2-3 | See below | 3 | | | |
| | Apply when mites first appear. Repeat as necessary to maintain control, making no more than three applications per year. Apply no more than 9 pounds per acre per year. For best control, thorough coverage of all leaf surfaces is required. Apply in sufficient water to fully cover the crop. | | | | | | |
| Raspberry (Black, Red) | Two-spotted Spider | 2 | Max. 200 | 3 | | | |
| in Washington and Oregon | Apply when mites first appear. Make only 1 application per year. | | | | | | |
| Christmas Trees in Washington and | Spruce Spider | 2 | Ground min. 20 Air min. 10 | NA | | | |
| Oregon | Apply when mites first appear and when the daily air temperature is expected to be at least 70°F. Make 1 application per year. For best results apply with air-blast ground equipment. Aerial application: Best results by air are obtained when the pilot applies 1/2 the dosage in one direction, then finishes by respraying in a pattern perpendicular to the first application. | | | | | | |
| † NOTE: One pound of p | product is equal to one soluble pack | et of product. | | | | | |

11/12

ORNAMENTAL USES

| CROP | MITES CONTROLLED | OUNCES PER 100 GALLONS DILUTE SPRAY | WATER SOLUBLE BAGS PER 400 GALLONS DILUTE SPRAY |
|---|--|--|---|
| Greenhouse and Outdoor Ornamentals (including nursery stock, flowers and plants grown for propagation purposes) | Spruce Spider, Oligonychus (Oak Mite, Southern Red Mite), Two-spotted Spider | 8-16 | 2-4 |
| | of visible spray residues on foliage. • Apply to the foliage only of chrysanthemu • Under greenhouse conditions, foliage a | necessary to maintain control. Frequent rep ims (pre-bloom) and poinsettias (prebract). nd flowers of certain species may demonst neral spraying. Occasional minor sensitivity is | rate sensitivity to repeat applications. If in |
| Established Landscape Ornamentals (Commercial application to established | Two-spotted Spider, Clover | 8-16 | 2-4 |
| landscape ornamentals, including trees, shrubs, flowering ornamentals, bedding plants, annuals and perennials.) | Under extreme weather conditions, foliag | necessary to maintain control. Do not apply e and flowers of certain species may demon neral spraying. Occasional minor sensitivity h | strate sensitivity to repeat applications. If in |

'Salix melanostachys (common willow), Asplenium bulbiferum (spleenwort garden fern), Dryopteris erythrosora (wood fern), Cercis canadensis (common redbud), Camellia japonica (red garden camellia), Pellaea rotundifolia (button fern), Davallia fejeensis (rabbit foot fern), Asparagus meyeri (Meyers asparagus fern), Asplenium nidus (birdnest fern), Adiantum cuneatum (maidenhair fern), Celosia argentea (cockscomb), Verbenea hortensis (verbena), Ageratum houstonianum (floss flower), Rosa sp. (common rose - some varieties, esp. yellow).

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store in a secure, dry and temperate area. Keep container closed when not in use. Do not store near food or feed. Do not use or store around the home. Avoid contact with water. In case of spill or leak, soak up with sand, earth or synthetic absorbent and dispose of wastes in compliance with local, State and Federal regulations.

PRODUCT DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available or, dispose of the empty outer foil pouch in the trash as long as WSP is unbroken.

NOTICE TO BUYER: Purchase of this material does not confer any rights under patents of countries outside of the United States.

IMPORTANT INFORMATION READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.