

70506-256

08/16/2012

1/14



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON D C 20460

OFFICE OF CHEMICAL SAFETY AND
POLLUTION PREVENTION

August 16, 2012

Sherry Hutcheson
Regulatory Affairs Manager
United Phosphorus, Inc
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

Subject Notification – Revision to Label Related to Transfer of Ownership
Phoenix Firebird Pro
EPA Registration Number 70506-256
Date of Submission July 5, 2012

Dear Ms Hutcheson

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration (PRN)98-10 dated July 5, 2012, for the above mentioned product. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the actions requested fall within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact BeWanda Alexander at (703) 305-7460 or e-mail alexander.bewanda@epa.gov

Sincerely,

A handwritten signature in black ink that reads "BeWanda Alexander for".

Richard Gebken
Product Manager Team 10
Insecticide Branch
Registration Division (7505P)

2/14

	United States Environmental Protection Agency Washington DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other Notification	OPP Identifier Number
--	--	--	-----------------------

Application for Pesticide - Section I

1 Company/Product Number 70506-256	2 EPA Product Manager Richard Gebken	3 Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4 Company/Product (Name) Firebird Pro	PM# 10	
5 Name and Address of Applicant (Include ZIP Code) United Phosphorus, Inc 630 Freedom Business Center, Suite 402 King of Prussia, PA 19406 <input type="checkbox"/> Check if this is a new address		6 Expedited Review In accordance with FIFRA Section 3(c)(3) (b)(I) my product is similar or identical in composition and labeling to EPA Reg No _____ Product Name _____

Section - II


<input type="checkbox"/> Amendment - Explain below	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> Me Too Application
<input checked="" type="checkbox"/> Notification Explain below	<input type="checkbox"/> Other Explain below

Explanation Use additional page(s) if necessary (For Section I and Section II)
 Changes to label related to EPA transfer of ownership
 This notification is consistent with the provisions of PR Notice 98 10 and EPA regulations at 40 CFR 152 46, and no other changes have been made to the labeling or the confidential statement of formula of this product I understand that it is a violation of 18 U S C Sec 1001 to willfully make any false statement to EPA I further understand that if this notification is not consistent with the terms of PR Notice 98 10 and 40 CFR 152 46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA

Section - III

1 Material This Product Will Be Packaged In			
Child Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If Yes No per Unit Packaging wgt container	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If Yes No per Package wgt container	2 Type of Container <input type="checkbox"/> Metal <input checked="" type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
* Certification must be submitted			
3 Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4 Size(s) Retail Container 1 gallon	5 Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6 Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

Section - IV

1 Contact Point (Complete items directly below for identification of individual to be contacted if necessary to process this application)		
Name Sherry Hutcheson	Title Regulatory Affairs Manager	Telephone # (Include Area Code) 229 247 0041
I certify that the statements I have made on this form and all attachments thereto are true accurate and complete and I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law		6 Date Application Received <div style="text-align: center; border: 1px solid black; padding: 5px;"> (Stamped) </div>
2 Signature 	3 Title Regulatory Affairs Manager	
4 Typed Name Sherry Hutcheson	5 Date 07/05/2012	

4/14

RESTRICTED USE PESTICIDE

Toxic to fish and aquatic organisms

For retail sale to and use only by certified applicators or persons under their direct supervision and only for the uses covered by the certified applicator's license



NOTIFICATION

AUG 16 2012

Firebird[®] Pro^{INSECTICIDE}

For Commercial Non-Food Use on Indoor and Outdoor Ornamentals, Greenhouses, Nurseries, Turf on Golf Courses, Sod Farms, and on ornamentals and lawns in landscaped areas around residential, institutional, public, commercial and industrial buildings, parks, recreational areas, golf courses, sod farms and athletic fields

DO NOT USE THIS PRODUCT ON GOLF COURSES IN NASSAU COUNTY OR SUFFOLK COUNTY NEW YORK NOT FOR USE ON SOD FARMS IN THE STATE OF NEW YORK

Active Ingredient
Bifenthrin
Other Ingredients
TOTAL

By Wt
7.9%
92.1%
100.0%

Phoenix Firebird Pro contains 2/3 pound active ingredient per gallon
Cis isomers 97% minimum trans isomers 3% maximum

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

IF SWALLOWED	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice • Have person sip a glass of water if able to swallow • Do not induce vomiting unless told to do so by the poison control center or doctor • Do not give anything by mouth to an unconscious person
IF INHALED	<ul style="list-style-type: none"> • Move person to fresh air • If person is not breathing call 911 or an ambulance then give artificial respiration preferably by mouth to mouth if possible • Call a poison control center or doctor for further treatment advice
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 20 minutes • Call a poison control center or doctor for treatment advice
IF IN EYES	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 20 minutes • Remove contact lenses if present after the first 5 minutes then continue rinsing eye • Call a poison control center or doctor for treatment advice

Have the product container or label with you when calling a poison control center or doctor or going for treatment **For medical emergency call the Rocky Mountain Poison Control Center at 1 866 673 6671**

NOTE TO PHYSICIAN

This product is a pyrethroid. If large amounts have been ingested the stomach and intestine should be evacuated. Treatment is symptomatic and supportive. Digestible fats, oils, or alcohol may increase absorption and so should be avoided.

See inside for additional Precautionary Statements, Directions For Use, and Storage and Disposal

FOR CHEMICAL EMERGENCY Spill, leak, fire, exposure or accident call CHEMTREC 1 800 424 9300

United Phosphorus Inc
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406 • 1 800 438 6071

EPA Reg. No. 70506 256

5/14

PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed inhaled or absorbed through skin Avoid contact with skin eyes or clothing Avoid breathing spray mist Wash thoroughly with soap and water after handling

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below If you want more options follow the instructions for category C on an EPA chemical resistance category selection chart

Applicators and other handlers (other than mixers and loaders) must wear

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate butyl rubber nitrile rubber neoprene rubber polyvinyl chloride or Viton
- Shoes plus socks

Mixers and Loaders must wear

- Long sleeved shirt and long pants
- Chemical resistant gloves such as barrier laminate butyl rubber nitrile rubber neoprene rubber polyvinyl chloride or Viton
- Shoes plus socks
- Protective eyewear

Follow manufacturer s instructions for cleaning/maintaining PPE If no such instructions for washables use detergent and hot water Keep and wash PPE separately from other laundry

USER SAFETY RECOMMENDATIONS

Users should

- Wash hands before eating drinking chewing gum using tobacco or using the toilet
- Remove clothing immediately if pesticide gets inside Then wash thoroughly and put on clean clothing
- Remove PPE immediately after handling this product Wash outside of gloves before removing As soon as possible wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates Drift and run off from treated areas may be hazardous to aquatic organisms in neighboring areas To protect the environment do not allow pesticide to enter or run off into storm drains drainage ditches gutters or surface waters Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds Do not apply this product or allow to drift to blooming crops if bees are visiting the treatment area

DIRECTIONS FOR USE

Restricted Use Pesticide

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling Do not apply this product in a way that will contact workers or other persons either directly or through drift Only protected handlers may be in the area during application For any requirements specific to your State or Tribe consult the agency responsible for pesticide regulation

Do not apply this product through any kind of irrigation system

Do not water the treated area to the point of run off

Do not make applications during rain

Do not apply by air

All outdoor applications must be limited to spot or crack and crevice treatments only except for the following permitted uses

- 1) Treatments to soil or vegetation around structures
- 2) Applications to lawns turf and other vegetation
- 3) Applications to building foundations up to a maximum height of 3 feet

Other than applications to building foundations all outdoor applications to impervious surfaces such as sidewalks driveways patios porches structural surfaces (such as windows doors and eaves) are limited to spot and crack and crevice applications only

Application is prohibited directly into sewers or drains or to any area like a gutter where drainage to sewers storm drains water bodies or aquatic habitat can occur Do not allow the product to enter any drain during or after application

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard (WPS) 40 CFR Part 170 This Standard contains requirements for the protection of agricultural workers on farms forests nurseries and greenhouses and handlers of agricultural pesticides It contains requirements for training decontamination notification and emergency assistance It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard

Do not enter or allow worker entry into treated areas during restricted entry interval (REI) of 12 hours

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants soil or water is

- Coveralls
- Chemical resistant gloves such as barrier laminate or butyl rubber nitrile rubber or neoprene rubber or polyvinyl chloride or Viton
- Shoes plus socks

6/14

California Specific Requirements for Greenhouse Applicators and Harvesters

In addition to following all applicable precautionary statements the following is required for greenhouse applicators and harvesters

Greenhouse Applicator Greenhouse applicators must wear a full body chemical resistant protective suit (such as barrier laminate butyl rubber nitrile rubber polyvinyl chloride or equivalent)

Reapplication Interval Reapplications to greenhouses must be at intervals of 30 days or longer

Greenhouse Harvesters Greenhouse harvesters must wear regular length gloves plus a long sleeved shirt or elbow length (gauntlet type) gloves during the 30 days following application

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170) The WPS applies when this product is used to produce agricultural plants on farms forests nurseries and greenhouses

Do not allow people or pets on treated surfaces until the spray has dried

Do not touch treated surface until dry

STORAGE AND DISPOSAL

Prohibitions Do not contaminate water food or feed by storage or disposal

Pesticide Storage Keep out of reach of children and animals Store in original containers only Store in a cool dry place and avoid excess heat Carefully open containers After partial use replace lids and close tightly Do not put concentrate or dilute material into food or drink container

In case of spill avoid contact isolate area and keep out animals and unprotected persons Confine spills

To Confine Spill If liquid dike surrounding area or absorb with sand cat litter or commercial clay If dry material cover to prevent dispersal Place damaged package in a holding container Identify contents

Pesticide Disposal Pesticide wastes are toxic Do not contaminate water food or feed by storage or disposal Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal Law Dispose of excess or waste pesticide by use according to label directions or contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

Container Handling Nonrefillable container Do not reuse or refill this container Clean container promptly after emptying

Triple rinse as follows Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip Fill the container 1/4 full with water and recap Shake for 10 seconds Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal Drain for 10 seconds after the flow begins to drip Repeat this procedure two more times Then offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration or if allowed by state and local authorities by burning If burned stay out of smoke

GENERAL APPLICATION INSTRUCTIONS

Phoenix Firebird Pro formulation mixes readily with water and other aqueous carriers and controls listed insects and mites on ornamentals trees shrubs foliage plants non bearing fruit and nut trees and flowers in greenhouses and outdoor nurseries and interiorscapes including hotels shopping malls office buildings etc and outdoor plantscapes such as around residential dwellings parks institutional public commercial and industrial buildings recreational athletic fields home lawns golf courses and sod farms Non bearing crops are perennial crops that will not produce a harvestable raw agricultural commodity during the season of application

Phoenix Firebird Pro may be tank mixed with other products including insect growth regulators When tank mixing Phoenix Firebird Pro with other products observe all precautions and limitations on each separate product label The addition of spreader stickers is not necessary The physical compatibility of Phoenix Firebird Pro may vary with different sources of pesticide products and local cultural practices Any tank mixture which has not been previously tested should be prepared on a small scale (pint or quart jar) using the proper proportions of chemicals and water to ensure the physical compatibility of the mixture

The following procedure is for preparation of a new tank mix unless specified otherwise in label directions (1) Add wettable powders to tank water (2) Agitate (3) Add liquids and flowables (4) Agitate (5) Add emulsifiable concentrates and (6) Agitate If a mixture is found to be incompatible following this order of addition try reversing the order of addition or increase the volume of water **Note** If the tank mixture is found to be compatible after increasing the amount of water then the sprayer will need to be recalibrated for a higher volume application Do not allow tank mix to stand overnight

Do not apply more than 0.1 lb a.i./acre (20 fl. ozs. of Phoenix Firebird Pro) in a single application and no more than 0.2 lb a.i./acre/year (40 fl. ozs.) for outdoor applications

Note For large infestations of ants imported fire ants and mole crickets a single application of 0.2 lb a.i./acre (40 fl. ozs.) may be applied once per year

Resistance Some insects are known to develop resistance to products used repeatedly for control Because the development of resistance cannot be predicted the use of this product should conform to resistance management strategies established for the use area Consult your local or state pest management authorities for details

If resistance to this product develops in your area this product or other products with a similar mode of action may not provide adequate control If poor performance cannot be attributed to improper application or extreme weather conditions a resistant strain of insect may be present If you experience difficulty with control and suspect that resistance is a reasonable cause immediately consult your local company representative or pest management advisor for the best alternative method of control for your area

7/14

Phoenix Firebird Pro Dilution Chart for Turf and Ornamental Uses

• Rate in amount of Active Ingredient per acre
Application Volume in Gallons per acre

Application Volume Gallons Per Acre	Application Rate lb a i /Acre	Fluid Ounces* of Phoenix Firebird Pro Diluted to These Volumes of Finished Spray			
		1 Gallon	25 Gallons	50 Gallons	100 Gallons
50	0 025	0 1	2 5	5 0	10 0
50	0 05	0 2	5 0	10 0	20 0
50	0 1	0 4	10 0	20 0	40 0
50	0 2	0 8	20 0	40 0	80 0
100	0 025	0 05	1 25	2 5	5 0
100	0 05	0 1	2 5	5 0	10 0
100	0 1	0 2	5 0	10 0	20 0
100	0 2	0 4	10 0	20 0	40 0
150	0 025	0 03	0 83	1 67	3 3
150	0 05	0 07	1 67	3 33	6 7
150	0 1	0 13	3 33	6 67	13 3
150	0 2	0 27	6 67	13 33	26 7
200	0 025	0 025	0 63	1 25	2 5
200	0 05	0 05	1 25	2 5	5 0
200	0 1	0 1	2 5	5 0	10 0
200	0 2	0 2	5 0	10 0	20 0
250	0 025	—	0 5	1 0	2 0
250	0 05	—	1 0	2 0	4 0
250	0 1	—	2 0	4 0	8 0
250	0 2	—	4 0	8 0	16 0
300	0 025	—	0 42	0 83	1 7
300	0 05	—	0 83	1 67	3 3
300	0 1	—	1 67	3 33	6 7
300	0 2	—	3 33	6 67	13 3

To convert to milliliters multiply by 29 57
1 fluid oz = 29 57 ml = 2 tablespoons = 6 teaspoons Do not use household utensils to measure Phoenix Firebird Pro

Phoenix Firebird Pro Dilution Chart for Turf and Ornamental Uses

Rate in amount of Active Ingredient per acre
Application Volume in Gallons per 1 000 Sq Ft

Application Volume Gallons Per 1 000 Sq Ft	Application Rate lb a i /Acre	Fluid Ounces of Phoenix Firebird Pro Diluted to These Volumes of Finished Spray				
		1 Gallon	10 Gallons	25 Gallons	50 Gallons	100 Gallons
0 5	0 05	0 5	5 0	12 5	25 0	50 0
0 5	0 2	0 1	10 0	25 0	50 0	100 0
0 5	0 1	0 2	20 0	50 0	100 0	200 0
0 75	0 05	0 33	3 33	8 33	16 67	33 33
0 75	0 1	0 67	6 67	16 67	33 33	66 67
0 75	0 2	1 33	13 33	33 33	66 67	133 33
1	0 05	0 25	2 5	6 25	12 5	25 0
1	0 1	0 5	5 0	12 5	25 0	50 0
1	0 2	1	10 0	25 0	50 0	100 0
1 5	0 05	0 17	1 67	4 17	8 33	16 67
1 5	0 1	0 33	3 33	8 33	16 67	33 33
1 5	0 2	0 67	6 67	16 67	33 33	66 67
2	0 05	0 13	1 25	3 13	6 25	12 5
2	0 1	0 25	2 5	6 25	12 5	25 0
2	0 2	0 5	5 0	12 5	25 0	50 0
2 3	0 05	0 11	1 09	2 72	5 43	10 87
2 3 ^a	0 1	0 22	2 17	5 43	10 87	21 74
2 3 ^a	0 2	0 43	4 35	10 87	21 74	43 48
3	0 05	—	0 83	2 09	4 17	8 33
3	0 1	0 17	1 66	4 17	8 33	16 67
3	0 2	0 33	3 33	8 33	16 67	33 33
4	0 05	—	0 63	1 56	3 13	6 25
4	0 1	0 13	1 25	3 13	6 25	12 5
4	0 2	0 25	2 5	6 25	12 5	25 0

To convert to milliliters multiply by 29 57
^a100 gallons per acre
1 fluid oz = 29 57 ml = 2 tablespoons = 6 teaspoons Do not use household utensils to measure Phoenix Firebird Pro

8/14

Phoenix Firebird Pro Dilution Chart for Turf and Ornamental Uses

- Rate in Fluid Ounces of Product per 1 000 Sq Ft
- Application Volume in Gallons per 1 000 Sq Ft

Application Volume Gallons Per 1 000 Sq Ft	Application Rate Fluid Ounces per 1 000 Sq Ft	Fluid Ounces of Phoenix Firebird Pro Diluted to These Volumes of Finished Spray			
		1 Gallon	5 Gallons	10 Gallons	100 Gallons
1 0	0 125	0 125	0 63	1 25	12 5
1 0	0 25	0 25	1 25	2 5	25 0
1 0	0 5	0 5	2 5	5 0	50 0
1 0	1 0	1 0	5 0	10 0	100 0
2 0	0 125	—	0 31	0 63	6 3
2 0	0 25	0 13	0 63	1 25	12 5
2 0	0 5	0 25	1 25	2 5	25 0
2 0	1 0	0 5	2 5	5 0	50 0
2 3 ^a	0 125	—	0 27	0 54	5 4
2 3	0 25	0 11	0 54	1 08	10 8
2 3 ^a	0 5	0 22	1 09	2 17	21 7
2 3	1 0	0 44	2 17	4 35	43 5
3 0	0 125	—	0 21	0 42	4 2
3 0	0 25	—	0 42	0 83	8 3
3 0	0 5	0 17	0 83	1 67	16 7
3 0	1 0	0 33	1 67	3 33	33 3
4 0	0 125	—	0 15	0 31	3 1
4 0	0 25	—	0 31	0 63	6 3
4 0	0 5	0 13	0 63	1 25	12 5
4 0	1 0	0 25	1 25	2 5	25 0
4 6 ^b	0 125	—	0 14	0 27	2 7
4 6 ^b	0 25	—	0 27	0 54	5 4
4 6 ^b	0 5	0 11	0 54	1 09	10 9
4 6 ^b	1 0	0 22	1 09	2 17	21 7
5 0	0 125	—	0 13	0 25	2 5
5 0	0 25	—	0 25	0 5	5 0
5 0	0 5	0 1	0 5	1 0	10 0
5 0	1 0	0 2	1 0	2 0	20 0
6 9 ^c	0 125	—	—	0 18	1 8
6 9 ^c	0 25	—	0 18	0 36	3 6
6 9	0 5	—	0 36	0 72	7 2
6 9 ^c	1 0	0 15	0 72	1 45	14 5
10 0	0 125	—	—	0 13	1 3
10 0	0 25	—	0 13	0 25	2 5
10 0	0 5	—	0 25	0 5	5 0
10 0	1 0	0 1	0 5	1 0	10 0

To convert to milliliters multiply by 29 57
^a100 gallons per acre
^b200 gallons per acre
^c300 gallons per acre
 Do not use household utensils to measure Phoenix Firebird Pro

Formula for Determining the Active Ingredient Content of the Finished Spray Mixture The following formula may be used to determine the percent active ingredient that is in the spray tank after mixing Phoenix Firebird Pro

$$\frac{(7.9)(\text{Fl Oz of Phoenix Firebird Pro added to tank})}{(\text{Gallons of finished spray mix})(128)} = \text{Percent Active Ingredient of spray mix}$$

ORNAMENTALS IN GREENHOUSES, LATH HOUSES, SHADE HOUSES AND OUTDOOR NURSERIES, INCLUDING NON-BEARING FRUIT AND NUT TREES

APPLICATION

Apply 0 025 to 0 2 lb a /A (5 to 40 fl ozs) or 0 125 to 1 0 fluid ounce per 1 000 square feet of Phoenix Firebird Pro Phoenix Firebird Pro may be diluted and applied in various volumes of water providing that the maximum label rate (0 2 lb a /A or 40 fl ozs 1 0 fluid ounce per 1 000 square feet) is not exceeded (refer to Dilution Chart for specific instructions) Phoenix Firebird Pro may be applied through low volume application equipment by dilution with water or other carriers and providing that the maximum label rate (0 2 lb a /A or 40 fl ozs 1 0 fluid ounce per 1 000 square feet) is not exceeded

ORNAMENTAL APPLICATION RATES

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions. However, at the discretion of the applicator, Phoenix Firebird Pro may be applied at up to 0.2 lb a.i./A (40 fl. ozs.) or 1 fluid ounce per 1,000 square feet to control each of the pests listed in this Table.

Application Rate of Phoenix Firebird Pro				
Pest	Fluid Ounces per 1,000 Square Feet	lb a.i./Acre	Fluid Ounces per Acre	Other Use Precautions and Restrictions
Aphids Bagworms ¹ Cutworms Elm Leaf Beetles Fall Webworms Lace Bugs Leaf Feeding Caterpillars Plant Bugs (including Lygus spp.) Tent Caterpillars	0.125 - 0.25	0.025 - 0.05	5 - 10	Certain cultivars may be sensitive to the final spray solution. A small number of plants should be treated and observed for one week prior to application to the entire planting. Apply with ground equipment only. Do not apply when wind direction favors downwind drift towards near by water bodies. Do not apply when wind velocity exceeds 10 mph. Avoid application when wind gusts approach 10 mph. Do not apply when a temperature inversion exists.
Beet Armyworm Black Vine Weevil (Adults) Broad Mites Brown Soft Scales Budworms California Red Scale (Crawlers) ² Centipedes Citrus Thrips Clover Mites Crickets Diaprepes (Adults) Earwigs European Red Mite Flea Beetles Fungus Gnats (Adults) Grasshoppers Gypsy Moth Caterpillars Leafhoppers Leafrollers Mealybugs Millipedes Mites Orchid Weevil Pillbugs Pine Needle Scales (Crawlers) ² San Jose Scales (Crawlers) ² Sowbugs Spider Mites Spiders Thrips Tip Moths Twig Borers ² Wasps Weevils Whiteflies	0.25 - 0.5	0.05 - 0.1	10 - 20	Apply the specified application rate as a full coverage foliar spray. Typical application volume is 100 gallons per acre. Repeat treatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity. Repeat applications are permitted provided that the maximum label rate is not exceeded. Apply using the largest nozzle size compatible with adequate coverage. Do not apply if rain is expected within 12 hours (or whatever time is necessary for the spray to dry). Do not apply by ground equipment within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds. When treating tall trees (>15 feet) from the ground with high pressure sprays or during any application with air assisted equipment (mist blower) do not apply within 150 feet of aquatic areas. Use of an alternate class of chemistry in a treatment program is recommended to prevent or delay pest resistance.
Ants Imported Fire Ants Japanese Beetle (Adults) Leafminers Pecan Leaf Scorch Mite Pine Shoot Beetle (Adults)	0.5 - 1.0	0.1 - 0.2	20 - 40	
¹ Bagworms Apply when larvae begin to hatch and spray larvae directly. Applications when larvae are young will be most effective. ² Scale Crawlers and Twig Borers Treat trunks, stems and twigs in addition to plant foliage. For foraging ants				

10/14

TURF (GOLF COURSES AND SOD FARMS) AND GRASS AREAS (INCLUDING GOLF COURSES, SOD FARMS, HOME LAWNS, LAWN AREAS AROUND PARKS, INSTITUTIONAL, PUBLIC, COMMERCIAL AND INDUSTRIAL BUILDINGS, RECREATIONAL AND ATHLETIC FIELDS)

APPLICATION

NOT FOR USE ON SOD FARMS IN THE STATE OF NEW YORK

Apply Phoenix Firebird Pro as a surface or sub surface treatment Use application volumes of up to 10 gallons per 1 000 square feet to get uniform coverage when treating dense and or long turf foliage

For low volume applications less than 2 gallons/1 000 square feet immediate irrigation of treated area with at least 0 25 inches of water following application to ensure efficacy of listed sub surface pests including Mole Crickets

In New York State this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake pond river stream wetland or drainage ditch)

In New York State do make a single repeat application of Phoenix Firebird Pro if there are signs of renewed insect activity but not sooner than two weeks after the first application

APPLICATION RATES

The application rates listed in the following table will provide excellent control of the respective pests under typical conditions However at the discretion of the applicator¹ Phoenix Firebird Pro may be applied at up to 0 1 lb a /A (20 fl ozs) to control each of the pests listed in this Table (0 2 lb a /A or 40 fl ozs of Phoenix Firebird Pro for ants imported fire ants and mole crickets)¹³

¹During periods of high pest pressure or for maximum residual control

Pest	lb Active Ingredient/Acre	Application Rate of Phoenix Firebird Pro		Other Use Precautions and Restrictions
		per acre	per 1 000 sq ft	
Armyworms ³ Cutworms ³ Sod Webworm ³	0 05	10 fl oz	0 25 fl oz	Apply with ground application equipment only (and apply with nozzles not more than two feet above the turf) Do not apply when wind conditions favor downwind drift towards near by water bodies Do not apply when wind velocity exceeds 10 mph Avoid application when wind gusts approach 10 mph Do not apply when a temperature inversion exists Apply using the largest droplet size compatible with adequate coverage Do not apply for surface feeding pests if rain is expected within 12 hours (or whatever time is necessary for the spray to dry) Do not apply by ground equipment within 25 feet of lakes reservoirs rivers permanent streams marshes or natural ponds estuaries and commercial fish farm ponds Do not apply when turf areas are water logged or soil is saturated with water (i e will not accept irrigation)
Annual Bluegrass Weevil (<i>Hyperodes</i>) (Adult) ⁴ Ants Billbugs (Adult) ⁵ Black Turfgrass Ataenius (Adult) ⁶ Centipedes Chinch Bugs ⁷ Crickets Earwigs Fleas (Adult) Grasshoppers Leafhoppers Mealybugs Millipedes Mites ⁸ Mole Cricket (Adult) ⁹ Mole Cricket (Nymph) ¹⁰ Pillbugs Sowbugs	0 05 - 0 1	10 - 20 fl oz	0 25 - 0 5 fl oz	
Fleas (Larvae) ¹¹ Imported Fire Ants Japanese Beetle (Adult) Ticks ¹²	0 1	20 fl oz	0 5 fl oz	
Ants Imported Fire Ants ¹⁴ Mole Crickets	0 2 ¹³	40 ¹³ fl oz	1 ¹³ fl oz	

Footnotes

³**Armyworms Cutworms and Sod Webworms** To ensure optimum control delay watering (irrigation) or mowing for 24 hours after application If the grass area is being maintained at a mowing height of greater than 1 inch then higher application rates (up to 0 1 lb a /A or 20 fl ozs of Phoenix Firebird Pro) may be required during periods of high pest pressure

⁴**Annual Bluegrass Weevil (*Hyperodes*) adults** Applications should be timed to control adult weevils as they leave their overwintering sites and move into grass areas This movement generally begins when *Forsythia* is in full bloom and concludes when flowering dogwood (*Cornus florida*) is in full bloom Consult your State Cooperative Extension Service for more specific information regarding application timing

⁵**Billbug adults** Applications should be made when adult billbugs are first observed during April and May Degree day models have been developed to optimize application timing Consult your State Cooperative Extension Service for information specific to your region In temperate regions spring applications targeting billbug adults will also provide control of over wintered chinch bugs

⁶**Black Turfgrass Ataenius adults** Applications should be made during May and July to control the first and second generation of black turfgrass ataenius adults respectively The May application should be timed to coincide with the full bloom stage of Vanhoutte spiraea (*Spiraea vanhouttei*) and horse chestnut (*Aesculus hippocastanum*) The July application should be timed to coincide with the blooming of Rose of Sharon (*Hibiscus syriacus*)

⁷**Chinch Bugs** Chinch Bugs infest the base of grass plants and are often found in the thatch layer Irrigation of the grass area before treatment will optimize the penetration of the insecticide to the area where the chinch bugs are located Use higher volume applications if the thatch layer is excessive or if a relatively long mowing height is being maintained Chinch Bugs can be one of the most difficult pests to control in grasses and the higher application rates (up to 0 1 lb a /A or 20 fl ozs) may be required to control populations that contain both nymphs and adults during the middle of the summer

⁸**Mites** To ensure optimal control of eriophyid mites apply in combination with the labeled application rate of a surfactant A second application five to seven days after the first may be necessary to achieve acceptable control

11/14

9Mole Cricket adults Achieving acceptable control of adult mole crickets is difficult because preferred grass areas are subject to continuous invasion during the early spring by this extremely active stage. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized. Grass areas that receive pressure from adult mole crickets should be treated at peak egg hatch to ensure optimum control of subsequent nymph populations (see below).

10Mole Cricket nymphs Grass areas that received intense adult mole cricket pressure in the spring should be treated immediately prior to peak egg hatch. Optimal control is achieved at this time because young nymphs are more susceptible to insecticides and they are located near the soil surface where the insecticide is most concentrated. Control of larger, more damaging nymphs later in the year may require both higher application rates and more frequent applications to maintain acceptable control. Applications should be made as late in the day as possible and should be watered in with up to 0.5 inches of water immediately after treatment. If the soil is not moist, then it is important to irrigate before application to bring the mole crickets closer to the soil surface where contact with the insecticide will be maximized.

11Flea larvae Flea larvae develop in the soil of shaded areas that are accessible to pets or other animals. Use a higher volume application when treating these areas to ensure penetration of the insecticide into the soil. Note: if the lawn area is being treated with Phoenix Firebird Pro at 0.05 lb a.i./A (10 fl. ozs) for adult flea control, then the larval application rate may be achieved by doubling the application volume.

12Ticks (including ticks that may transmit Lyme Disease and Rocky Mountain Spotted Fever) Do not make spot applications. Treat the entire area where exposure to ticks may occur. Use higher spray volumes when treating areas with dense ground cover or heavy leaf litter. Ticks may be reintroduced from surrounding areas on host animals. Retreatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed activity. Repeat application must be limited to no more than once per seven days. Do not allow public use of treated areas during application or until sprays have dried.

Deer ticks (Ixodes spp.) have a complicated life cycle that ranges over a two-year period and involves four life stages. Applications should be made in the late fall and/or early spring to control adult ticks that are usually located on brush or grass above the soil surface and in mid to late spring to control larvae and nymphs that reside in the soil and leaf litter.

American dog ticks may be a considerable nuisance in suburban settings, particularly where homes are built on land that was previously field or forest. These ticks commonly congregate along paths or roadways where humans are likely to be encountered. Applications should be made as necessary from mid-spring to early fall to control American dog tick larvae, nymphs, and adults.

13Note: For large infestations of ants, imported fire ants and mole crickets, a single application of 0.2 lb a.i./A (40 fl. ozs of Phoenix Firebird Pro) may be applied once per year.

14Imported Fire Ants Control will be optimized by combining broadcast applications that will control foraging workers and newly mated fly-in queens with mound drenches that will control existing colonies. If the soil is not moist, then it is important to irrigate before application or use a high volume application. Broadcast treatments should apply 0.2 lb a.i./A (40 fl. ozs of Phoenix Firebird Pro). Mounds should be treated by diluting 1 teaspoon of Phoenix Firebird Pro Insecticide per gallon of water and applying 1 to 2 gallons of finished spray per mound. The mounds should be treated with sufficient force to break their apex and allow the insecticide solution to flow into the ant tunnels. A four-foot diameter circle around the mound should also be treated. For best results, apply in cool weather (65–80 °F) or in early morning or late evening hours. Note: a spray rig that is calibrated to apply 0.2 lb a.i./A (40 fl. ozs) of Phoenix Firebird Pro in 5 gallons per 1,000 square feet contains the approximate dilution (1 teaspoon per gallon) that is required for fire ant mound drenches in the spray tank.

IMPORTED FIRE ANT QUARANTINE TREATMENT

Against Imported Fire Ants (IFA) in Potting Media (including balled and containerized nursery grown ornamental trees, shrubs, plants, flowers, conifers, bushes, Christmas trees, and non-bearing fruit and nut trees), Phoenix Firebird Pro can be used in accordance with the USDA Imported Fire Ant Quarantine Program. Phoenix Firebird Pro may be applied either soil incorporated, as a topical application, or as a high volume drench treatment.

Soil Incorporation: Incorporate the appropriate volume of Phoenix Firebird Pro (see table below) per cubic yard of potting media by diluting it in water (typically 1 quart to 1 gallon per cubic yard of media) and sprinkling or spraying it onto the media. The applications are based on the dry bulk density of the potting media. When used in accordance with USDA guidelines, this application will provide a 6-month certification period.

Soil Incorporation Rate of Phoenix Firebird Pro for Control of IFA in Potting Media

Potting Media Bulk Density (lbs per cubic yard)	Fluid ounces of Phoenix Firebird Pro in one cubic yard
200	1.9
400	3.8
600	5.7
800	7.6
1000	9.5
1200	11.4
1400	13.3

Use proportional amounts of Phoenix Firebird Pro for potting media with bulk densities not listed.

Topical Application: Mix Phoenix Firebird Pro in 1,000 ounces of water based on container size and bulk density of the potting media (see table below). Apply one (1) ounce of the mix to each container evenly distributed over the surface of the potting media. Irrigate all treated containers with 1.5 inches of water following application. When used in accordance with USDA guidelines, this application will provide a 6-month certification period.

Topical Drench Application Rate of Phoenix Firebird Pro for Control of IFA in Potting Media

Potting Media Bulk Density (lbs per cubic yard)	Fluid ounces of Phoenix Firebird Pro per 1,000 ounces of water	
	3 Qt. Container	4 Qt. Container
200	3.6	5.2
400	7.2	10.4
600	10.8	15.6
800	14.4	20.8
1000	18.0	26.0
1200	21.6	31.2
1400	25.2	36.4

Use proportional amounts of Phoenix Firebird Pro for potting media with bulk densities not listed.

High Volume Drench Apply Phoenix Firebird Pro as a high volume drench by mixing the appropriate amount of product based on the bulk density in 100 gallons of water (see table below) Apply mix to individual containers to the point of saturation The amount of mix used for each plant is generally 1/5 the volume of the container When used in accordance with USDA guidelines this application will provide a 6 month certification period

High Drench Application Rate of Phoenix Firebird Pro for Control of IFA in Potting Media

Potting Media Bulk Density (lbs per cubic yard)	Fluid ounces of Phoenix Firebird Pro in 100 gallons
200	2.4
400	4.8
600	7.2
800	9.6
1000	12.0
1200	14.4
1400	16.8

Use proportional amounts of Fluid ounces of Phoenix Firebird Pro for potting media with bulk densities not listed

LARVAL CONTROL IN POTTING MEDIA OF CONTAINERIZED PLANTS

Black Vine Weevil Larval Control – Preventative Treatment – Topical Drench For preventative control of black vine weevil larvae in containerized plants dilute Phoenix Firebird Pro at the rate of 10 to 40 fl ozs (0.05 to 0.2 lb a.i.) per 100 gallons and apply as a drench at the rate of 4 to 8 fluid ounces of finished spray per 6 inch (diameter) container Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter Ideally the media should be treated to the point of saturation which generally requires 1/5 the volume of the container Diluting 10 fluid ounces of Phoenix Firebird Pro per 100 gallons and applying 8 fluid ounces of finished spray per 6 inch (diameter) container will provide black vine weevil larval control for one growing season when the application is made in the spring Diluting 20 to 40 fluid ounces of Phoenix Firebird Pro per 100 gallons and applying 8 fluid ounces of finished spray per 6 inch (diameter) container will provide black vine weevil larval control for two growing seasons when the application is made in the spring

White Grub Control – Preventative Treatment – Topical Drench For preventative control of white grubs (including but not limited to Japanese beetle, oriental beetle and European chafer) in containerized plants dilute Phoenix Firebird Pro at the rate of 40 to 80 fluid ounces (0.2 to 0.4 lb a.i.) per 100 gallons and apply as a drench at the rate of 4 to 8 fluid ounces of finished spray per 6 inch (diameter) container Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter Ideally the media should be treated to the point of saturation which generally requires 1/5 the volume of the container Use the higher application rate for a longer period of control

Black Vine Weevil and White Grub Larval Control – Preventative Treatment – Media Incorporation For preventative control of black vine weevil and white grub larvae in containerized plants incorporate the appropriate volume of Phoenix Firebird Pro (see table below) per cubic yard of potting media by diluting it in water (typically 1 quart to 1 gallon per cubic yard of media) and sprinkling or spraying it onto the media Use the higher application rates for longer periods of control

Potting Media Bulk Density (lbs per cubic yard)	Fluid ounces of Phoenix Firebird Pro in one cubic yard			
	10 ppm	15 ppm	20 ppm	25 ppm
200	0.4	0.6	0.8	1.0
300	0.6	0.9	1.2	1.5
400	0.8	1.2	1.6	2.0
500	1.0	1.5	2.0	2.5
600	1.2	1.8	2.4	3.0
700	1.4	2.1	2.8	3.5
800	1.6	2.4	3.2	4.0
900	1.8	2.7	3.6	4.5
1000	2.0	3.0	4.0	5.0

The application rates listed above are based on the dry bulk density of the potting media

Use proportional volumes of Phoenix Firebird Pro for potting media with dry bulk densities that are not listed above

Black Vine Weevil Larval Control – Curative Treatment – Topical Drench To control black vine weevil larvae infesting containerized plants dilute Phoenix Firebird Pro at the rate of 10 to 40 fl ozs (0.05 to 0.2 lb a.i.) per 100 gallons and apply as a drench at the rate of 8 to 16 fluid ounces of finished spray per 6 inch (diameter) container Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter Ideally the media should be treated to the point of saturation which generally requires 1/5 the volume of the container

Bare root Treatment for Preventative Root Weevil Larvae – Control To protect treated roots of field grown nursery stock from feeding by root weevil larvae dilute one gallon of Phoenix Firebird Pro in 100 gallons of water and treat the bare roots of plants that are being re-planted into the field either by dipping the roots into the insecticide solution for ten seconds or by spraying the insecticide solution onto the roots

Diaprepes Weevil Larval Control – Curative Treatment – Topical Drench To control *Diaprepes* weevil larvae infesting containerized plants dilute Phoenix Firebird Pro at the rate 10 to 40 fl ozs (0.05 to 0.2 lb a.i.) per 100 gallons and apply as a drench at the rate of 8 to 16 fluid ounces of finished spray per 6 inch (diameter) container Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter Ideally the media should be treated to the point of saturation which generally requires 1/5 the volume of the container

Fungus Gnat Larval Control – Preventative Treatment – Topical Drench For preventative control of fungus gnat larvae in containerized plants dilute Phoenix Firebird Pro at the rate of 20 to 40 fl ozs (0.1 to 0.2 lb a.i.) per 100 gallons and apply as a drench at the rate of 4 to 8 fl ozs of finished spray per 6 inch (diameter) container Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter Ideally the media should be treated to the point of saturation which generally requires 1/5 the volume of the container Use the higher application rate for a longer period of control

Fungus Gnat Larval Control – Curative Treatment – Topical Drench To control fungus gnat larvae infesting containerized plants dilute Phoenix Firebird Pro at the rate of 10 to 40 fl ozs (0.05 to 0.2 lb a.i.) per 100 gallons and apply as a drench at the rate of 8 to 16 fluid ounces of finished spray per 6 inch (diameter) container Use a proportional volume of finished spray for containers less than or greater than 6 inches in diameter Ideally the media should be treated to the point of saturation which generally requires 1/5 the volume of the container

IMPREGNATION AND APPLICATION OF PHOENIX FIREBIRD PRO ON DRY BULK TURF FERTILIZERS

Phoenix Firebird Pro may be impregnated on dry bulk fertilizers. When applied as directed, Phoenix Firebird Pro/dry bulk fertilizer mixtures provide insect control equal to that provided by the same rates of Phoenix Firebird Pro applied in water.

Impregnation: Apply using a minimum 100 pounds of dry bulk fertilizer per acre with the specified amount of Phoenix Firebird Pro per acre. Use a closed rotary drum mixer or a similar type of closed blender equipped with suitable spray equipment. The spray nozzle(s) should be positioned to provide a uniform fine spray pattern over the tumbling fertilizer for thorough coverage. The physical properties of fertilizers vary particularly in liquid absorptive capacity. When absorptivity is sufficient, simple spray impregnation of the fertilizer with Phoenix Firebird Pro provides a satisfactory dry mixture. If the absorptive capacity is inadequate, use of a highly absorptive powder is required to provide a dry, flowable mixture. Microcel E (Johns Manville Products Corporation) is a recommended absorbent powder. Generally less than 2% by weight of Microcel E is required. **DO NOT** impregnate Phoenix Firebird Pro onto straight coated ammonium nitrate or straight limestone because these materials will not absorb the insecticide. Dry fertilizer blends containing mixtures of ammonium nitrate or limestone may be impregnated with Phoenix Firebird Pro.

The amount of Phoenix Firebird Pro actually required in the preparation of individual fertilizer mixtures should be determined carefully for each production operation. This is necessary to ensure that the amount of pesticide actually contained in the mixture applied to the soil represents the correct rate of use. Bulk fertilizer impregnated with Phoenix Firebird Pro should be applied immediately, not stored.

All individual Federal and State regulations relating to bulk dry fertilizer blending, registration, labeling, and application of the mixtures are the responsibility of the individual and/or company selling the fertilizer and Phoenix Firebird Pro mixture.

Fertilizer for this use should be Turf fertilizers recommended for specific regions.

PEST CONTROL ON OUTSIDE SURFACES AND AROUND BUILDINGS

For control of Ants, Bees, Biting Flies, Boxelder Bugs, Centipedes, Crickets, Earwigs, Elm leaf beetle, Fleas, Flies, Millipedes, Mosquitoes, Roaches (including Cockroaches), Silverfish, Sowbugs (Pillbugs), Spiders (including Black Widow Spiders), Ticks (including Brown Dog Ticks), and Wasps.

Follow additional application restrictions for residential outdoor surface and space sprays under Directions for Use.

Applications to vertical exterior surfaces (e.g., foundations) are permitted to a maximum height of 3 feet from ground level. Sections of vertical exterior surfaces that abut non-porous horizontal surfaces can only be treated if either 1) these sections are protected from rainfall and spray from sprinklers or 2) they do not drain into a sewer, storm drain, or curbside gutter (e.g., not to sections that abut driveways or sidewalks that drain into streets).

Apply Phoenix Firebird Pro using a 0.03 to 0.06% emulsion as a residual spray to outside surfaces of buildings including, but not limited to, exterior siding, foundations, porches, window frames, eaves, patios, garages, refuse dumps, lawns such as grass areas adjacent or around private homes, duplexes, townhouses, condominiums, house trailers, apartment complexes, carports, garages, fence lines, storage sheds, barns, and other residential and non-commercial structures, soil, trunks of woody ornamentals, and other areas where pests congregate or have been seen. Use a spray volume of 2 to 10 gallons of emulsion per 1,000 square feet.

Mixing Directions: For 0.03% emulsion, mix 0.5 fluid oz. of Phoenix Firebird Pro per gallon of water. For 0.06% emulsion, mix 1 fluid oz. Phoenix Firebird Pro per gallon of water (1 fluid oz. = 2 tablespoons). Do not use household utensils to measure Phoenix Firebird Pro. Use the higher rate for heavy pest infestation, quicker knockdown, or longer residual control. Repeat treatment may be necessary to achieve and/or maintain control during periods of high pest pressure. Repeat application is necessary only if there are signs of renewed insect activity. Repeat applications are permitted provided that the maximum label rate is not exceeded.

Perimeter Treatment: Treat a band of soil and vegetation 6 to 10 feet wide around and adjacent to the structure. Also, treat the foundation of the structure to a height of 2 to 3 feet. Apply 0.5 to 1.0 fluid oz. of Phoenix Firebird Pro per 1,000 square feet in sufficient water to provide adequate coverage (refer to Dilution Chart). For sections of foundation that abut non-porous horizontal surfaces, the treated areas must be protected from rainfall and spray from sprinklers or they do not drain into a sewer, storm drain, or curbside gutter (e.g., not to sections that abut driveways or sidewalks that drain into streets).

For Ant and Fire Ant Mounds use Phoenix Firebird Pro 0.06% emulsion as Drench Method: Apply 1.2 gallons of emulsion to each mound area by sprinkling the mound until it is wet and treat a 4-foot diameter circle around the mound. Use the higher volume for mounds larger than 12". For best results, apply in cool weather, such as in early morning or late evening hours, but not in the heat of the day.

GENERAL PRECAUTIONS

For turf treatments, apply with ground application equipment only (and apply with nozzles not more than two feet above the grass).

Do not apply when wind conditions favor downwind drift to nearby water bodies.

Do not apply when wind velocity exceeds 10 miles per hour.

Avoid application when wind gusts approach 10 mph.

Do not apply when a temperature inversion exists.

Apply using nozzles that provide the largest droplet size compatible with adequate coverage.

Do not apply for surface feeding pests if rain is expected within 12 hours (or whatever time is necessary for the spray to dry).

Do not apply by ground equipment within 25 feet of lakes, reservoirs, rivers, permanent streams, marshes, or natural ponds, estuaries, and commercial fish farm ponds.

Do not apply when grass areas are water logged or the soil is saturated with water (i.e., will not accept irrigation).

DEALERS MUST SELL IN ORIGINAL PACKAGES ONLY

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Firebird is a registered trademark of United Phosphorus, Inc.

Rev. 5/16/2012

70506 256(052912 4258)

