

70506-284

12/20/2013

1/25



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

Ms Sherry B. Hutcheson
United Phosphorus, Inc
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

DEC 20 2013

Subject: Notification for Label Change per PRN 98-10: General label changes
Application Date: 11/12/13
Product name: Siskin
EPA Reg. No. 70506-284
Decision Number 485496

Dear Ms Hutcheson:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 dated 11/12/13 for the above product. The Registration Division has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have questions concerning this letter, please call Banza Djapao at 703-305-7269 or via email at djapao.banza@epa.gov, or you may call me at 703-308-3194.

Sincerely,

A handwritten signature in cursive script that reads "Shaja B. Joyner".

Shaja B. Joyner
Product Manager 20
Fungicide Branch
Registration Division (7504P)

2/25

	EPA United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other: Notification	OPP Identifier Number
---	---	---	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 70506-284	2. EPA Product Manager William Cutchin	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Siskin	PM#	
5. Name and Address of Applicant (Include ZIP Code) United Phosphorus, Inc. 630 Freedom Business Center, Suite 402 King of Prussia, PA 19406		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____
<input type="checkbox"/> Check if this is a new address		

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application
<input type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below

Explanation: Use additional page(s) if necessary. (For Section I and Section II.)


Notification of minor changes.

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
*Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 1 gallon		5. Location of Label Directions <input checked="" type="checkbox"/> On Label <input type="checkbox"/> On labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input checked="" type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application)		
Name Sherry Hutcheson	Title Regulatory Affairs Manager	Telephone No. (Include Area Code) 229-247-9041
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Regulatory Affairs Manager	
4. Typed Name Sherry Hutcheson	5. Date 11/12/2013	

3/25



United Phosphorus, Inc.
Sherry B. Hutcheson
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
Phone: (229) 247-9041
Fax: (229) 241-9699

November 12, 2013

William Cutchin (Acting PM 21)
Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
U.S. Environmental Protection Agency
Ariel Rios Building
1200 Pennsylvania Ave., N.W.
Washington, DC 20460

RE: Notification of label changes for Siskin (EPA Reg. No. 70506-284)

Dear Mr. Cutchin,

United Phosphorus, Inc. (UPI) is notifying the Agency of our intent to make a couple of minor revisions to our product label, Siskin (EPA Reg. No. 70506-284) containing the active ingredient, Myclobutanil. We believe this qualifies as a notification under PR Notice 98-10. We have made the following changes to the label:

1. To the table for non-residential turf grass and residential turf grass, the section marked RESTRICTIONS applies to the entire table, not just the first two diseases.
- ~~X~~ 2. Under SPECIFIC USER DIRECTIONS FOR CHRYSANTHEMUM, New York has requested we revise the NOTE for use in New York.
3. We have corrected the word PLUMES to read PLUMS.

In addition to this letter the following are enclosed:

- EPA application form, 8570-1,
- Two copies of the label clearly marked to show the proposed changes,
- One clean copy of the proposed label.

If you have any questions, please feel free to contact me at 229-247-9041 or sherry.hutcheson@uniphos.com

Thank you for your kind attention to this matter.

Best regards,

Sherry B. Hutcheson
Regulatory Affairs Manager

4/25

NOTIFICATION

DEC 20 2013

Siskin Draft Label
Non - Notification
Clean Copy
November 11, 2013

Group	3	Fungicide
-------	---	-----------

Siskin (Alternate Brand Name: Phoenix Siskin)

A systemic, protectant, and curative fungicide for disease control in turf grass, landscape ornamentals, greenhouse and nursery ornamentals, apples, stonefruit, and grapes.

Active Ingredient:
 Myclobutanil [alpha-butyl-alpha-(chlorophenyl)-1H-1,2,4, triazole-1-propanenitrile]..... 19.7%
 Other Ingredients: 80.3%
 Total: 100.0%

Contains Petroleum Distillates
 Contains 1.67 lbs of myclobutanil per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Immediately call a poison control center or doctor. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Note to Physician: Contains petroleum distillate - vomiting may cause aspiration pneumonia.	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For medical emergency, call the Rocky Mountain Poison Control Center at 1-866-673-6671.	
FOR CHEMICAL EMERGENCY: spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.	

Net Content ___ Gallons

EPA Reg. No: 70506-284
 EPA Est. No. _____

United Phosphorus, Inc.
 630 Freedom Business Center, Suite 402
 King of Prussia, PA 19406
 1-800-438-6071

PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals
CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Personal Protective Equipment (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants,
- Chemical-resistant gloves made from barrier laminate,
- Shoes plus socks.

Non-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meet the requirements listed in the Worker Protection Standards (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY REQUIREMENTS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

PHYSICAL AND CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift or runoff from areas treated.

DIRECTIONS FOR USE

Shake Well Before Using.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all Directions for Use carefully before applying.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls,
- Barrier laminate gloves,
- Shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated area until sprays have dried.

7/25

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool and dry area. Store at temperatures above 32 °F.

PESTICIDE DISPOSAL: Wastes resulting in the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

For Non-refillable Containers (1, 2.5, 30 and 55 gallon): Do not reuse or refill this container. Offer for recycling, if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

For Non-refillable Containers (< 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

For Non-refillable Containers (> 5 gallons): Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows (all sizes): Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use for disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For Refillable Containers (250 gallon and bulk): Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing process two more times.

Steps to be taken in Case Material is Released or Spilled: Contain spills immediately with inert materials (e.g., sand, earth). Transfer liquids and solid diking material to separate suitable containers for recovery or disposal. Keep spills and cleaning runoff out of municipal sewers and open bodies of water. Spills on porous surfaces can contaminate groundwater.

PRODUCT INFORMATION

Siskin fungicide is a systemic, protectant, and curative fungicide for the control of diseases listed on this label in established turf grass, (including, but not limited to residential and commercial lawns, ornamental turf, grounds, or lawns around business and office complexes, and golf course fairways, roughs, tee boxes and greens), landscape ornamentals, greenhouse and nursery ornamentals, apples, stone fruit, and grapes. Optimum disease control is achieved when Siskin is applied in a regularly scheduled preventative spray program.

USE PRECAUTIONS

Fungicide Resistance Management:

Siskin belongs to the sterol demethylation inhibitor (DMI) class of fungicides and is classified as a Group 3 Fungicide by EPA. Since certain fungi can develop resistance to this class of products, the use of Siskin fungicide should be part of a resistance management strategy that includes alternation and/or tank mixing with fungicides of different modes of action. Consult your local or state agricultural authorities for resistance management strategies that are appropriate for your disease management program.

MIXING DIRECTIONS

Be sure sprayer is clean and not contaminated with other materials prior to use. Fill the spray tank $\frac{1}{4}$ to $\frac{1}{2}$ of the total amount of water required for the load. Start agitation and maintain agitation throughout mixing and application. Add the required amount of Siskin directly into the spray tank. Complete filling the tank. Always add Siskin to the spray tank before adding other materials.

Compatibility: Siskin is compatible with most commonly used fungicides, insecticides, growth regulators, micronutrients, and spray adjuvants. When preparing tank mixtures or spray compatibility charts State Cooperative Extension Service Specialist should be consulted prior to use.

When tank mixing Siskin with other pesticides, observe the more restrictive label limitations and precautions. No label dosage rates should be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing.

APPLICATION GUIDELINES

Carefully read, understand, and follow label use rates and restrictions. For proper application, determine the size of the area to be treated, the label use rate, and the gallonage to be applied to the area. Prepare only the amount of spray solution required to treat the measured area. Careful calibration of spray equipment is recommended prior to use.

GROUND APPLICATION: Thorough coverage sprays will result in optimum disease control. Application equipment should be properly calibrated and provide uniform spray coverage.

HANDGUN OR PRESSURIZED SPRAYERS: For best results when applying this product on a protectant schedule, ensure thorough coverage of all plant parts.

CHEMIGATION (SPRINKLER IRRIGATION): Siskin must be applied on a regular protectant fungicide schedule, not an irrigation schedule. If irrigation cycles are less frequent than the recommended application intervals for Siskin, ground or handgun applications must supplement chemigation applications to achieve adequate disease control. Apply this product only through solid set or hand-move sprinkler irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of fungicidal effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution

9/25

of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. Before applying Siskin through sprinkler irrigation equipment, the chemigation system must meet the following specifications.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Before applying Siskin through sprinkler irrigation equipment, the chemigation system must meet the following specifications:

- Public water system means a system for the provision to the public of piped water for human consumption if such system that has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
- Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone (RPZ), back flow preventer or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the flow outlet end of the pipe fill and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
- Systems not connected to a public water supply must contain a functional check valve, vacuum relief valve, and low-pressure drain appropriately located in the irrigation pipeline to prevent water source contamination from back flow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
- The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.

- Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Solid-Set and Hand Move Irrigation Equipment:

- Determine area covered by sprinkler.
- Fill injector solution tank with water and adjust flow rate to use the contents over a 10 to 30 minute interval.
- Determine the amount of Siskin required for the area to be treated.
- Add the required amount of Siskin into the same quantity of water used to calibrate the injection equipment.
- Maintain constant solution tank agitation during the injection period.
- Operate system at normal pressures recommended by the manufacturer of the injection equipment and used for the time interval established during calibration.
- Inject Siskin at the end of an irrigation cycle or as a separate application to maximize foliar absorption and retention.
- Stop injection equipment after treatment is completed. Continue to operate the system until the Siskin solution has cleared the last sprinkler head.

USE DIRECTIONS FOR TURF GRASS

Product Information

Use Siskin in conjunction with turf management practices that promote good plant health and optimum disease control. The key to selecting a fungicide is the proper diagnosis of the organism causing the disease. Kits, extension experts, or other identification methods should be used when developing disease control strategies.

In non-residential turf grass (including, but not limited to commercial lawns, ornamental turf, grounds or lawns around business and office complexes, and golf course fairways, roughs, tee boxes, and greens), optimum disease control is achieved when Siskin is applied in a preventative disease control program at a rate of 1.0 to 2.4 fluid ounces per 1000 square feet. In residential turf grass, optimum disease control is achieved when Siskin is applied in a preventative disease control program at a rate of 1.2 fluid ounces per 1000 square feet.

See the tables below for specific application rates for various diseases. Apply Siskin in sufficient water to ensure thorough coverage. For foliar diseases, use approximately one gallon of water per 1000 square feet. Use 2-3 gallons of spray solution per 1000 square feet to control diseases causing root and crown rots. Under conditions favorable for high disease development, reduce the spray interval between applications of Siskin. Under light to moderate disease pressure, apply Siskin at the lower use rate and/or longer treatment interval.

When disease pressure is high or when used as a curative treatment, use a higher rate of Siskin and a shorter treatment interval unless otherwise specified.

11/25

NON-RESIDENTIAL TURF GRASS ¹				
DISEASE	SISKIN™ (fl. oz./1000 sq. ft.)	APPLICATION INTERVAL (Days)	USE INSTRUCTIONS	RESTRICTIONS
Anthracnose Red thread Septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 13.8 fl. oz. of Siskin per 1000 sq. ft. per year.
Brown patch	1.2	14	Begin applications when conditions are favorable for disease development, but before disease symptoms are apparent. If disease is present, mix Siskin with an EPA registered contact fungicide, such as Wingman. Under conditions of high temperature and humidity, use the shorter spray interval.	
Copper spot Zonate leaf spot	1.2	14	Apply when conditions are favorable for disease development.	For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl. oz. of Siskin per 1000 sq. ft. per year (1.95 lbs. myclobutanil per acre).
Crown rot Leaf sot Melting-out	1.2	14	Apply when conditions are favorable for disease development.	
Dollar Spot	0.5	7	Apply when conditions are favorable for disease development.	
	0.5	14	Tank mix with a low label rate of Phoenix Pegasus chlorothalonil products.	
	1	21-28	Tank mix with the label rate of Phoenix Pegasus chlorothalonil products.	
	1 - 2.4	14 - 28	If using this rate without tank mixing, make no more than 3 consecutive applications for control of dollar spot before rotating to a registered fungicide with a different mode of action.	
Fusarium blight	1.2 - 2.4	14 - 21	Apply when conditions are favorable for disease development.	
Fusarium patch (pink snow mold)	1.2 - 2.4	Fall - Winter	Apply prior to snow cover.	

12/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

NON-RESIDENTIAL TURF GRASS ¹				
DISEASE	SISKIN™ (fl. oz./1000 sq. ft.)	APPLICATION INTERVAL (Days)	USE INSTRUCTIONS	RESTRICTIONS
Gray leaf spot	1.2 – 2.4	14	Apply when conditions are favorable for disease development. If using the lower rate, tank mix with a registered contact fungicide at its registered rate.	
Leaf smuts	1.2	14	Apply in the fall after turf grass enters dormancy and/or in the spring prior to the initiation of growth.	
Necrotic ring spots	1.2 – 2.4	Spring: 28	Make applications on a preventative basis in early to mid-spring.	
		Fall: 28	Make 2 applications beginning in August before the turf goes dormant. Apply 2.4 fl. oz./1000 sq. ft. followed by a second application one month later.	
Powdery mildew Rusts	1.2	14 – 28	Apply when conditions are favorable for disease development.	
Spring dead spot	2.4	Fall: 28	Make one to two applications in the fall before turf dormancy. Make 2 nd application one month later.	
Summer patch	1.2 – 2.4	14 – 28	Begin applications in spring when conditions are favorable for disease development. Make 2 to 4 applications depending on recommendations from local turf grass extension experts. Use at least 2 to 3 gallons of water per 1000 sq. ft. to increase spray penetration to crown and roots.	
Take-All patch	2.4	Spring/Fall: 28	Apply Siskin to reduce the severity of take-all patch. Make 1 to 2 fall applications in September and October or when night temperatures drop to 55°F, and 1 to 2 spring applications in April and May depending on local	

13/25

			recommendations.	
--	--	--	------------------	--

NON-RESIDENTIAL TURF GRASS ¹				
DISEASE	SISKIN™ (fl. oz./1000 sq. ft.)	APPLICATION INTERVAL (Days)	USE INSTRUCTIONS	RESTRICTIONS
Zoysi large patch	2.4	Fall: 28	Make applications in fall before turf dormancy.	

¹Including, but not limited to commercial lawns, ornamental turf, grounds or lawns around business and office complexes, and golf course fairways, rough, tee boxes, and greens.

RESIDENTIAL TURF GRASS				
DISEASE	SISKIN™ (fl. oz./1000 sq. ft.)	APPLICATION INTERVAL (Days)	USE INSTRUCTIONS	RESTRICTIONS
Anthracnose Red thread Septoria leaf spot	1.2	14 - 21	Apply when conditions are favorable for disease development.	Do not apply more than 13.8 fl. oz. Siskin per 1000 sq. ft. per year. For Nassau and Suffolk Counties in New York State, do not apply more than 3.43 fl. oz. of Siskin per 1000 sq. ft. per year (1.95 lbs. myclobutanil per acre).
Brown patch	1.2	14	Begin applications when conditions are favorable for disease development, and before disease symptoms are apparent. If disease is present, mix Siskin with an EPA registered contact fungicide, such as Wingman. Under conditions of high temperature and humidity, use the shorter spray interval.	
Copper spot Zonate leaf spot	1.2	14	Apply when conditions are favorable for disease development.	
Crown rot Leaf spot Melting-out	1.2	14	Apply when conditions are favorable for disease development.	
Dollar spot	1.2	14	Apply when conditions are favorable for disease development. Make no more than 3 consecutive applications for control of dollar spot before rotating to a registered fungicide with a different mode of action.	
Fusarium blight	1.2	14	Apply when conditions are favorable for disease	

14/25

			development.	
--	--	--	--------------	--

RESIDENTIAL TURF GRASS				
DISEASE	SISKIN™ (fl. oz./1000 sq. ft.)	APPLICATION INTERVAL (Days)	USE INSTRUCTIONS	RESTRICTIONS
Fusarium patch (pink snow mold)	1.2	Fall – Winter	Apply prior to snow cover.	
Gray leaf spot	1.2	14	Apply when conditions are favorable for disease development.	
Leaf smuts	1.2	14	Apply in the fall after turf grass enters dormancy and/or in the spring prior to the initiation of growth.	
Necrotic ring spot	1.2	Spring: 28	Make applications on a preventative basis in early to mid-spring.	
		Fall: 28	Make 2 applications beginning in August before the turf goes dormant. Apply 2.4 fl. oz./1000 sq. ft. followed by a 2 nd application one month later.	
Powdery mildew Rusts	1.2	14 – 28	Apply when conditions are favorable for disease development.	
Summer patch	1.2	14 – 28	Begin applications in the spring when conditions are favorable for disease development. Make 2 to 4 applications depending on recommendations from local turf grass extension experts. Use at least 2-3 gallons of water per 1000 sq. ft. to increase spray penetration to crown and roots.	

USE DIRECTIONS FOR LANDSCAPE, GREENHOUSE AND NURSERY ORNAMENTALS

Siskin is a locally systemic fungicide having protectant and curative properties that will translocate to new growth. For best control of labeled diseases, achieve thorough coverage of all plant parts on a protective application schedule. For dilute application sprays (> 100 gallons of spray volume per acre) applied to ornamental plants in greenhouses, field-grown plantings or in commercial and residential landscapes, apply Siskin at the rate of 6 to 12 fl. oz. per 100 gallons of spray volume on a 10 to 14 day application schedule, unless otherwise directed. Use the higher rate under conditions of high disease pressure and/or optimum conditions for infection.

For concentrate sprays (< 100 gallons of spray volume per acre) apply 8.0 fl. oz. per acre on a 10- to 14-day application schedule.

The addition of a non-phytotoxic spray adjuvant will improve spray coverage and fungicidal performance. Treated plants should be maintained in a vigorous growing condition. Plants under nutritional or water stress will not respond as well to treatment as well-maintained plants. Overdosage of Siskin can result in observable foliar greening, thickened leaves, and/or shortened internodes. If this condition is observed, reduce the fungicide use rate but do not extend the recommended application schedule.

Crop Tolerance

Plant tolerances are acceptable in the specific plants listed on this label. It is not possible to evaluate all ornamental plant species or varieties for tolerance to Siskin. The user should test for possible phytotoxic responses by treating a limited number of plants, at recommended use rates, prior to initiating "large-scale" use. The effects of spraying Siskin in combination with plant growth regulators are not fully understood at this time. If the use of a plant growth regulator is planned in an area being treated, the user should test for possible enhanced growth regulatory effects by treating a small number of plants, at the recommended use rates of all products, prior to initiating large-scale use. Since the effectiveness of such products depends not just on plant species or cultivar but also weather and seasonable differences (e.g., daylight hours), it is recommended that tests be repeated on previously tested varieties as environmental factors change and that observations for growth regulatory responses be made at regular intervals.

SPECIFIC USE DIRECTIONS FOR CHRYSANTHEMUM

Foliar Sprays: Best control is achieved by thorough coverage sprays, applied to point of runoff on a protectant application schedule. Use Siskin at a rate of 8 fl. oz. per 100 gallons of spray mixture. (Do not apply more than 19 fl. oz. of Siskin (0.25 lbs. myclobutanil) per acre per application). Application should be made on a 10 to 14 day schedule (not to exceed 21 days).

Prestick Dip Treatment: Chrysanthemum cuttings may be treated by a dip procedure prior to planting as follows: Prepare a dip suspension at a concentration equivalent to 8 fl. oz. of Siskin per 100 gallons of water. Cuttings must be fully submerged in the dip suspension until wet throughout (cuttings should not remain submerged longer than 2 minutes). If cuttings are dipped, this procedure will represent the first spray under the quarantine program. Used dip suspension should be disposed of if it becomes contaminated with soil, plant debris or other foreign matter. Dispose of used dip suspension by spraying it onto registered crops (but not onto previously dipped cuttings) after filtering, or in a manner consistent with local, state, and federal guidelines.

NOTE: All infected plant material must be destroyed if your state is under quarantine directive.

NOTE: This product is not for use on ornamentals in Nassau and Suffolk Counties, New York.

16/25

RESTRICTIONS ON USE ON ORNAMENTALS

- Do not apply more than 20 fl. oz. of Siskin (0.25 lbs. myclobutanil) per acre per application. On a total volume per acre basis, do not apply more than 333 gallons of spray per acre at the 6 fl. oz. per 100 gallons rate or 167 gallons per acre at the 12 fl. oz. per 100 gallons rate per application.
- Do not apply more than 153 fl. oz. of Siskin (2 lbs. myclobutanil) per acre per year.
- Do not use treated plant materials for food or feed.

USE DIRECTIONS FOR ORNAMENTALS

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Abelia	Cercospora leaf spot Powdery mildew		
Acalypha (Copper-leaf)	Cercospora leaf spot Powdery mildew		
Achillea (Yarrow)	Powdery mildew Rust		
African violet	Powdery mildew		
Ageratum	Rust Powdery mildew		
Alder	Powdery mildew Rust		
Almond, flowering	Blossom blight (<i>monilinia</i> spp.)	Apply prebloom, 50% bloom and at petal fall.	
Amelanchier (Juneberry, Shadbush)	Fabraea leaf spot Powdery mildew Rust		
Amorpha (False indigo)	Cercospora leaf spot Powdery mildew Rust		
Anemone	Rust		
Angelica	Cercospora leaf spot Rust		
Ash	Rust		
Aster	Rust Powdery mildew		
Australian pine	Diplodia tip blight		
Azalea	Petal blight (<i>Ovulinia</i> spp.) Powdery mildew	Begin applications when flowers start to exhibit color.	
Barberry	Powdery mildew Rust		May cause temporary damage to "crimson pigmy" and other "atropurposis" varieties.
Begonia	Powdery mildew		
Bellflower	Cercospora leaf spot Powdery mildew Rust		
Birch	Rust		
Bittersweet	Powdery mildew		

17/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Buckeye	Powdery mildew		
Butonbush	Cercospora leaf blight Powdery mildew Rust		
Clendula	Cercospora leaf spot		
California poppy	Powdery mildew		
Canna lily	Rust		
Carnation	Powdery mildew Rust		
Catalpa	Cercospora leaf spot Powdery mildew		
Cherry, flowering	Leaf spot Powdery mildew		
Chestnut, horse	Powdery mildew		
China aster	Rust		
Chokeberry	Rust Twig and Fruit blight		
Christmas trees	Rust		
Chrysanthemum	Rust White rust Ascochyta blight		
Columbine	Rust		
Cornflower	Rust		
Cosmos	Powdery mildew		
Cottonwood	Powdery mildew		
Crabapple, flowering	Powdery mildew Rust Scab		
Crepe-myrtle	Powdery mildew		
Daffodil	Rust		
Dahlia	Powdery mildew		
Delphinium	Powdery mildew Rust		
Dogwood	Anthracoese Powdery mildew Septoria leaf spot		
Douglas fir	Needle rust	Apply 12 to 18 fl. oz. per acre starting early spring. Continue applications at 2-3 week intervals until the threat of infection has passed. Spray adjuvants must be added to spray solutions to obtain good spray coverage and disease control.	
Dianthus	Rust		
Elm	Powdery mildew		
Euonymus	Powdery mildew		
Fern	Rhizoctonia aerial blight		

18/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Fleabane	Cercospora leaf spot Powdery mildew Rust		
Four O'clock	Rust		
Fuchsia	Rust		
Gaillardia	Powdery mildew Rust		
Gardenia	Powdery mildew Rust		
Geranium	Powdery mildew Rust		
Gerbera daisy	Powdery mildew		
Gourd, ornamental	Powdery mildew		
Grape leaf ivy	Powdery mildew		
Hackberry	Cercospora leaf spot Powdery mildew		
Hawthorn	Fabraea leaf spot Powdery mildew Rust Scab		
Hibiscus	Powdery mildew		
Holly	Powdery mildew		
Hollyhock	Powdery mildew Rust		
Honeysuckle	Cercospora leaf spot Powdery mildew		
Hydrangea	Cercospora leaf spot		
Iris	Didymellina leaf spot Rust	Apply 12 fl. oz. per 100 gallons of spray solution.	
Juniper	Rust		
Leucothoe	Cercospora leaf spot		
Leyland Cyprus	Cercospora leaf spot		
Lilac	Powdery mildew		
Loblolly pine	Fusiform rust	Refer to Douglas fir.	
Locust	Powdery mildew		
Maple	Powdery mildew		Treated trees may not be used for syrup production. Do not apply to Abutilon (Flowering Maple).
Marigold	Cercospora leaf spot Rust		
Mock-orange	Powdery mildew Rust		
Moonflower	Rust		
Mountain laurel	Cercospora leaf spot Ovulinia petal blight Powdery mildew	Refer to Azalea.	
Nephtytis	Cephalosporium leaf spot		
Ninebark	Rust		
Oak	Powdery mildew		

19/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

Pansy	Powdery mildew Rust		
-------	------------------------	--	--

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Pear, flowering	Powdery mildew Rust Scab		
Petunia	Powdery mildew Rust		
Phlox	Cercospora leaf spot Powdery mildew Rust		
Photinia	Entomosporium leaf spot Powdery mildew Rust		
Poinsettia	Powdery mildew Poinsettia scab		
Poplar	Rust		
Potentilla	Rust		
Privet	Cercospora leaf spot Powdery mildew		
Pyracantha (Firethorn)	Fusicladium scab		
Quince, flowering	Blossom and Twig blight Cercospora leaf spot Fabraea leaf spot Rust		
Rhododendron	Cercospora leaf spot Ovlinia petal blight Powdery mildew	Refer to Azalea.	
Rose	Black spot Powdery mildew Rust	Apply on a 7 to 10 day protectant schedule. In areas where black spot is not a problem, spray intervals may be increased to a maximum of 14 days. Greenhouse rose varieties vary in their sensitivity to Siskin. User should evaluate for possible abnormal response by treating a limited number of plants at recommended rates, prior to initiating large-scale use.	
Russian olive	Cercospora leaf spot Rust		
Salvia	Powdery mildew Rust		
Sedum	Powdery mildew		
Slash pine	Fusiform rust	Refer to Douglas fir.	
Smoke-tree (Cotinus)	Cercospora leaf spot Rust		

20/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

Snapdragon	Powdery mildew Rust		
Spirea	Powdery mildew		

CROP	DISEASE	USE INSTRUCTIONS	RESTRICTIONS
Sunflower	Cercospora leaf spot Powdery mildew Rust		Seeds from treated plants may not be used for food or feed.
Sycamore	Powdery mildew		
Trumpet creeper	Cercospora leaf blight Powdery mildew		
Viburnum	Powdery mildew Rust		
Walnut	Powdery mildew		Nuts from treated trees may not be used for food purposes.
Willow	Powdery mildew		
Zinnia	Cercospora leaf spot Powdery mildew		

USE DIRECTIONS FOR HOME ORCHARDS, VINEYARDS, OR FRUIT TREES

Best control of labeled diseases is achieved when Siskin is applied on a 7 to 10 day protectant schedule. Siskin is a systemic fungicide and does not redistribute after application. Application equipment spray nozzles should be adjusted to apply a uniform spray throughout the entire tree canopy.

Dilute (thorough coverage) applications are recommended and are based on the amount of spray solution required to thoroughly wet plants to the point of run-off. Refer to use directions for specific tree fruits and vines to determine actual use rate per 100 gallons of spray for control of labeled diseases. The following specific use directions are based on a dilute spray volume of 300 gallons per acre.

USE DIRECTIONS FOR APPLES

DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Powdery mildew	4-6	Begin application at tight cluster and continue through the second cover spray. Additional sprays beyond second cover may be needed on susceptible varieties or under heavy disease pressure. Use high rate in rate range if powdery mildew was present in previous years.	Do not apply within 14 days of harvest. Do not apply more than 153 fl. oz. of Siskin (2 lbs. myclobutanil) per acre per season.
Rusts (<i>Gymnosporangium</i> spp.)	4-6	Begin applications at pink stage and continue through the second cover spray.	

2/1/25

DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Scab (<i>Venturis</i> spp.)	4-6	Begin application at green tip or when environmental conditions become favorable for primary scab development. Apply Siskin alone or tank mixed with a protectant fungicide on a 7-10 day schedule.	
Prebloom			
Bloom, Postbloom	4-6	Use Siskin in a tank mixture with the recommended rate of a protectant fungicide, registered for use on apples, for improved fruit scab and summer disease control.	
Post Infection	6	Siskin provides 96-hour post-infection control or curative activity. Apply as soon as possible after infection period. Follow with a standard preventative spray schedule.	

USE DIRECTIONS FOR STONE FRUIT

APRICOTS			
DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Brown rot Blossom blight (<i>Monilinia</i> spp.)	2-3	Begin application at early red bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 84 fl. oz. of Siskin (1.1 lbs. myclobutanil) per acre per season. Applications may be made up to the day of harvest.
Brown rot (<i>Monilinia</i> spp.)		Apply 12 fl. oz. (0.16 lbs. myclobutanil) per acre on a 7-14 day protectant schedule. Apply whenever environmental conditions favor disease development during the month prior to harvest.	
Powdery mildew (<i>Podosphaera</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 10-14 day intervals until terminal growth ceases.	
Shothole (<i>Stigmina</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 7-10 day intervals as long as needed.	

22/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

CHERRIES			
DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Brown rot Blossom blight (<i>Monilinia</i> spp.)	2-3	Begin application at early popcorn stage, before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 100 fl. oz. of Siskin (1.3 lbs. myclobutanil) per acre per season. Applications may be made up to the day of harvest.
Brown rot (<i>Monilinia</i> spp.)		Refer to Apricots.	
Powdery mildew (<i>Podosphaera</i> and <i>Sphaerotheca</i> spp.)		Refer to Apricots.	
Leaf spot (<i>Blumeriella</i> spp.)		Follow the brown rot blossom blight schedule. Reapply at 7-10 day intervals. Make additional applications after harvest.	

23/25

Siskin Draft Label
 Non - Notification
 Clean Copy
 November 11, 2013

NECTARINES			
DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Brown rot Blossom blight (<i>Monilinia</i> spp.)	2-3	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 100 fl. oz. of Siskin (1.3 lbs. myclobutanil) per acre per season.
Brown rot (<i>Monilinia</i> spp.)		Refer to Apricots.	
Powdery mildew (<i>Podosphaera</i> and <i>Shaerotheca</i> spp.)		Refer to Apricots.	Applications may be made up to the day of harvest.
Shothole (<i>Stigmia</i> spp.)		Follow brown rot blossom blight schedule. Reapply at 7-10 day intervals as long as needed.	

PEACHES			
DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Brown rot Blossom blight (<i>Monilinia</i> spp.)	2-3	Begin application at early pink bud stage before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall.	Do not apply more than 100 fl. oz. of Siskin (1.3 lbs. myclobutanil) per acre per season.
Brown rot (<i>Monilinia</i> spp.)		Refer to Apricots.	
Powdery mildew (<i>Podosphaera</i> spp.)		Refer to Apricots.	Applications may be made up to the day of harvest.
Rust (<i>Tranzschelia</i> spp.)		Apply 12 fl. oz. (0.16 lbs. myclobutanil) per acre. Begin application approximately 8 weeks after flowering if environmental conditions are favorable for disease development. For optimum disease control, do not	

24/25

		apply on a protectant schedule exceeding 21 days.	
--	--	---	--

PLUMS/PRUNES			
DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Brown rot Blossom blight (<i>Monilinia</i> spp.)	2-3	Begin application at green tip, before infection occurs. If conditions are favorable for disease development, reapply at full bloom and petal fall	Do not apply more than 84 fl. oz. of Siskin (1.1 lbs. myclobutanil) per acre per season. Applications may be made up to the day of harvest.
Rust (<i>Tranzschelia</i> spp.)		Refer to Peaches.	

USE DIRECTIONS FOR GRAPES

GRAPES			
DISEASE	SISKIN™ (fl. oz./100 gal)	USE INSTRUCTIONS	RESTRICTIONS
Anthracnose (<i>Elsinoe</i> spp.)	6-10	Begin application when new shoots are 1 to 3 inches in length and continue on an application schedule that does not exceed 14 days.	Preharvest Interval: Do not apply within 14 days of harvest. Do not apply more than 46 fl. oz. of Siskin (0.6 lbs. myclobutanil) per acre per season.
Black rot (<i>Guignardia</i> spp.)		Preventative Schedule: Begin application when new shoots are 1 to 3 inches in length. Reapply on a protectant schedule that does not exceed 14 days. Use a higher rate under heavy disease pressure. Post-infection Schedule: Apply within 72 hours after the beginning of an infection period.	
Powdery mildew (<i>Uncinula</i> spp.)		Begin application at prebloom (12 to 18 inch shoots) and do not extend applications beyond a 21 day interval. Use a higher rate or shorter spray interval on susceptible varieties or under heavy disease pressure.	

25/25

IMPORTANT INFORMATION

READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. To the extent consistent with applicable law all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Siskin is a registered trademark of United Phosphorus, Inc.

Rev. 11/11/2013