

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460



OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

October 21, 2014

Sherry Hutcheson
Regulatory Affairs Manager
United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

Subject: Label Amendment – addition of respirator for broccoli, cabbage, and pepper uses-
master and supplemental labels
Product Name: Elixir Fungicide
EPA Registration Number: 70506-298
Application Date: 5/6/2014, resubmissions dated 7/24/2014 & 8/15/2014
Decision Number: 491583


Dear Ms. Hutcheson:

The amended label and supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable.

A stamped copy of your labeling is enclosed for your records. The master labeling supersedes all previously accepted master labeling. The supplemental labeling supercedes the previously accepted version of this supplemental labeling. The next label printing of this product must use this labeling unless subsequent changes have been approved. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.


Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact me by phone at 703-305-5410, or via email at johnson.hope@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Hope Johnson". The signature is fluid and cursive, with the first name "Hope" and the last name "Johnson" clearly distinguishable.

Hope Johnson, Product Manager 21
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure



Elixir Fungicide
Label amendment – Clean copy
July 21 , 2014

GROUP	M3, M5	FUNGICIDES
-------	--------	------------

ELIXIR™ fungicide

ACTIVE INGREDIENTS	BY WEIGHT
Mancozeb: A coordination product of zinc ion and manganese ethylenebisdithiocarbamate	62.5%
in which the ingredients are:	
Manganese++	12.5%
Zinc++	1.6%
Ethylenebisdithiocarbamate ion (C ₄ H ₆ N ₂ S ₄)--	48.4%
Chlorothalonil (tetrachloroisophthalonitrile).....	12.5%
OTHER INGREDIENTS	<u>25.0%</u>
TOTAL	100.0%

Contains 0.625 pound of mancozeb and 0.125 pound of chlorothalonil per pound of product

Patent pending.
EPA Reg. No. 70506-298 EPA Est. No. XXX-XXX-XXX

KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.
Contact the Rocky Mountain Poison Center at 1-866-673-6671 for emergency medical treatment information. .

**FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure,
or accident, call CHEMTREC at 1-800-424-9300.**

Net Contents: 45 pounds


United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. Causes substantial but temporary eye injury. Do not get in eyes or on clothing. e Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes and clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

PHYSICAL AND CHEMICAL HAZARDS

Mancozeb decomposes in acid and alkaline conditions, with heat, and upon exposure to moisture and air. May be ignited by heat or open flame.Keep away from fire or sparks.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for Category A on an EPA chemical resistance category selection chart.

Mixers,loaders, applicators, and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Chemical resistant gloves made of any waterproof material (except pilots, groundboom applicators, and airblast applicators)
- Shoes and socks
- Protective eyewear

Aerial application of Elixir on broccoli, cabbage, and peppers requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R, or P filter, NIOSH approval prefix TC-84A.

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

ENGINEERING CONTROL STATEMENTS:

When handlers use enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR part 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

During aerial application, human flaggers must be in enclosed cabs.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/ PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through

drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical resistant gloves made of any waterproof material
- Shoes and socks
- Protective eyewear

Special Eye Irritation Provisions: Chlorothalonil in this product is a severe eye irritant. Although the restricted entry interval expires after 24 hours, for the next 6 days entry is permitted only when the following safety measures are provided:

- (1) At least one container designed specifically for flushing eyes must be available in operating condition at the WPS required decontamination site intended for workers entering the treated area.
- (2) Workers must be informed, in a manner they can understand:
 - that residues in the treated area may be highly irritating to their eyes
 - that they should take precautions, such as refraining from rubbing their eyes to keep the residues out of their eyes
 - that if they do get residues in their eyes, they should immediately flush their eyes using the eyeflush
 - container that is located at the decontamination site, or using other readily available clean water
 - how to operate the eyeflush container

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Application to lawn grasses, golf courses, industrial (office park) and municipal lawns are not within the scope of the Worker Protection Standard.

Do not enter treated areas until sprays have dried.

ELIXIR™ fungicide is a water dispersible granule (WDG) labeled for use as a spray for the control of many important plant diseases.

APPLICATION INSTRUCTIONS

AS A SPRAY (Ground or Aerial Equipment) - Apply ELIXIR fungicide at the rate shown; use sufficient water to provide thorough coverage. Use 20 to 100 gallons per acre for ground equipment, and no less than 2 gallons per acre for aircraft. Add ELIXIR fungicide slowly to water in the spray tank with agitation, or premix thoroughly in separate holding tank for concentrate or aircraft sprayers. Continuous agitation is

required to keep the product in suspension. A spreader-sticker spray adjuvant may be used with this product if needed; contact your local product distributor or United Phosphorus, Inc. representative for specific recommendations. If tank mixed, follow the more restrictive labeling of any tank mix partner. Do not tank mix with any product that contains a prohibition on tank mixing.

RESTRICTIONS

Foliar Applications

Where EBDC Products Used Allow the Same Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient (mancozeb or metiram) is used on a crop during the same growing season and the EBDC products used allow the same maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed any one of the specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

Where EBDC Products Used Allow Different Maximum Poundage of Active Ingredient Per Acre Per Season

If more than one product containing an EBDC active ingredient is used on a crop during the same growing season and the EBDC products used allow different maximum poundage of active ingredient per acre per season, then the total poundage of all such EBDC products used must not exceed the lowest specified individual EBDC product maximum seasonal poundage of active ingredient allowed per acre.

CHEMIGATION

Apply ELIXIR fungicide only through sprinkler systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set or hand move irrigation systems. Do not apply ELIXIR fungicide through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact State Extension Service Specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems:

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve

located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Sprinkler Irrigation Systems:

1. The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Good agitation is required in the injection tank.
9. In moving systems, apply specified dosage of ELIXIR fungicide as a continuous injection. In non-moving systems inject ELIXIR fungicide for 15 to 30 minutes at end of cycle. Use the least amount of water possible consistent with uniform coverage.
10. Mix the amount of ELIXIR fungicide needed for acreage to be treated into the quantity of water determined during prior calibration. For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.
11. Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all ELIXIR fungicide is flushed from system.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g. wind direction, wind speed, temperature, relative humidity) and method of application (e.g. ground, aerial, airblast, chemigation) can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Wind Speed

Do not apply at wind speeds greater than 15 mph.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of mancozeb. Where states have more stringent regulations, they must be observed.

Equipment

All aerial and ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates.

Additional requirements for aerial applications:

1. The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.
2. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy unless a greater height is required for aircraft safety.
3. When applications are made with a crosswind, the swath must be displaced downwind. The applicator must compensate for this displacement at the up and downwind edge of the application area by adjusting the path of the aircraft upwind.

Additional requirements for ground boom application:

1. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

USE DIRECTIONS

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
Almonds	Anthrachnose, Blossom Blight (<i>Monilinia spp.</i>), Shothole (<i>Stigmima spp.</i>), Rust, Scab	6.4-7.7 lbs/A	Apply by ground or air. Begin application at dormant to popcorn stage, full bloom or petal fall. Reapply every 7 to 10 days if bloom is staggered and weather is rainy. Apply in sufficient water and with proper calibration to obtain uniform coverage of tree canopy. Do not use less than 10 gallons of spray volume per acre if aerially applied.	Do not apply more than 23.0 lbs of product per acre per season. Do not make last application later than 150 days before harvest (150 day PHI) Do not graze livestock in treated area. Minimum retreatment interval is 7 days. Do not apply this product with a U-boom device.
Asparagus	Cercospora Leaf Spot, Rust, Purple Spot	2.0-2.4 lbs/A	Apply by ground. Start applications when rust first appears and repeat at 14 to 28 day intervals. Use 25-50 gallons of water per acre.	Apply only on asparagus ferns after spears have been harvested. Do not apply more than 9.6 lbs of product per acre per season. Do not apply within 190 days (120 days in CA and AZ) of harvest. (190 day PHI; 120 day PHI in CA and AZ)
Broccoli	Alternaria Leaf Spot, Downy Mildew	1.6 – 2.5 lbs/A	Apply by ground, air, or chemigation. Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when	Do not apply more than 15.4 lbs of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI). Do not apply this

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
			conditions favor disease.	product with a U-boom device. Aerial application of Elixir on broccoli, requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R. or P filter, NIOSH approval prefix TC 84-A.

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
Cabbage	Alternaria Leaf Spot, Downy Mildew	1.6 – 2.5 lbs/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10-day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 15.4 lbs of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI). Do not apply this product with a U-boom device. Aerial application of Elixir on cabbage, requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R. or P filter, NIOSH approval prefix TC 84-A.
Corn sweet corn for fresh use and sweet corn for seed production, including hybrid seed	Common Rust, Helminthosporium Leaf Blight, Gray Leaf Spot	1.5-1.8 lbs/A	Apply by ground, air, or chemigation. Use sufficient water for thorough coverage. Start applications when disease first appears and repeat at 7 day intervals.	East of the Mississippi River, Arkansas and Louisiana: Do not apply more than 27.0 lbs of product per acre per season. West of the Mississippi River except Arkansas and Louisiana: Do not apply more than 9.0 lbs of product per acre per season. Do not apply to sweet corn in home gardens. Do not allow livestock to graze in treated fields. Do not ensile treated corn or use as livestock forage. Minimum preharvest interval is 14 days (14 day PHI).
Cranberry	Fruit Rot, Lophodermium	3.0-7.2 lbs/A	Apply by ground, air, or chemigation. When applying	Do not apply more than 21.6 lbs of

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
	Leaf/Twig Blight		by chemigation, use 300 gallons of water per acre through solid set systems only. Start applications at mid-bloom and repeat at 10 day intervals.	product per acre per season. Minimum preharvest interval is 50 days (50 day PHI).
Cucurbit crop group Cucumber, Muskmelon, Pumpkin, Squash (summer), Squash (winter), Watermelon Additional Cucurbit crops: Chayote, Chinese waxgourd, Citron melon, Gherkin, Gourds edible (<i>Momordica spp.</i>)	Alternaria Leaf Spot, Anthracnose, Cercospora Leaf Spot, Downy Mildew, Gummy Stem Blight, Scab	2.0-3.6 lbs/A	Start applications when the plants are in the two-leaf stage and repeat at 7 to 10 day intervals. Use sufficient water and direct spray to provide thorough coverage of both upper and lower leaf surfaces. For aerial applications, the minimum spray volume is 2 gallons per acre. Some cantaloupe varieties (i.e. Harvest Queen, Gold Star, Super Star, Sweet and Early, and Saticoy) are sensitive to ELIXIR fungicide. Consult your State Cooperative Extension Service Specialist prior to use.	Do not apply more than 30.7 lbs of product per acre per year. Do not apply more than 8 applications per year. Minimum preharvest interval is 5 days (5 day PHI)
Ginseng	Alternaria Blight, Gray Mold	2.0-2.4 lbs/A	Start applications when disease first threatens and repeat every 7 to 10 days as needed. In Wisconsin, apply with ground equipment and a minimum of 80 gallons of water per acre.	Do not apply more than 28.8 lbs of product per acre per year. Do not apply more than 12 applications per year. Minimum preharvest interval is 30 days (30 day PHI).
Mango	Anthracnose, Phytophthora Fruit Rot, Black Spot (<i>Cercospora</i>)	2.0-3.0 lbs/A	Start applications at flowering and continue at 14 to 21 day intervals. Direct spray to crown and blossom area. Use 20 to 100 gallons water per acre.	Do not apply more than 44.8 lbs. of product per acre per year. Do not apply more than 14 applications per year. Minimum preharvest interval is 21 days (21

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
				day PHI).
Onion (Dry Bulb), Garlic, Shallots (Furrow Drench)	Botrytis Leaf Blight, Downy Mildew, Neck Rot, Purple Blotch	3.0-3.6 lbs/A	Apply by air, ground, or chemigation. Follow a protective spray schedule starting when diseases are first reported in the area and repeat at 7 day intervals throughout the season.	Do not apply more than 36.0 lbs of product per acre per crop. Do not apply to exposed bulbs. Minimum preharvest interval is 7 days (7 day PHI).
	Smut	3.0-3.6 lbs/A	Apply 3 to 3.6 lbs per acre as a furrow drench at time of planting onion seeds. Use 75 to 125 gallons of water per acre.	Do not apply more than 3.6 lbs of product per acre (29,000 linear feet of furrow) with an 18- inch row spacing.
Papaya	Anthracnose (<i>Colletotricum</i>), Phytophthora Fruit Rot, Black Spot (<i>Cercospora</i>), Alternaria	2.0-3.0 lbs/A	Begin at flowering; treat central column crown, blossom area and developing fruit. Repeat at 14 to 21 day intervals. Use a minimum 50 gallons of water per acre.	Do not apply more than 42.0 lbs of product per acre per crop. Minimum preharvest interval 0 days (0 day PHI)
Peanut	Ascochyta Web Blotch, Cercospora Leaf Spot, Rust, Late Leaf Spot, Pepper Spot	1.0-2.4 lbs/A	Apply by ground, air, or chemigation. Start application when disease first appears or is reported in area. Repeat sprays at 14 day intervals.	Do not apply more than 19.2 lbs of product per acre per crop. Do not feed treated vines, hay, or threshings to livestock. Do not allow livestock to graze in treated areas. Minimum preharvest interval is 14 days (14 day PHI).

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
Pepper	Anthracnose, Cercospora Leaf Spot (Frogeye Spot), Phytophthora Blight, Ripe Rot, Botrytis Leaf Mold	West of the Mississippi River 1.6–2.5 lbs/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 15.4 lbs of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI). Do not apply this product with a U-boom device. Aerial application of Elixir on pepper, requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R. or P filter, NIOSH approval prefix TC 84-A.
		East of the Mississippi River 1.6–3.8 lbs/A	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 23.0 lbs of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI). Do not apply this product with a U-boom device. Aerial application of Elixir on pepper, requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R. or P filter, NIOSH approval prefix TC 84-A.
Potato	Early Blight,	1.2-2.4	Apply by ground, air, or	Do not apply more

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
	<p>Late Blight, Black Dot</p> <p>Disease Suppression: Botrytis</p>	lbs/A	<p>chemigation. Begin applications when plants are 4 to 6 inches high by applying 1.2 lb of product per acre. As the vines increase in size, apply 1.5 to 2.4 lbs of product per acre at intervals of 5 to 10 days; or 1.2 lb of product per acre at 5 day intervals. Use the highest rate and shortest interval when plants are rapidly growing.</p>	<p>than 18.0 lbs of product per acre per crop. Vine kill should occur 14 days before harvest. It is recommended that this product be used in an Integrated Pest Management Program. Minimum preharvest interval is 14 days except CT, DE, FL, MA, ME, MI, NY, OH, PA, RI, VT and WI (14 day PHI). Minimum preharvest interval is 7 days in CT, DE, FL, MA, ME, MI, NH, NY, OH, PA, RI, VT, and WI (7 day PHI).</p>
Tomato	<p>Anthracnose, Early Blight, Gray Leaf Spot, Late Blight, Leaf Mold, Septoria Leaf Spot</p>	<p>(West of the Mississippi River) 0.75-1.2 lbs/A</p> <p>(East of the Mississippi River) 0.75-1.8 lbs/A</p>	<p>Start application when seedlings emerge or transplants are set. Apply at 7 day intervals throughout the season.</p>	<p>West: Do not apply more than 10.2 lbs of product per acre per crop. East: Do not apply more than 26.9 lbs of product per acre per crop.</p> <p>Minimum preharvest interval is 5 days (5 day PHI).</p> <p>FOR BACTERIAL SPECK AND SPOT: Use a full rate of a fixed copper fungicide (such as Cuprofix® Ultra Disperss®) in a tank mix combination with a ½ to full rate of ELIXIR fungicide. Follow the application interval specified on the copper fungicide label.</p>
	Bacterial Speck and Spot	<p>(West of the Mississippi River) 1.5-2.4 lbs/A</p> <p>(East of the Mississippi River) 1.5-3.6 lbs/A</p>	<p>Start application when seedlings emerge or transplants are set. Apply at 7 to 10 day intervals throughout the season.</p>	

**FLOWERS, FOLIAGE PLANTS, AND ORNAMENTALS
 NOT INTENDED FOR USE ON FRUIT TREES BY HOMEOWNERS.**

**TREATED PLANTS, FRUITS, NUTS OR SYRUP FROM MAPLE TREES MUST NOT BE USED FOR
 FOOD OR FEED PURPOSES.**

Apply in the field, nursery or greenhouse as a thorough coverage spray, using 1.0 to 2.4 lbs. ELIXIR fungicide per acre (1 ½ to 3 ⅔ tsp. per gal.). Do not use in residential greenhouses.

Plant sensitivities to ELIXIR fungicide have been found to be acceptable in specific genera and species listed on this label, however, phytotoxicity may occur. Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test each one for sensitivity to ELIXIR fungicide. Neither the manufacturer nor seller has determined whether or not ELIXIR fungicide can be safely used on ornamental or nursery plants not listed on this label. The user is responsible for determining if ELIXIR fungicide can be used safely prior to commercial use. In a small area, apply the specified rates to the plants in question, i.e. bedding plants, foliage, etc., and observe for 7 to 10 days for symptoms of phytotoxicity prior to commercial use. Use ELIXIR fungicide in commercial greenhouses and nurseries for control of fungal diseases of flowers, foliage and ornamentals. Do not make more than 20 applications per year.

Aerial application: For aerial applications made to field-planted ornamentals, apply 1.0 to 2.4 lbs. per acre; use a minimum rate of 5 gals of spray per acre during aerial applications.

Application of dilute sprays: Apply as a thorough coverage spray using 1.0 to 2.4 lbs. per acre or 1.0 to 2.4 lbs. per 100 gals of water. Begin application at first sign of disease and repeat at 7 to 10 day intervals or as needed; use shorter interval during periods of frequent rains or when severe disease conditions persist. ELIXIR fungicide may be used alone or in combination with other fungicides as maintenance spray. Use higher rate and shorter intervals during periods of excessive wetness and rapid growth.

ELIXIR fungicide is labeled for use on certain flower, foliage and ornamental plants listed in the table below for control of the following diseases and pathogens

PLANT	PATHOGEN CONTROLLED:
Aglaonema	Alternaria
Almond, ornamental	Botrytis, Cladosporium, Coryneum, Gloeosporium, Monilinia
Andromeda	Exobasidium, Rhytisma, Venturia
Ash	Cercospora, Cylindrosporium, Gloeosporium, Puccinia, Rhizoctonia, Sphaeropsis
Aster	Alternaria, Ascochyta, Botrytis, Colletotrichum, Fusarium, Phomopsis, Phyllosticta, Puccinia, Ramularia, Rhizoctonia, Septoria, Uromyces
Azalea	Alternaria, Botrytis, Cladosporium, Colletotrichum, Cylindrocladium, Ovulinia
Baby's Breath	Botrytis, Rhizoctonia
Begonia	Botrytis, Cercospora, Gloeosporium, Rhizoctonia
Bougainvillea	Colletotrichum
Buckeye	Cercospora, Glomerella, Guignardia, Monchaetia, Phyllosticta, Septoria, Taphrina
Camellia	Botrytis, Cercospora, Elsinoe, Exobasidium, Glomerella, Pestalotia, Phomopsis, Phyllosticta
Carnation	Alternaria, Botrytis, Cladosporium, Colletotrichum, Fusarium, Helminthosporium, Septoria, Stemphylium, Uromyces
Cherry, ornamental	Alternaria, Cercospora, Cladosporium, Cocomyces, Coryneum, Fusicladium, Monilinia, Phomopsis, Phyllosticta, Taphrina
Christmas cactus	Alternaria, Cercospora, Colletotrichum, Fusarium, Phomopsis
Chrysanthemum	Alternaria, Ascochyta, Bipolaris, Botrytis, Cercospora, Cylindrosporium, Helminthosporium, Phyllosticta, Septoria, Stemphylium

Crabapple	Gymnosporangium, Marssonina, Phyllosticta, Septoria, Venturia
Croton	Gloeosporium
Daisy	Botrytis, Cercospora, Whetzelinia
Dogwood	Ascochyta, Botrytis, Cercospora, Colletotrichum, Elsinoe, Phyllosticta, Septoria
Douglas Fir	Phaeocryptopus
Dracaena	Alternaria, Cercospora, Colletotrichum, Fusarium, Phyllosticta
Euonymus	Cercospora, Colletotrichum, Gloeosporium, Marssonina, Ramularia, Septoria, Whetzelinia
Fatsia	Alternaria, Cercospora, Colletotrichum, Phyllosticta
Ficus	Alternaria, Ascochyta, Cephalosporium, Cercospora, Cladosporium, Colletotrichum, Fusarium, Gloeosporium, Glomerella, Mycosphaerella, Phomopsis, Stemphylium
Firethorn	Fusarium, Fusicladium, Rhizoctonia
Geranium	Alternaria, Ascochyta, Bipolaris, Botrytis, Cercospora, Cyindrosporium, Helminthosporium, Puccinia, Ramularia, Rhizoctonia, Septoria, Uromyces, Venturia
Gladiolus*	Alternaria, Botrytis, Cladosporium, Curvularia, Rhizoctonia, Septoria, Stemphylium
Hawthorn	Cercospora, Cyindrosporium, Gloeosporium, Gymnosporangium, Monilinia, Mycosphaerella, Phyllosticta, Septoria, Venturia
Holly	Phyllosticta
Hollyhock	Alternaria, Ascochyta, Cercospora, Colletotrichum, Puccinia, Septoria
Hydrangea (foliage only)	Ascochyta, Botrytis, Cercospora, Colletotrichum, Phyllosticta, Rhizoctonia, Septoria
Impatiens	Cercospora, Phyllosticta, Rhizoctonia, Septoria
Iris	Ascochyta, Botrytis, Cladosporium, Fusarium, Kabatiella, Phyllosticta, Puccinia, Rhizoctonia
Jade plant	Gloeosporium, Phomopsis
Laurel, Cherry	Alternaria, Cercospora, Coccomyces, Monilinia, Phyllosticta, Septoria
Lilac	Botrytis, Cercospora, Cladosporium, Cyindrocladium, Gloeosporium
Lily	Botrytis, Cercospora, Cladosporium, Colletotrichum, Fusarium, Puccinia, Ramularia, Rhizoctonia
Magnolia	Alternaria, Cercospora, Cladosporium, Colletotrichum, Glomerella, Rhizoctonia
Mahonia	Cercospora, Cyindrocladium, Gloeosporium, Leptosphaeria, Phomopsis, Phyllosticta, Puccinia
Maple	Alternaria, Cercospora, Ciborinia, Fusarium, Marssonina, Monochaetia, Phomopsis, Phyllosticta, Rhizoctonia, Rhytisma, Septoria, Sphaeropsis, Taphrina, Venturia
Narcissus	Botrytis, Sclerotinia
Oak (red group only)	Cephalosporium, Cercospora, Cladosporium, Cronartium, Elsinoe, Fusarium, Gloeosporium, Gnomonia, Marssonina, Phyllosticta, Septoria, Taphrina, Venturia
Palm, Areca	Alternaria, Cercospora, Colletotrichum, Phomopsis, Phyllosticta, Septoria
Pansy	Alternaria, Botrytis, Cercospora, Colletotrichum, Peronospora, Phyllosticta, Ramularia, Rhizoctonia
Peach, ornamental	Cercospora, Cladosporium, Coryneum, Fusarium, Glomerella, Monilinia, Mycosphaerella, Phomopsis, Phyllosticta, Taphrina
Peperomia	Colletotrichum, Gloeosporium, Rhizoctonia
Petunia	Cercospora, Puccinia, Rhizoctonia, Stemphylium
Philodendron	Gloeosporium, Colletotrichum
Phlox	Ascochyta, Botrytis, Cercospora, Colletotrichum, Phyllosticta, Puccinia, Ramularia, Septoria, Stemphylium, Volutella
Photinia	Cercospora, Gloeosporium, Gymnosporangium, Lophodermium, Pestalotia, Phyllosticta, Septoria
Pine	Alternaria, Botrytis, Cronartium, Fusarium, Lophodermium, Monochaetia, Rhizoctonia, Septoria, Sirococcus
Pine, Norfolk Island	Botrytis, Colletotrichum, Cronartium, Cyindrocladium, Fusarium, Lophodermium, Pestalotia, Rhizoctonia, Septoria, Sirococcus
Plum, ornamental	Botrytis, Cercospora, Cladosporium, Coccomyces, Coryneum, Monilinia, Phyllosticta, Taphrina

Poinsettia**	Botrytis, Cercospora, Fusarium, Uromyces
Poplar	Cercospora, Ciborinia, Colletotrichum, Cylindrocladium, Fusarium, Marssonina, Melampsora, Mycosphaerella, Phyllosticta, Septoria, Stigmata, Taphrina, Venturia
Prayer plant	Alternaria, Drechslera, Glomerella, Puccinia
Privet	Cercospora, Glomerella, Phomopsis, Phyllosticta, Ramularia
Pyracantha	Botrytis, Cercospora, Diplodia, Phomopsis, Phyllosticta, Sphaeropsis
Quince, flowering	Cercospora, Fabraea, Gymnosporangium, Septobasidium
Rhododendron	Alternaria, Cercospora, Coryneum, Gloeosporium, Glomerella, Guignardia, Lophodermium, Mycosphaerella, Pestalotia, Phomopsis, Rhizoctonia, Septoria, Venturia
Rose ¹	Alternaria, Bipolaris, Botryosphaeria, Botrytis, Cercospora, Cladosporium, Cylindrocladium, Diplocarpon, Elsinoe, Gloeosporium, Helminthosporium, Leptosphaeria, Monochaetia, Mycosphaerella, Peronospora, Phyllosticta, Septoria
Spirea	Cylindrosporium
Spruce	Ascochyta, Botrytis, Cladosporium, Lophodermium, Rhizoctonia
Statice	Alternaria, Ascochyta, Botrytis, Cercospora, Colletotrichum, Rhizoctonia, Uromyces
Syngonium	Cephalosporium, Erwinia, Fusarium
Tulip	Botrytis
Viburnum	Botrytis, Cercospora, Cladosporium, Helminthosporium, Monochaetia, Phomopsis, Ramularia
Walnut	Cercospora, Cladosporium, Cylindrocladium, Cylindrosporium, Gnomonia
Yucca	Cercospora, Cylindrosporium, Gloeosporium, Puccinia
Zebra plant	Alternaria, Cercospora, Colletotrichum
Zinnia	Alternaria, Botrytis, Cercospora, Rhizoctonia

*Do not exceed 0.9 lb product per 100 gallons per acre on flower spikes.
 **Do not exceed 1.8 lbs product per 100 gallons per acre.
 Do not use this product for the treatment of marigolds due to highly variable plant responses.

GRASSES: SODFARMS (AGRICULTURAL CROP USE)				
For sodfarm applications, follow provisions within the Agricultural Use Requirements box. Harvesting of treated turf is prohibited until 120 hours following application.				
- Do not apply more than 4 applications per year at the maximum rate per acre per application.				
- Do not allow less than a 10-day interval between applications.				
CROP	DISEASE/PEST	RATE (oz or lbs product)	TIMING/INTERVAL	COMMENTS
Sod Farm	Algae	6.0 to 7.2 oz. in 3 to 5 gal/1000 sq. ft. (16.0 to 19.2 lbs in 130-220 gals/acre)	Begin when algae begins to appear. Repeat at 7-day intervals as long as condition persists.	Do not use on grasses grown for seed. Do not use on

¹ Knock Out® and Double Delight roses may be sensitive resulting in damage to foliage under certain growing conditions.

	<p>Copper Spot (<i>Gloeocercospora sorghi</i>), Fusarium Blight (<i>Fusarium spp.</i>), Red Thread (<i>Laetisaria fuciformis</i>), Slime Molds (<i>Mucilago, Physarum, Fuligo</i>)</p>	<p>4.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (11.0 to 26.4 lbs in 130-220 gal/acre) When conditions are unusually favorable for disease, use 6.0 to 9.6 oz./1000 sq. ft. (16.0 to 26.4 lbs/Acre.)</p>	<p>Begin application when disease appears. Repeat at 7-day intervals as long as condition persists.</p>	<p>grasses intended for grazing, such as range or pasture grasses. Do not graze treated areas of feed clippings to livestock.</p>
	<p>Gray Leaf Spot (<i>Pyricularia grisea</i>) Pythium Blight (<i>Pythium spp.</i>)</p>	<p>8.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (22.0 to 26.4 lbs in 130-220 gal/acre)</p>		
	<p>Dollar Spot (<i>Sclerotinia</i>)</p>	<p>6.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (16.0 to 26.4 lb in 130-220 gals/acre) When conditions are unusually favorable for disease, use 6.0 to 9.6 oz./1000 sq. ft. (16.0 to 26.4 lbs/Acre.)</p>		
	<p>Leaf Spot (<i>Helminthosporium spp.</i>) Rhizoctonia solani Brown Patch</p>	<p>3.0 to 4.8 oz. in 3 to 5 gals/1000 sq. ft. (8.0 to 13.2 lbs in 130-220 gals/acre)</p>		
	<p>Fusarium Snow Mold</p>	<p>6.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (16.0 to 26.4 lbs in 130-220 gals/acre) When conditions are unusually favorable for disease, use 6.0 to 9.6 oz./1000 sq. ft. (16.0 to 26.4 lbs/Acre.)</p>	<p>Apply at 2-6 week intervals during winter.</p>	
	<p>Leaf Rust, Stem Rust, Stripe Rust</p>	<p>3.0 to 4.8 oz. in 3 to 5 gals/1000 sq. ft. (8.0-13.2 lbs in 130-220 gals/acre.)</p>	<p>Begin when disease threatens. Repeat at 7 to 10-day intervals as long as disease persists.</p>	

GRASSES: TURF USES (NON-AGRICULTURAL USE)

For use on golf courses, industrial and commercial lawns, and other nonresidential lawns. Do not use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park athletic fields, athletic fields located on or next to schools (i.e. elementary, middle, and high schools), campgrounds, churches, and theme parks. Follow provisions within the Non-Agricultural Use Requirements Box.

Do not apply by chemigation.

Golf Courses: for cool season grasses; greens, tees and aprons – do not apply more than 5 applications per year at a maximum application rate of 26.4 lbs/A per application.

For cool season grasses; fairways – do not apply more than 4 applications per year at a maximum application rate of 26.4 lbs/A per application.

For warm season grasses; greens, tees and aprons – do not apply more than 4 applications per year at a maximum application rate of 26.4 lbs/A per application.

All Other Turf:

- do not apply more than 4 applications per year at a maximum application rate of 26.4 lbs/A per application
- do not allow less than a 10-day interval between applications

CROP	DISEASE/PEST	RATE (oz. or lbs. product)	TIMING/INTERVAL	COMMENTS
Golf courses, industrial (office park), and municipal lawns	Algae	6.0 to 7.2 oz. in 3 to 5 gal/1000 sq. ft. (16.0 to 19.2 lbs in 130-220 gals/acre)	Begin when algae begins to appear. Repeat at 7-day intervals as long as condition persists.	Do not use on grasses grown for seed. Do not use on grasses intended for grazing, such as range or pasture grasses.
	Copper Spot (<i>Gloeocercospora sorghi</i>), Fusarium Blight (<i>Fusarium spp.</i>), Red Thread (<i>Laetisaria fuciformis</i>), Slime Molds (<i>Mucilago</i> , <i>Physarum</i> , <i>Fuligo</i>)	4.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (11.0 to 26.4 lbs in 130-220 gal/acre) When conditions are unusually favorable for disease, use 6.0 to 9.6 oz./1000 sq. ft. (16.0 to 26.4 lbs/Acre).	Begin application when disease appears. Repeat at 7-day intervals as long as condition persists.	Do not graze treated areas of feed clippings to livestock.
	Gray Leaf Spot (<i>Pyricularia grisea</i>)	8.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (22.0 to 26.4 lbs in 130-220 gal/acre)		

CROP	DISEASE/PEST	RATE (oz. or lbs. product)	TIMING/INTERVAL	COMMENTS
	Dollar Spot (<i>Sclerotinia</i>)	6.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (16.0 to 26.4 lb in 130-220 gals/acre) When conditions are unusually favorable for disease, use 6.0 to 9.6 oz./1000 sq. ft. (16.0 to 26.4 lbs/Acre).		
	Leaf Spot (<i>Helminthosporium spp.</i>) Rhizoctonia solani Brown Patch	3.0 to 4.8 oz. in 3 to 5 gals/1000 sq. ft. (8.0 to 13.2 lbs in 130-220 gals/acre)		
	Fusarium Snow Mold	6.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (16.0 to 26.4 lbs in 130-220 gals/acre) When conditions are unusually favorable for disease, use 6.0 to 9.6 oz./1000 sq. ft. (16.0 to 26.4 lbs/Acre).	Apply at 2-6 week intervals during winter.	
	Pythium Blight (<i>Pythium spp.</i>)	8.0 to 9.6 oz. in 3 to 5 gal/1000 sq. ft. (22.0 to 26.4 lbs in 130-220 gals/acre)	Repeat at 5-day intervals or more frequently if conditions are favorable for disease development.	
	Leaf Rust, Stem Rust, Stripe Rust	3.0 to 4.8 oz. in 3 to 5 gals/1000 sq. ft. (8.0-13.2 lbs in 130-220 gals/acre.	Begin when disease threatens. Repeat at 7 to 10-day intervals as long as disease persists.	

CONIFERS INCLUDING CHRISTMAS TREES: Plantations and Nurseries

USE SITE	DISEASES CONTROLLED	RATE LBS/ACRE	DIRECTONS
Conifers (including Christmas trees)	Ascochyta, Alternaria, Botrytis, Cephalosporium, Cladosporium,	By ground: 1 – 2.4 lbs per acre in 100 gallons water	Apply by ground or air at 7 to 10 day intervals in sufficient water and with proper calibration to obtain uniform and thorough coverage of the tree canopy.

	Cronartium, Fusarium, Lophodermium, Melampsora, Monochaetia, Phomopsis, Rhizoctonia, Septoria, Sirococcus, Sphaeropsis	By air: 1 – 2.4 lbs per acre in a minimum of 10 gallons spray per acre	Begin application at first sign of disease. Use the shortest spray interval during periods of frequent rain, when severe disease conditions persist or during periods of rapid plant growth. This product may be used alone or in combination with other fungicides. Do not allow livestock to graze in treated areas. Do not apply Elixir within one week before or after application of oil or an oil-based pesticide. Do not apply to forests.
--	---	---	--

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal

STORAGE: Important-Never allow ELIXIR to become wet during storage. This may lead to certain chemical changes which will reduce the effectiveness of ELIXIR as a fungicide and create vapors which may be flammable. Keep container closed when not in use. Store product in original container only, away from other pesticides, fertilizer, food or feed.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ATTENTION: This product contains chlorothalonil, mancozeb and ETU, chemicals known to the State of California to cause cancer in laboratory animals. ETU is also known to the State of California to cause birth defects or other reproductive harm in laboratory animals.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

**CONDITIONS OF SALE AND LIMITATION OF
WARRANTY AND LIABILITY**

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of United Phosphorus, Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of United Phosphorus, Inc. and Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold United Phosphorus, Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UNITED PHOSPHORUS, INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, United Phosphorus, Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UNITED PHOSPHORUS, INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UNITED PHOSPHORUS, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

United Phosphorus, Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of United Phosphorus, Inc.

Elixir is a pending trademark of United Phosphorus, Inc.
Cuprofix® Ultra Disperss® is a registered trademark of United Phosphorus, Inc.

Rev. 07/21/2014



Supplemental Label

ELIXIR FUNGICIDE

EPA Reg. No. 70506-298

This supplemental label expires on December 31, 2017 and must not be used or distributed after this date.

DIRECTIONS FOR USE For Broccoli, Cabbage, and Peppers

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

The supplemental labeling and the entire ELIXIR FUNGICIDE container label, EPA Registration Number 70506-298, must be in the possession of the user at the time of the application. Read the label affixed to the container for ELIXIR FUNGICIDE before applying. Carefully follow all precautionary statements and applicable use directions. Use ELIXIR FUNGICIDE according to this supplemental labeling is subject to the use precautions and limitations imposed by the label affixed to the container for ELIXIR FUNGICIDE.

CROP	DISEASES CONTROLLED	RATE LBS PRODUCT/ACRE	TIMING/INTERVALS	RESTRICTIONS & COMMENTS
Broccoli	Alternaria Leaf Spot, Downy Mildew	1.6-2.5	Apply by ground, air, or chemigation. Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 15.4 lbs of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI). Do not apply this product with a U boom device. Aerial application of Elixir on broccoli requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R, or P filter, NIOSH approval prefix TC 84-A.

ACCEPTED

10/21/2014

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 70506-298

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071



Supplemental Label

Cabbage	Alternaria Leaf Spot, Downy Mildew	1.6-2.5	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	<p>Do not apply more than 15.4 lbs. of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI).</p> <p>Do not apply this product with a U boom device.</p> <p>Aerial application of Elixir on cabbage requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R, or P</p>



Supplemental Label

				<p>filter, NIOSH approval prefix TC 84-A.</p>
<p>Pepper</p>	<p>Anthrachnose, Cercospora Leaf Spot (Frogeye Spot), Phytophthora Blight, Ripe Rot, Botrytis Leaf Mold</p>	<p>West of the Mississippi River 1.6-2.5</p>	<p>Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.</p>	<p>Do not apply more than 15.4 lbs of product per acre per season. Minimum retreatment interval is 7 days. Mimimum preharvest interval is 7 days (7 day PHI). Do not apply this product with a U Boom device.</p> <p>Aerial application of Elixir on pepper requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R, or P</p>



Supplemental Label

				filter, NIOSH approval prefix TC 84-A.
		East of the Mississippi River 1.6-3.8	Begin applications prior to disease development and when conditions are favorable for disease development. Apply at 7 to 10 day intervals, if needed. Use higher specified rates when conditions favor disease.	Do not apply more than 23.0 lbs of product per acre per season. Minimum retreatment interval is 7 days. Minimum preharvest interval is 7 days (7 day PHI). Do not apply this product with a U Boom device. Aerial application of Elixir on pepper requires that occupational handlers performing mixing/loading operations observe the additional mitigation measures of wearing a particulate respirator with an N,R, or P filter, NIOSH approval prefix TC 84-A.

Read the Limitation of Warranty and Liability on the container labeling before buying or using ELIXIR FUNGICIDE. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the container label.

Rev. 08/15/2014

United Phosphorus, Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406
1-800-438-6071