



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

July 14, 2021

Jeanette Covert
Regional Regulatory Manager
UPL NA Inc.
630 Freedom Business Center, Ste. 402
King of Prussia, PA 19406

Subject: Registration Review Label Mitigation for Bifenthrin
Product Name: BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE
EPA Registration Number: 70506-480
Application Date: 03/09/2021
Decision Number: 571974

Dear Ms. Covert:

The Agency, in accordance with the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Bifenthrin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Page 2 of 2
EPA Reg. No. 70506-480
Decision No. 571974

If you have any questions about this letter, please contact Quinn Gavin by phone at 703-347-0325, or via email at gavin.quinn@epa.gov.

Sincerely,

A handwritten signature in blue ink, appearing to read 'Linda Arrington', with a stylized flourish at the end.

Linda Arrington, Branch Chief
Risk Management and Implementation Branch 4
Pesticide Re-Evaluation Division
Office of Pesticide Programs

Enclosure

RESTRICTED USE PESTICIDE

Toxic to Fish and Aquatic Organisms

For retail sale to and use only by certified applicators, or persons under their direct supervision and only for those uses covered by certified applicator's certification.

Bifenthrin	GROUP	3A	INSECTICIDE
------------	-------	----	-------------

BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE

INSECTICIDE

For control of Insect Pests including imported fire ants and mole crickets on lawns and perimeters around buildings (residential, commercial, industrial, public and institutional) and parks. For Use on Sod Farms and Recreational Areas including Golf Courses and Athletic Fields.

For outdoor use only.

DO NOT USE THIS PRODUCT ON GOLF COURSES AND SOD FARMS IN NASSAU COUNTY OR SUFFOLK COUNTY, NEW YORK.

ACTIVE INGREDIENT:	By Wt.
---------------------------	---------------

Bifenthrin*.....	0.2%
------------------	------

OTHER INGREDIENTS:	99.8%
---------------------------------	--------------

TOTAL:	100.0%
---------------------	---------------

*Cis isomers 97% minimum, trans isomers 3% maximum.

KEEP OUT OF REACH OF CHILDREN CAUTION

See other panels for additional precautionary information

FIRST AID

IF ON SKIN OR CLOTHING Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN: This pesticide is a pyrethroid. If significant quantities have been swallowed, promptly eliminate the ingested quantities from the stomach and intestines. Treatment is symptomatic and supportive. Alcohol, milk, cream, and other digestible fats and oils should NOT be administered as they may increase absorption.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

FOR 24-HOUR EMERGENCY MEDICAL ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident call CHEMTREC 1-800-424-9300

EPA Reg. No. 70506-480
EPA Est. No. 51036-GA-001

NET CONTENTS: _____

Manufactured For:

UPL NA Inc.

630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

ACCEPTED

Jul 14, 2021

Under the Federal Insecticide, Fungicide
and Rodenticide Act as amended, for the
pesticide registered under

EPA Reg. No. 70506-480

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution: Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing.

The following Personal Protective Equipment requirements apply to sod farm use only.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

1. Long sleeved shirt and long pants
2. Waterproof gloves
3. Shoes and socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:

1. Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
2. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish and aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems. Do not apply when weather conditions favor drift from treated areas. Drift and run-off from treated areas may be hazardous to aquatic organisms in neighboring areas.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Apply using ground equipment only; do not apply by air.

For Use around residential sites such as homes, apartments, condominiums, preschools, day care facilities, outbuildings that are not used for agricultural purposes, private (non-commercial) greenhouses etc. Apply no more than 100 pounds of BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE (0.2 lb. active ingredient (a.i.)) per acre per application on residential use sites. For applications near institutional, public, commercial or industrial buildings, parks, recreational areas or athletic fields, use up to 0.4 lb. a.i. per acre per application.

Per acre, apply no more than 200 lbs. BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE (0.4 lbs. a.i.) in a year.

Spot treatments must not exceed two square feet in size (for example, 2 ft. by 1 ft. or 4 ft. by 0.5 ft.).

For soil or foliar applications, do not apply within a 25-feet of lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries, and commercial fish farm ponds.

Do not apply the product into fish pools, ponds, streams, or lakes. Do not apply directly to sewers or storm drains, or to any area like a drain or gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur.

Do not allow the product to enter any drain during or after application.

Do not apply directly to impervious horizontal surfaces such as sidewalks, driveways, and patios except as a spot or crack-and-crevice treatment.

Do not apply or irrigate to the point of runoff.

Do not apply on the following: nurseries, commercial greenhouses, and grass grown for seed.

Apply this product directly to the lawn or garden area. Water treated area as directed on this label. Do not water to the point of run-off.

For applications at rates above 0.34 lb ai/A (170 lbs product/A), users must water the treated area with directions for use on immediately after application.

Do not make applications during rain. Avoid making applications when rainfall is expected before the product has sufficient time to dry (minimum 4 hours). Rainfall within 24 hours after application may cause unintended runoff of pesticide application.

Do not apply when the wind speed is greater than 15 mph.

Treat surfaces to ensure thorough coverage but avoid runoff.

To treat insects harbored in voids and cracks-and-crevices, applications must be made in such a manner to limit dripping and avoid runoff onto untreated structural surfaces and plants.

Application is prohibited directly into sewers or drains, or to any area like a gutter where drainage to sewers, storm drains, water bodies, or aquatic habitat can occur. Do not allow the product to enter any drain during or after application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry to treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

1. Coveralls
2. Waterproof gloves
3. Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standards for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries and greenhouses.

If watering-in of the granules is necessary, do not allow people or pets on treated surfaces until the grass or soil is dry.

Do not touch treated surfaces until dry.

VEGETATIVE FILTER STRIPS

Construct and maintain a vegetative filter strip, according to the width specified below, of grass or other permanent vegetation between the field edge and nearby down gradient aquatic habitat (such as, but not limited to, lakes; reservoirs; rivers; streams; marshes or natural ponds; estuaries; and commercial fish farm ponds).

Only apply products containing (name of pyrethroid) onto fields where a maintained vegetative filter strip of **at least 25 feet** exists between the field edge and where a down gradient aquatic habitat exists. This minimum required width of 25 feet may be reduced or removed under the following conditions:

- For Western irrigated agriculture, a maintained vegetative filter strip of at least 10 feet wide is required. Western irrigated agriculture is defined as irrigated farmland in the following states: WA, OR, CA, ID, NV, UT, AZ, MT, WY, CO, NM, and TX (west of I-35).
 - For Western irrigated agriculture, if a sediment control basin is present, a vegetative filter strip is not required.
- In all other areas, a vegetative filter strip with a minimum width of 25 feet is required, unless the following conditions are met. The vegetative filter strip requirement may be reduced from 25 feet to 15 feet if at least one of the following applies:
 - The area of application is considered prime farmland (as defined in 7 CFR § 657.5).
 - Conservation tillage is being implemented on the area of application. Conservation tillage is defined as any system that leaves at least 30% of the soil surface covered by residue after planting. Conservation tillage practices can include mulch-till, no-till, or strip-till.
 - A functional terrace system is maintained on the area of application. o Water and sediment control basins for the area of application are functional and maintained.
 - The area of application is less than or equal to 10 acres.

For further guidance on vegetated filter strips, refer to the following publication for information on constructing and maintaining effective buffers: Conservation Buffers to Reduce Pesticide Losses. Natural Resources Conservation Services. <https://www.regulations.gov/document?D=EPA-HQ-OPP2008-0331-0175>.

Buffer Zones – Ground Application

- Do not apply within 25 feet of aquatic habitats (such as, but not limited to, lakes, reservoirs, rivers, streams, marshes, ponds, estuaries, and commercial fish ponds).

APPLICATION TO SOD (including Sod Farms, Lawns and Golf Courses) AND PERIMETERS

SOD APPLICATION: Use equipment designed to give uniform coverage of the area to be treated to Broadcast BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE.

The following table provides recommended application rates of this product for control of the respective pests under normal conditions. However, for each pest listed BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE can be applied at a rate up to 0.4 lbs ai/A (200 pounds per acre). When used on residential sites such as lawns, do not apply more than 0.2 abs ai/A (100 pounds) per application.

INSECT	POUNDS PER ACRE OF ACTIVE INGREDIENT	POUNDS PER ACRE OF BIFENTHRIN PRO 0.2% DG GRANULAR	POUNDS PER 1000 SQ.FT. OF BIFENTHRIN PRO 0.2% DG GRANULAR
Armyworms ¹ Cutworms ¹ Sod Webworm ¹	0.1 lbs. a.i./A	50 lbs. per acre	1.15 lbs. per 1000 sq.ft.
Annual Bluegrass Weevil (<i>Hyperodes</i>)(Adult) ² Billbugs (Adult) ³ Black Turfgrass Ataenius (Adult) ⁴ Leafhoppers Mealybugs	0.1 - 0.2 lbs. a.i./A	50 - 100 lbs. per acre	1.15 - 2.3 lbs. per 1000 sq.ft.
Ants Centipedes Chinch Bugs ⁵ Fleas (Adults) Fleas (Larvae) ⁶ Imported Fire Ants (Adult) ⁷ Imported Fire Ants (Mounds) ⁷ Millipedes Mole Cricket (Adult) ⁸ Mole Cricket (Nymph) ⁹	0.2 - 0.4 lbs. a.i./A	100 - 200 lbs. per acre	2.3 - 4.6 lbs. per 1000 sq.ft.
Pillbugs Sowbugs	0.4 lbs. a.i./A	200 lbs. per acre	4.6 lbs. per 1000 sq.ft.

In New York State, this product may NOT be applied to any grass or turf area within 100 feet of a water body (lake, pond, river, stream, wetland, or drainage ditch).

In New York State, do make a single repeat application of BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE if there are signs of renewed insect activity, but not sooner than two weeks after the first application.

¹ **Armyworms, Cutworm, and Sod Webworm** – Immediately after application irrigate with up to 0.1 inches of water in order to release the insecticide from the granule.

² **Annual Bluegrass Weevil (*Hyperodes*) (adult)**: Time applications to control adult insects as they move from overwintering sites and to grass areas. Movement generally starts when *Forsythia* is in full bloom and ends when flowering dogwood is in full bloom. Consult your State Cooperative Extension Service for more information.

³ **Billbug (adult)**: Apply when adult insects are first observed (April and May). Degree-day models are available to determine best application timing. Spring applications will also control over-wintered chinch bugs in temperate regions. Consult your State Cooperative Extension Service for information.

⁴ **Black Turfgrass Ataenius (adult)**: Apply during May and July for control of the first and second generation of insects. Apply in May when Vanhoutte spiraea and horse chestnut are blooming. Apply in July when Rose of Sharon is blooming.

⁵ **Chinch Bug**: Immediately after application irrigate with up to 0.25 inches water in order to release the insecticide from the granule. To optimize treatment of the base of grass plants and the thatch area, where the insects are often located, irrigate the grass area before treatment. Chinch bugs are difficult pests to control in grasses; thus, higher application rates may be required during the summer when both nymphs and adults are present.

⁶ **Flea larvae**: Immediately after application irrigate with up to 0.5 inches water in order to release the insecticide from the granule. Flea larvae are known to develop in the soil in shaded areas that are accessible to animals (including pets).

⁷ **Imported Fire Ant**: To obtain the best result, combine broadcast applications (this will provide control of foraging

workers and newly mated fly-in queens) with mound drenches (this will control existing colonies). When the soil is dry, irrigate before application or use a high volume application. For broadcast treatments, apply up to 0.4 lb ai/A. Mounds should be treated either by diluting 1 teaspoon of **Bifenthrin Pro Multi Insecticide** (EPA Reg. No. 51036-392) formulation per gallon of water and applying 1 to 2 gallons of finished spray per mound or by applying 1/2 cup of **BIFENTHRIN PRO 0.2% DG GRANULAR**. **BIFENTHRIN PRO 0.2% DG GRANULAR** is used, drench the mound with 1 to 2 gallons of water. Spray with sufficient force to break the mound apex and allow the insecticide solution to flow into the ant tunnels. In addition, treat the four foot diameter circle surrounding each mound. For optimal control, treat in cool weather (65 - 80°F) or in early morning or late evening hours.

8Mole Cricket (adult): Apply as late in the day as possible. If the soil is dry, irrigate before application to bring the insects closer to the soil surface (maximizing contact with the product). Immediately after application, water in with up to 0.5 inches of water. Treat grassy areas that receive pressure from adult mole crickets at peak egg hatch for best control of subsequent nymph populations (see below).

9Mole Cricket Nymph: Apply as late in the day as possible. If the soil is dry, irrigate before application to bring the insects closer to the soil surface (maximizing contact with the product). Immediately after application, water in with up to 0.5 inches of water. Treat grassy areas that received intense adult mole cricket pressure in the spring immediately prior to peak egg hatch. Higher application rates and more frequent applications may be required to control larger nymphs later in the year.

When treating perimeters of structures, make the application to the ground from 5-10 feet adjacent to the structure.

For Perimeter Applications:

Application Rate: Lbs./ 1,000 sq.ft.	Band Width In Feet	Lbs. of BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE/ Linear Distance (Feet) Around Structure							
		100	150	200	250	300	400	500	1,000
2.3	5	1.3	1.9	2.5	3.0	3.6	4.8	5.9	11.7
2.3	10	3.0	4.2	5.3	6.5	7.6	9.9	12.2	23.7
3.0	5	1.7	2.5	3.3	3.9	4.7	6.3	7.7	15.2
3.0	10	3.9	5.4	6.9	8.5	9.9	12.9	15.9	30.9
3.5	5	2.0	2.9	3.8	4.5	5.4	7.2	8.9	17.5
3.5	10	4.5	6.3	8.0	9.8	11.4	14.9	18.3	35.5
4.6	5	2.6	3.8	5.0	6.0	7.2	9.6	11.8	23.4
4.6	10	6.0	8.4	10.6	13.0	15.2	19.8	24.4	47.4

NOTE: This table provides BIFENTHRIN 0.2%G GC amounts which include the total area to be treated. This area includes the area of a circle whose radius is the bandwidth (this accounts for the area treated at the four corners of the structure) plus the linear distance treated times the bandwidth.

APPLICATION TO INDUSTRIAL NON-AGRICULTURAL SITES (Such as Industrial Sites, Ball Fields, Parks and Cemeteries)

BROADCAST SPOT APPLICATIONS: To control actively foraging ants (including Imported Fire Ants) uniformly broadcast 2.3 to 4.6 lbs. BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE per 1000 sq. ft. (this rate is equivalent to 0.2 to 0.4 lbs. a.i./Acre). Additional rate recommendations can be found in the footnotes in the Sod Application Section.

ELECTRICAL SITES (Pad Mounted): Apply 1 to one and one-half Tablespoons of BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE per 4 sq. ft of pad in order to reduce or prevent mound building activity by ants. Care must be taken to ensure that the product is evenly distributed. If the existing transformer is filled or partially filled with soil already excavated by ants the fill must be removed prior to application. Once the excess is removed, and prior to application, relevel the soil.

Best results are obtained by treatment in the early Spring or Fall as ants appear to be attracted to transformers in cooler months.

For Broadcast Applications:

Recommended Spreader Settings					
	Application Rate lbs / 1,000 sq. ft.			Pattern Setting	Width
Equipment	1.15	2.30	4.60		
LESCO Broadcast	F	G	I	—	10
Prizelawn PBR-III	E ½	F ½	H ½	—	10
Scotts RBA	G ½	H ½	J ½	3	10
Scotts SR 2000	H ½	I ½	L ½	8	12
Scotts Handy Green	2	3	4	—	10
Scotts Accupro	4 ½	4 ¾	5	8	12
Spyker	3 ½	3 ¾	4 ¼	—	10
Vicon 402*	25	28 ½	38	—	40
Lely W*	3 ¾	4 ¼	5	III-B	52
Cycloneway	1.5	3	5 ¾	—	4
Earthway 2400	8 ½	10 ½	12 ½	—	10

Speed Applied: Vicon 5.6 mph, Lely 4.5 mph

Equipment should always be calibrated for use prior to application. The settings recommended above represent starting points for final calibration. In order to insure continuing delivery of the correct amount of

BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE further adjustments to the settings should be made periodically during application. Consult the Owner's Manual of the equipment being used for calibration instructions.

RESISTANCE

For resistance management, BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE contains a Group 3A insecticide. Any insect/mite population may contain individuals naturally resistant to BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE and other Group 3A insecticides. The resistant individuals may dominate the insect population if this group of insecticides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay insecticide resistance, take the following steps:

- Rotate the use of BIFENTHRIN PRO 0.2% DG GRANULAR INSECTICIDE or other Group 3A insecticides within a growing season, or among growing seasons, with different groups that control the same pest.
- Use tank mixtures with insecticides from a different group that are equally effective on the target pest when such use is permitted. Do not rely on the same mixture repeatedly for the same pest population. Consider any known cross-resistance issues (for the targeted pests) between the individual components of a mixture. In addition, consider the following recommendations provided by the Insecticide Resistance Action Committee (IRAC):
 - Individual insecticides selected for use in mixtures should be highly effective and be applied at the rates at which they are individually registered for use against the target species.
 - Mixtures with components having the same IRAC mode of action classification are not recommended for insect resistance management.
 - When using mixtures, consider any known cross-resistance issues between the individual components for the targeted pest(s).
 - Mixtures become less effective if resistance is already developing to one or both active ingredients, but they still may provide pest management benefits.
 - The insect resistance management benefits of an insecticide mixture are greatest if the two components have similar periods of residual insecticidal activity. Mixtures of insecticides with unequal periods of residual insecticide activity may offer an insect resistance management benefit only for the period where both insecticides are active.
- Adopt an integrated pest management program for insecticide/acaricides use that includes scouting, uses historical information related to pesticide use, crop rotation, record keeping, and which considers cultural, biological and other chemical control practices.
- Monitor after application for unexpected target pest survival. If the level of survival suggests the presence of resistance, consult with your local university specialist or certified pest control advisor.

- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for that specific site and pest problems in your area.

For further information or to report suspected resistance, contact UPL at 1-800-438-6071.

Dealers Should Sell in Original Packages Only.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Do not store this product near fertilizers, seeds, or other pesticides.

Keep this product out of reach of children and animals at all times. Store this product in a cool, dry place in its original container only. Do not expose the product to excessive heat. Open the product container carefully and, after use, seal tightly.

To Contain a Spill: Cover the spilled product to prevent the material from spreading. If the product container is damaged, place the entire container along with the product in a secure receptacle and clearly label the contents.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Paper and plastic bags: Non refillable container. Do not use or refill this container. Completely empty bag into application equipment. Offer for recycling, if available or dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of UPL NA Inc. ("UPL"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer.

UPL warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to UPL, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF UPL IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF UPL, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT UPL'S ELECTION, THE REPLACEMENT OF THE PRODUCT.