



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

5/6/2026

Meshea J. Brodie  
Sr. Regional Regulatory Manager  
UPL NA, Inc.  
630 Freedom Business Center, Ste. 402  
King of Prussia, PA 19406  
Meshea.brodie@upl-ltd.com

Subject: Label Amendment - Registration Review Mitigation for Carboxin  
Product Name: VITAVAX FLOWABLE FUNGICIDE  
EPA Registration Number: 70506-514  
Case Number: 475017  
Application Date: June 25, 2021

Dear Meshea Brodie:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Carboxin and Oxycarboxin Interim Decision, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Bibiana Oe by phone at (202)566-1538, or via email at [oe.bibiana@epa.gov](mailto:oe.bibiana@epa.gov).

Sincerely,

A handwritten signature in cursive script that reads "C. Lavelle".

Candice Lavelle, Branch Chief  
Worker Protection and Intergovernmental Affairs  
Branch  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs

ENCLOSURE: Stamped label

# Vitavax<sup>®</sup> Flowable Fungicide

Active Ingredient: (% by weight)  
 carboxin (5,6-dihydro-2-methyl-N-phenyl-1,4-oxathiin-3-carboxamide)..... 34%\*  
 Other Ingredients:..... 66%  
 Total: ..... 100%

\*Contains 3.34 lbs. active ingredient per gallon, or 400 grams active ingredient per liter.


**KEEP OUT OF REACH OF CHILDREN  
 CAUTION**

FIRST AID	
<b>IF INHALED:</b>  <b>then</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<p><b>EMERGENCY ASSISTANCE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment:  <b>FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671</b>  <b>FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or 1-703-527-3887.</b></p>	
For PRODUCT USE INFORMATION: Call 1-866-761-9397	

**Net Contents:**

Manufactured for:  
 UPL NA, Inc.  
 PO Box 12219  
 Research Triangle Park, NC 27709

EPA REG. NO. 70506-514  
 EPA EST. NO.  
 Rev. 02/25/2026



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Harmful if inhaled. Avoid breathing vapor or spray mist.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical-resistant to this product are listed below.

**All Mixers, Loaders, Applicators and Other Handlers Must Wear:** long-sleeved shirt and long pants; socks plus shoes; chemical-resistant gloves made of barrier laminate, neoprene, viton, PVC, butyl rubber, or nitrile rubber *except* when bagging or sewing bags of treated seeds; and chemical-resistant aprons when mixing, loading or participating in dip treatments.

FOR COMMERCIAL SEED TREATMENT APPLICATIONS TO CORN (FIELD AND SWEET) AND SUCCULENT BEANS (LIMA, SNAP): Workers carrying out activities involving cleaning and maintenance of seed treatment equipment, including but not limited to scraping, wiping, brushing, and vacuuming of the mixing, delivery, or treatment chamber using compressed air and/or pressurized water, must also: Wear a minimum of a NIOSH-approved elastomeric half-mask respirator with organic vapor (OV) cartridges and combination R or P filters; OR a NIOSH-approved full-face respirator with OV cartridges; OR a gas mask with OV canisters; OR a powered air purifying respirator with OV cartridges.

**Respirator fit testing, medical qualification, and training**

Using a program that conforms to OSHA's requirements (see 29 CFR Part 1910.134), employers must verify that any handler who uses a respirator is:

- Fit-tested and fit-checked,
- Trained, and
- Examined by a qualified medical practitioner to ensure physical ability to safely wear the style of respirator to be worn. A qualified medical practitioner is a physician or other licensed health care professional who will evaluate the ability of a worker to wear a respirator. The initial evaluation consists of a questionnaire that asks about medical conditions (such as a heart condition) that would be problematic for respirator use. If concerns are identified, then additional evaluations, such as a physical exam, might be necessary. The initial evaluation must be done before respirator use begins. Handlers must be reexamined by a qualified medical practitioner if their health status or respirator style or use conditions change.

Upon request by local/state/federal/tribal enforcement personnel, employers must provide documentation demonstrating how they have complied with these requirements.

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. When handlers use closed systems in a manner that meet the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **ENGINEERING CONTROLS**

When handlers use closed systems designed by the manufacturer to enclose the pesticide to prevent it from contacting handlers or other people AND the system is functioning properly and is used and maintained in accordance with the manufacturer's written operating instructions, the handlers need not wear the chemical-resistant apron or gloves. However, the chemical-resistant apron and gloves must be immediately available, for use in an emergency, such as a spill or equipment breakdown.

## **ENVIRONMENTAL HAZARDS**

Do not apply to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate or by disposal of wastes. Treated seed exposed on soil surface may be hazardous to birds. Cover or collect spilled seeds.

Carboxin and its degradates are known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of carboxin and carboxin sulfoxide from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

This product may have effects on federally listed threatened or endangered species or their critical habitat in some countries. It is a violation of federal law to kill, harm or harass listed animal species without authorization. To limit the potential for such impacts when using this product, consult and follow the instructions provided in the EPA endangered Species bulletin for the County or Parish in which you are applying the seed. To determine whether your County or Parish has a bulletin, consult <http://www.epa.gov/espp> before each season's use of this product.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 12 hours**.

*Exception:* Once the seeds, corms, or bulbs are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no contact with the soil/media subsurface.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, or treated seed is: coveralls; chemical resistant gloves made of barrier laminate, neoprene, viton or nitrile rubber; shoes plus socks.

## RESISTANCE MANAGEMENT

For resistance management, VITAVAX Flowable Fungicide contains a Group 7 (carboxin) fungicide. Any fungal population may contain individuals naturally resistant to VITAVAX Flowable Fungicide and other Group 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of VITAVAX Flowable Fungicide or Group 7 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact UPL at 1 (800) 438-6071 or at [www.upl-ltd.com](http://www.upl-ltd.com). You can also contact your pesticide distributor or university extension specialist to report resistance.

## PRODUCT INFORMATION

VITAVAX® Flowable Fungicide is a systemic broad-spectrum fungicide for seed treatment that protects against a wide variety of seed- and soil-borne diseases. VITAVAX Flowable Fungicide is effective against loose smuts, covered smut, common bunt, *Rhizoctonia* and other diseases. See the Disease Control table for details. Seed should be sound and well-cured before treatment.

Do not use VITAVAX Flowable Fungicide in combination with other seed treatment products unless compatibility has been verified. Read and follow carefully all label directions of each combination product before use. When using combinations of products, the most restrictive of label limitations and precautions must be followed. Do not tank mix with any product that has a label prohibition against tank mixing.

Do not plant treated seeds by broadcasting to the soil surface. Treated seed exposed on soil surfaces may be hazardous to birds or mammals. Cover or collect seed spilled during loading. If seeds are not thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.

Areas planted with carboxin-treated seed may be replanted immediately with barley, beans (dry shelled and succulent), canola, corn (field, sweet, pop), cotton, oat, onion (bulb), peanuts, rape seed, rice, safflower, soybean, triticale, or wheat. Do not plant any other crop not listed above in the treated area for at least four months after treated seeds are planted.

### **SEED TREATMENT EQUIPMENT**

VITAVAX Flowable Fungicide may be applied with mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. DO NOT apply this product as a planter box or hopper box treatment. Uniform application to seed is necessary to assure best disease protection and optimum performance. Seed should be sound and well cured before treatment. Refer to the label rates below.

VITAVAX Flowable Fungicide may be diluted with water to achieve more uniform seed coverage (depending on rate, treater and crop). When diluted, VITAVAX Flowable Fungicide and water should be mixed

thoroughly into a slurry prior to treating seed. Recalibrate treating equipment to compensate for the addition of water. The amount of dilution water that is most suitable depends on the seed, treater type and temperature. Contact your local UPL representative or supplier for specific recommendations.

## APPLICATION AND DISEASE CONTROL - SEEDS

Crop	Disease	Application Rate (fl.oz./100 lb seed)	Comments
Wheat	Loose smut ( <i>Ustilago tritici</i> ) Common bunt ( <i>Tilletia caries</i> , <i>T. foetida</i> ) Flag smut *( <i>Urocystis agropyri</i> ) Seed rots, damping off and seedling blight caused by <i>Cochliobolus sativus</i>	2 – 3	The higher rate will provide increased protection on highly infected seed. Do not graze or feed livestock on treated areas for 6 weeks after planting. *Use for control of flag smut not registered by the California Department of Pesticide Regulation.
Barley	True loose smut ( <i>Ustilago nuda</i> ) Covered smut ( <i>U. hordei</i> ) False loose smut ( <i>U. nigra</i> ) Seed rots, damping off and seedling blight caused by <i>C. sativus</i>	2 – 3	The higher rate will provide increased protection on highly infected seed. Do not graze or feed livestock on treated areas for 6 weeks after planting.
Oats	Loose smut ( <i>U. avenae</i> ) Covered smut ( <i>U. koleri</i> ) Seed rots, damping off and seedling blight caused by <i>C. sativus</i>	2 – 3	The higher rate will provide increased protection when high disease pressure is expected. Do not graze or feed livestock on treated areas for six weeks after planting.
Triticale	Seed rots, damping off and seedling blight caused by <i>C. sativus</i>	2 – 3	The higher rate will provide increased protection when high disease pressure is expected. Do not graze or feed livestock on treated areas for 6 weeks after planting.
Corn (field & sweet)	<i>Rhizoctonia solani</i> seed rots, damping off and seedling blight	2 - 4	The higher rate will provide increased protection on highly infected seed, or use for control of seed borne head smut as described below.  VITAVAX Flowable Fungicide may be used for application to seed that has been or will be treated with protectant fungicides such as captan or thiram to obtain a wider spectrum of control. Follow the more restrictive labeling of any tank mix partner. Do not tank mix with any product that has a label prohibition against tank mixing. Do not graze or feed livestock on treated areas for 6 weeks after planting.
	Seed-borne head smut ( <i>Sphacelotheca reiliana</i> )	4	Provides systemic control of seedborne head smut. Avoid planting corn under cool, wet conditions, because the incidence of head smut may increase during the first few weeks after planting under such conditions. Do not graze or feed livestock on treated areas for 6 weeks after planting.
Cotton	<i>Rhizoctonia solani</i> seed rots, damping off and seedling blight	4 – 8	Use the higher rate when planting cottonseed early season in cold soils, or to achieve a higher level of control. Mix with enough water to apply mixture (slurry) at the rate of 20 fluid ounces per 100 pounds of seed. Do not graze or feed livestock on cotton hay grown from treated seed.

†Where a rate range is stated, the lower specified rate should be used where a given seed lot or field is known to have low disease populations or potential for low disease. The higher specified rate should be used where a given seed lot or field is known to have high disease populations or potential for high disease.

Soybeans	<i>Rhizoctonia solani</i> seed rots and seedling blight	3 - 4	The higher rate will provide increased protection when high disease pressure is expected. Do not graze or feed livestock on soybean forage or hay grown from treated seed.
Safflower*	<i>Rhizoctonia solani</i> seed rots, damping off and seedling blight	2	*Use on safflower not registered by the California Department of Pesticide Regulation
Beans	<i>Rhizoctonia solani</i> seed rots, damping off and seedling blight	3 - 4	The higher rate will provide increased protection when high disease pressure is expected. Do not allow livestock to graze or feed on bean forage until 60 days after treated seed is planted.

### APPLICATION AND DISEASE CONTROL – ORNAMENTAL BULBS AND CORMS\*

Crop	Disease	Dip Mixture (Quarts VITAVAX Flowable /100 gallons water)	Comments
Easter Lilies *	<i>Rhizoctonia</i> root rot	1.5 - 3	Dip bulbs for at least 5 to 15 minutes in solution. A sticking agent may be added. Maintain good agitation in dip tanks. Use higher rate for increased disease control.
Other Ornamental Bulbs / Corms *	<i>Rhizoctonia</i> root rot	1.5 - 3	A preliminary trial must be conducted prior to large scale dipping to determine individual plant tolerance. Dip bulbs for at least 5 to 15 minutes in solution. A sticking agent may be added. Maintain good agitation in dip tanks. Use higher rate for increased disease control.

#### USE PRECAUTIONS AND RESTRICTIONS FOR BULB AND CORM DIPS:

- Do not mix insecticides in this dipping solution.

\*Use on bulbs and corms not registered by the California Department of Pesticide Regulation.

#### COMPATIBILITY

VITAVAX Flowable Fungicide can be used in combination with other flowable seed protectant fungicides and insecticides. Consult a seed treatment specialist regarding combining of products with VITAVAX Flowable Fungicide for specific seed crop applications and slurry compatibility. Pretest tank mixes to determine physical compatibility between formulations, and to assure there will be no adverse phytotoxic reactions.

When using combinations of products, the most restrictive of label limitations and precautions must be followed. Do not tank mix with any product that has a label prohibition against tank mixing.

#### USE RESTRICTIONS

When using formulations that do not contain dye, to comply with 21CFR, Part 2.25, treated seed must be colored with an approved dye (40 CFR, Part 153.155) to distinguish and prevent subsequent and inadvertent use as food for man or feed for animals. Treated seed must not be used for or mixed with food or animal feed, or processed for oil. Treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Do not store excess treated seeds beyond planting time.

#### TREATED SEED LABELING

Seed, bulbs or corms treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

“This bag contains seed, bulbs or corms treated with carboxin. Treated Seed, Bulbs or Corms - Do Not Use for Food, Feed or Oil Purposes. Care must be exercised in the handling of treated seed.

When opening this bag or loading/pouring the treated seeds, bulbs or corms, wear long-sleeved shirt, long pants, shoes, socks and chemical resistant gloves.

This product may have effects on federally listed threatened or endangered species or their critical habitat in some counties. It is a violation of federal law to kill, harm or harass listed animal species without authorization. To limit the potential for such impacts when using this product, consult and follow the instructions provided in the EPA endangered Species bulletin for the County or Parish in which you are applying the seed. To determine whether your County or Parish has a bulletin, consult <http://www.epa.gov/espp> before each season's use of this product.

Do not plant treated seeds by broadcasting to the soil surface. Treated seeds exposed on soil surface may be hazardous to birds or mammals. Cover or collect spilled seeds during loading. If seeds are not thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.

Areas planted with carboxin-treated seed may be replanted immediately with barley, beans (dry shelled and succulent), canola, corn (field, sweet, pop), cotton, oat, onion (bulb), peanuts, rape seed, rice, safflower, soybean, triticale, or wheat. Do not plant any other crop not listed above in the treated area for at least four months after treated seeds are planted.

Carboxin and its degradates are known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow. This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of carboxin and carboxin sulfoxide from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

After the seeds, bulbs or corms have been planted, do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. Exception: Once the seeds, bulbs or corms are planted in soil or other planting media, the Worker Protection Standard allows workers to enter the treated area without restriction if there will be no worker contact with the seeds or soil/media subsurface.

Do not graze or feed livestock on treated areas for six weeks after planting.

Exceptions: Do not allow livestock to graze or feed on bean forage until 60 days after treated seed is planted. Do not graze or feed livestock on soybean forage or hay grown from treated seed.

Do not graze or feed livestock on cotton hay grown from treated seed.”

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container only. Store in a secure location that is cool and dry. Do not store in direct sunlight. Protect from extreme temperatures.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL: Nonrefillable container:**

**Plastic Containers:** Do not reuse or refill this container. Triple rinse (or equivalent) container promptly after emptying.

**Triple rinse** as follows:

*For containers small enough to shake:* Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container  $\frac{1}{4}$  full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

*For containers too large to shake:* Empty remaining contents into application equipment or a mix tank. Fill the container  $\frac{1}{4}$  full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Then** offer container for recycling, if available, reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Refillable Container:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**To clean the container before final disposal,** empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times.

**Recycling:** Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or [www.agrecycling.org](http://www.agrecycling.org).

## **IMPORTANT INFORMATION**

### **READ BEFORE USING PRODUCT**

#### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials, or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, UPL NA Inc or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

VITAVAX® is a trademark of a UPL NA Inc.

20260225 Registration Review