



OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 18, 2025

Zack Chesser
Regional Regulatory Manager
UPL NA, Inc.
640 Freedom Business Center, Suite 402
King of Prussia, PA 19406

Subject: PRIA Label Amendment – Add me-too uses on canola, mustard seed, and rapeseed; Metalaxyl Interim Registration Review Decision label mitigation
Product Name: Belmont 2.7 FS
EPA Registration Number: 70506-598
Application Date: 10/26/2023, 6/7/2023
Case Number: 493926, 474187

Dear Zack Chesser:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

The Agency, in accordance with FIFRA, as amended, has completed reviewing all of the information submitted with your application to support the Registration Review of the above referenced product in connection with the Metalaxyl/Mefenoxam Interim Registration Review Decision, and has concluded that your submission is acceptable.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the

Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6.


If you have any questions, please contact Stephanie Suarez at suarez.stephanie@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Manjula Unnikrishnan". The signature is written in a cursive, flowing style.

Manjula Unnikrishnan
Product Manager 21
Fungicide Branch
Registration Division
Office of Pesticide Programs

Enclosure



Belmont 2.7 FS
Fast track amendment: IRRD V2 and Canola Addition Changes V5
Clean copy
July 16, 2025

[Text in brackets is optional]

| | | | |
|-----------|-------|---|-----------|
| Metalaxyl | Group | 4 | Fungicide |
|-----------|-------|---|-----------|

Belmont™-2.7 FS

Seed Treatment Fungicide
for control of seed rot and damping-off diseases of certain crops

| ACTIVE INGREDIENT: | % BY WT. |
|-------------------------------------|----------|
| metalaxyl (CAS No. 57837-19-1)..... | 28.98%* |
| OTHER INGREDIENTS..... | 71.02% |
| TOTAL: | 100.00% |

*Contains 2.65 lbs. metalaxyl per gallon, or 317 grams per liter.

KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you DO NOT understand the label, find someone to explain it to you in detail.)

See [back][side] Panel for First Aid Instructions and [Leaflet][Booklet] for Complete Precautionary Statements and Directions for Use.

| FIRST AID | |
|---|---|
| IF ON SKIN OR CLOTHING: | <ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice. |
| IF SWALLOWED: | <ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• DO NOT induce vomiting unless told to do so by a poison control center or doctor.• DO NOT give anything by mouth to an unconscious person. |
| IF IN EYES: | <ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671. FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887. | |
| For Product Use Information Call 1-800-438-6071 | |

EPA Reg. No. 70506-598
UPL NA, Inc.

EPA Est. No.:
NET CONTENTS:

P.O. Box 12219
Research Triangle Park, NC 27709-2119

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. Causes skin irritation. Harmful if swallowed. Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes. **DO NOT** get on skin or on clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE):

Some materials that are chemical-resistant to this product are listed below.

Applicators and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants;
- Socks and chemical-resistant footwear;
- Chemical-resistant gloves made of (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber > 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, and/or viton ≥ 14 mils).
- When mixing and loading, and when cleaning equipment, wear a chemical-resistant apron.

NOTE: Persons involved in bagging treated seed, sewing or moving bags of treated seed, or cleaning up bagging areas or seed treatment equipment are pesticide handlers and must wear the PPE required on this label for pesticide handlers.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing them. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. **DO NOT** apply when weather conditions favor drift from treated areas. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

DO NOT apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and green-houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the **restricted entry interval (REI) of 24 hours**. Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no worker contact with the soil sub-surface or treated seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- coveralls
- socks and chemical-resistant footwear
- Chemical-resistant gloves; and protective eyewear

PRODUCT INFORMATION

BELMONT™2.7 FS is a systemic fungicide seed treatment specifically for control of systemic downy mildews, *Pythium* and *Phytophthora* spp. For control of other soilborne diseases, use in combination with other seed treatment fungicides.

VITAVAX®-34 (carboxin), VITAFLO®-280 (carboxin + thiram), RANCONA® 3.8 FS (ipconazole), RANCONA® SUMMIT (ipconazole + metalaxyl), RANCONA® V 100 PRO FS (ipconazole + carboxin) and RANCONA® V RTU (ipconazole + carboxin + metalaxyl) are compatible with **BELMONT 2.7FS**. **DO NOT** use **BELMONT 2.7 FS** in combination with other seed treatment products unless compatibility and safety to the crop has been verified. **DO NOT** mix with any pesticide that has a prohibition against tank-mixing. Observe the most restrictive of labeling limitations and precautions of all products used in mixtures.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

BELMONT 2.7 FS may be applied on its own, as water-based slurry, or in combination with other registered seed treatment insecticides and fungicides through standard slurry or mist-type commercial seed treatment equipment.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, METALAXYL 317 FS contains a Group 4 fungicide. Any fungal population may contain individuals naturally resistant to METALAXYL 317 FS and other Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one of more of the following steps:

- Rotate the use of METALAXYI 317 FS or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance- management and/or IPM recommendations for specific crops and pathogens.
- For further information or to repost suspected resistance contact United Phosphorus, Inc. at 610-491-2800. You can also contact your pesticide distributor or university extension specialist to report resistance.

Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season. **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

NOTE: **BELMONT 2.7 FS** is a systemic fungicide with a specific mode of action. **BELMONT 2.7 FS** could be subject to development of resistant strains of fungi. Development of resistance cannot be predicted. Therefore, the company cannot assume liability for crop damage resulting from resistant strains of fungi. Consult your State Agricultural Experiment Station or Extension Service Specialist for guidance and ways to control any possible **BELMONT 2.7 FS** resistant strains of fungi which may occur.

FAILURE TO FOLLOW DIRECTIONS ON THIS LABEL MAY RESULT IN CROP INJURY OR POOR DISEASE CONTROL.

USE RESTRICTIONS

- Seed treated commercially or for sale must be dyed an un-natural color using an EPA approved colorant before being sold. When using formulations that **DO NOT** contain dye, to comply with 40 CFR 153.155, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man or feed for animals.
- IN CALIFORNIA: ONLY for use in commercial seed-treatment establishments. NOT allowed for "on-farm" use in California.
- IMPORTANT: MAXIMUM USAGE when applying multiple metalaxyl and mefenoxam containing products to the same crop within the same season: **DO NOT** apply more than the maximum seasonal total for the active ingredient as stated on the label of the lowest seasonal total on that crop.
- **DO NOT** use other metalaxyl or mefenoxam products for control of Pythium in water-seeded rice.
- **DO NOT** apply **BELMONT 2.7 FS** to spinach which is destined to be grown in the greenhouse.

As stated by regulations specified in 40 CFR 153.155, all seed treated with V-10491 FS must be colored with an EPA-approved dye or colorant to visually distinguish seed treated with a pesticide. This product contains such dye or colorant. This regulation is necessary to prevent treated seed from accidental use as a food for man or feed for animals.

- **DO NOT** use treated seed for feed, food, or oil purposes.
- Store treated seed away from feeds and foodstuffs.

- Seed treated with this product must be labeled in accordance with all applicable requirements of the Federal Seed Act.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- **DO NOT** allow children, pets, or livestock to have access to treated seed.
- If excess treated seed remains after planting, dispose of it by burying away from bodies of water.
- The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb ai/A per calendar year or the lesser value listed in the crop use section.
- **DO NOT** harvest mustard greens. Rape green seeds grown and harvested from BELMONT 2.7 FS treated seed are not for human or feed consumption. Rapeseed grown and harvested from BELMONT 2.7 FS treated seed is only for industrial uses and cannot be used for edible oil or any other human consumption or animal feed.

SEED CONTAINER LABEL REQUIREMENTS

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with metalaxyl fungicide.
- **DO NOT** use treated seed for feed, food or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with metalaxyl:

- Store treated seed away from food and feedstuffs.
- When opening this bag or loading/pouring the treated seed/seed-pieces, wear long sleeved shirt, long pants, shoes, socks, chemical-resistant gloves.
- Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate water bodies when disposing of planting equipment wash waters.
- **DO NOT** allow children, pets or livestock to have access to treated seeds.
- **DO NOT** use other metalaxyl or mefenoxam products for control of pythium in water-seeded rice.
- **DO NOT** apply metalaxyl to spinach which is destined to be grown in the greenhouse.
- The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb ai/A

Groundwater Advisory: Metalaxyl has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow. Therefore, contain any product spills or equipment leaks and dispose of wastes according to the disposal instructions on this label.

APPLICATION INSTRUCTIONS

| Crop | Application Rate Per 100 lbs of seed | Diseases Controlled | Restrictions |
|--|--|--|---|
| Cotton | 0.75 to 1.5 fl oz (0.016 to 0.032 lb ai) To deliver 0.0156 - 0.0312 mg ai per seed, apply 0.167 - 0.333 fl oz of product per 100,000 seeds (Based on average of 4,500 seeds per lb) | <i>Pythium</i> seed rot and damping-off | DO NOT apply by seed treatment more than 0.004 lb metalaxyl/A per year. |
| Soybeans | 0.75 to 1.5 fl oz (0.016 to 0.032 lb ai) To deliver 0.0251 - 0.0502 mg ai per seed, apply 0.038 - 0.075 fl oz of product per 140,000 seeds (Based on average of 2,800 seeds per lb) | <i>Pythium</i> damping-off and early season <i>Phytophthora</i> | DO NOT apply by seed treatment more than 0.0166 lb metalaxyl/A per year. |
| Peanuts[*] [*Not registered for use by California] | 0.75 fl oz (0.016 lb ai) To deliver 0.1004 mg ai per seed, apply 1.071 fl oz of product per 100,000 seeds (Based on average of 700 seeds per lb) | <i>Pythium</i> seed rot and damping-off | |
| Rice[*] [*Not registered for use by California] | 0.75 to 1.5 fl oz (0.016 to 0.032 lb ai) To deliver 0.0037 - 0.0075 mg ai per seed, apply 0.040 - 0.080 fl oz of product per 100,000 seeds (Based on average of 18,800 seeds per lb) | <i>Pythium</i> damping-off | Use Restriction: DO NOT use other metalaxyl or mefenoxam containing products in water-seeded rice. Containers or bags containing BELMONT 2.7 FS treated rice must be labeled with this restriction |
| Sorghum / milo Note: Treat only high quality seed (at least 90% germination) with BELMONT 2.7 FS . | 0.375 to 0.75 fl oz (0.008 to 0.016 lb ai) To deliver 0.0020 - 0.0039 mg ai per seed, apply 0.021 - 0.042 fl oz of product per 100,000 seeds (Based on average of 18,000 seeds per lb) | <i>Pythium</i> damping-off | |

| Crop | Application Rate Per 100 lbs of seed | Diseases Controlled | Restrictions |
|---|---|--|--------------|
| | 0.75 to 1.5 fl oz (0.016 to 0.032 lb ai) To deliver 0.0039 - 0.0078 mg ai per seed, apply 0.042 - 0.083 fl oz of product per 100,000 seeds (Based on average of 18,000 seeds per lb) | Systemic downy mildew (races susceptible to chemical control, and cultivars which possess sorghum downy mildew resistance) | |
| | 1.5 to 3.0 fl oz (0.032 to 0.063 lb ai) To deliver 0.0078 - 0.0156 mg ai per seed, apply 0.083 - 0.167 fl oz of product per 100,000 seeds (Based on average of 18,000 seeds per lb) | Systemic downy mildew (races susceptible to chemical control, and cultivars which DO NOT possess sorghum downy mildew resistance) | |
| Sunflowers (oil and confection type) | 1.5 to 3.0 fl oz (0.032 to 0.063 lb ai) For oil type: To deliver 0.0201 - 0.0402 mg ai per seed, apply 0.214 - 0.429 fl oz of product per 100,000 seeds (Based on average of 7,000 seeds per lb) For confection type: To deliver 0.0402 - 0.0804 mg ai per seed, apply 0.429 - 0.857 fl oz of product per 100,000 seeds (Based on average of 3,500 seeds per lb) | Systemic downy mildew | |
| Beans (except soybeans): Field beans, green beans, chick- peas (garbanzos), kidney beans, lima beans, lupines, navy beans, okra, pole beans, snap beans, string beans, wax beans, and lentils. | 0.75 fl oz (0.016 lb ai) To deliver 0.0439 mg ai per seed, apply 0.469 fl oz of product per 100,000 seeds (Based on average of 1,600 seeds per lb) | <i>Pythium</i> seed rot and damping-off and early season <i>Phytophthora</i> | |
| Peas: Black-eyed peas, cowpeas, field peas, garden peas | 0.75 fl oz (0.016 lb ai) For small size seeds (average of 3,300 seeds per lb): To deliver 0.0213 mg ai per seed, apply 0.227 fl oz of product per 100,000 seeds For larger size seeds (average of 2,100 seeds per lb): To deliver 0.0338 mg.ai. per seed, apply 0.361 fl oz of product per 100,000 seeds | <i>Pythium</i> seed rot and damping-off and early season <i>Phytophthora</i> | |

| Crop | Application Rate Per 100 lbs of seed | Diseases Controlled | Restrictions |
|---|---|--|--------------|
| | 1.5 fl oz (0.032 lb ai) For field peas: To deliver 0.0426 mg ai per seed, apply 0.455 fl oz of product per 100,000 seeds (Based on average of 3,300 seeds per lb) For garden peas: To deliver 0.0676 mg ai per seed, apply 0.721 fl oz of product per 100,000 seeds (Based on average of 2,080 seeds per lb) | Systemic downy mildew and <i>Pythium</i> damping-off | |
| Corn, sweet | 0.75 fl oz (0.016 lb ai) To deliver 0.0223 mg ai per seed, apply 0.238 fl oz of product per 100,000 seeds (Based on average of 3,150 seeds per lb) | <i>Pythium</i> damping-off | |
| | 1.5 fl oz (0.032 lb ai) To deliver 0.0446 mg ai per seed, apply 0.476 fl oz of product per 100,000 seeds (Based on average of 3,150 seeds per lb) | Systemic downy mildew | |
| Corn, field | 0.75 fl oz (0.016 lb ai) To deliver 0.0419 mg ai per seed, apply 0.446 fl oz of product per 100,000 seeds (Based on average of 1,680 seeds per lb) | <i>Pythium</i> damping-off | |
| Corn, pop | 0.75 fl oz (0.016 lb ai) To deliver 0.0230 mg ai per seed, apply 0.245 fl oz of product per 100,000 seeds (Based on average of 3,060 seeds per lb) | <i>Pythium</i> damping-off | |
| Cereal Grains: Wheat, barley, oats, rye and triticale | 0.75 fl oz (0.016 lb ai) For wheat, barley, oats: To deliver 0.0054 mg ai per seed, apply 0.058 fl oz of product per 100,000 seeds (Based on average of 13,000 seeds per lb) For rye / triticale: To deliver 0.0037 mg ai per seed, apply 0.040 fl oz of product per 100,000 seeds (Based on average of 18,850 seeds per lb) | <i>Pythium</i> damping-off | |

| Crop | Application Rate Per 100 lbs of seed | Diseases Controlled | Restrictions |
|--|---|--------------------------------|--|
| Millet | 0.75 fl oz (0.016 lb ai) To deliver 0.0009 mg ai per seed, apply 0.009 fl oz of product per 100,000 seeds (Based on average of 80,000 seeds per lb) | <i>Pythium</i> damping- off | |
| Buckwheat | 0.75 fl oz (0.016 lb ai) To deliver 0.0054 mg ai per seed, apply 0.058 fl oz of product per 100,000 seeds (Based on average of 13,000 seeds per lb) | <i>Pythium</i> damping- off | |
| Canola, (cultivars, varieties and/or hybrids, including Brassica napus, Brassica rapa, and Brassica juncea) [*] Mustard Seed[*] Rapeseed[*] [*Not registered for use by California] | 0.25-0.50 fl. oz. (0.005 lb ai to 0.01 lb ai) To deliver 0.00023 – 0.00047 mg ai per seed, apply 0.0025 – 0.005 fl. oz. of product per 100,000 seeds. (Based on an average of 100,000 seeds per pound.) | <i>Pythium</i> damping- off | Use Restriction: DO NOT harvest mustard greens. Rape green seeds grown and harvested from BELMONT 2.7 FS treated seed are not for human or feed consumption. Rapeseed grown and harvested from BELMONT 2.7 FS treated seed is only for industrial uses and cannot be used for edible oil or any other human consumption or animal feed. |
| Cucumber | 0.75 fl. oz. (0.016 lb ai) | <i>Pythium</i> damping- off | |

| Crop | Application Rate Per 100 lbs of seed | Diseases Controlled | Restrictions |
|---|--|--|---|
| Spinach ¹ | 0.1 to 0.75 fl oz (0.0021 to 0.016 lb ai) | <i>Pythium</i> damping-off | ¹ DO NOT apply to spinach seed to be grown in greenhouses. Containers or bags containing BELMONT 2.7 FS –treated spinach seed must be labeled with this restriction. |
| Forage Legumes: Alfalfa, clover, lespedeza, beans (for forage), soybeans (for forage), soybean hay, peanuts (for forage), peanut hay, peas (for forage), pea vine hay, cowpeas (for forage), cowpea hay, trefoil, vetch and velvet beans (for forage). | 0.75 to 1.5 fl oz (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off and early season <i>Phytophthora</i> | |
| Root and Tuber Vegetables[*]: Beets (garden and sugar), carrot [*Not registered for use by California] | 0.75 fl oz (0.016 lb ai) | <i>Pythium</i> damping-off | |
| Turf Grasses: On golf courses, home lawns, reclamation and erosion control | 1.5. fl oz (0.032 lb ai) | <i>Pythium</i> damping-off | |
| Forage Grasses | 0.75 fl oz (0.016 lb ai) | <i>Pythium</i> damping-off | |
| Use Restriction The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb ai/A | | | |

SEED TREATMENT DIRECTIONS:**USE AT REDUCED RATES IN COMBINATION WITH OTHER FUNGICIDES**

BELMONT 2.7 FS systemic fungicide seed dressing may be applied at the reduced rates shown in the table below and still provide good *Pythium* control when used in combination with broad spectrum seed treatment fungicides. Apply as a commercial seed treatment only in combination at EPA registered rates of broad-spectrum seed treatment fungicides for which compatibility and crop safety have been verified. **DO NOT** mix with any pesticide that has a prohibition against tank-mixing. **BELMONT 2.7 FS** may be applied on its own or as a water-based slurry through standard slurry or mist- type commercial seed treatment equipment.

| Crop | Application Rate Per 100 lb of seed | Diseases Controlled | |
|--|--|--|--|
| Cereal Grains Wheat, barley, oats, rye | 0.10 to 0.375 fl oz (0.002 to 0.008 lb ai) For wheat, barley, oats: To deliver 0.0007 to 0.0027 mg ai per seed, apply 0.008 to 0.029 fl oz of product per 100,000 seeds (Based on average of 13,000 seeds per lb) For rye: To deliver 0.0005 to 0.0019 mg ai per seed, apply 0.005 to 0.020 fl oz of product per 100,000 seeds (Based on average of 18,850 seeds per lb) | Seed decay and damping-off caused by <i>Pythium</i> | |
| Corn, field | 0.10 to 0.375 fl oz (0.002 to 0.008 lb ai) To deliver 0.0056 to 0.0209 mg ai per seed, apply 0.060 to 0.223 fl oz of product per 100,000 seeds (Based on average of 1,680 seeds per lb.) | Seed decay and damping-off caused by <i>Pythium</i> | |
| Peanuts[*] [*Not registered for use by California] | 0.20 to 0.375 fl oz (0.004 to 0.008 lb ai) To deliver 0.0268 - 0.0502 mg ai per seed, apply 0.286 - 0.536 fl oz of product per 100,000 seeds (Based on average of 700 seeds per lb) | Seed decay and damping-off caused by <i>Pythium</i> | |
| Rice[*] [*Not registered for use in California] | 0.10 to 0.375 fl oz (0.002 to 0.008 lb ai) To deliver 0.0005 - 0.0019 mg ai per seed, apply 0.005 - 0.020 fl oz of product per 100,000 seeds (Based on average of 18,800 seeds per lb) | Seed decay and damping-off caused by <i>Pythium</i> | |

| Crop | Application Rate Per 100 lb of seed | Diseases Controlled | |
|--|--|---|--|
| Soybeans | 0.20 to 0.375 fl oz (0.004 to 0.008 lb ai) To deliver 0.0067 - 0.0126 mg ai per seed, apply 0.10 - 0.19 fl oz of product per 140,000 seeds (Based on average of 2,800 seeds per lb) | <i>Pythium</i> damping- off and early season <i>Phytophthora</i> | |
| Peas | 0.20 to 0.375 fl oz (0.004 to 0.008 lb ai) To deliver 0.0059 - 0.0110 mg ai per seed, apply 0.063 - 0.117 fl oz of product per 100,000 seeds (Based on average of 3,200 seeds per lb) | <i>Pythium</i> seed rot and damping-off | |
| Beans (Field, kidney, navy, lentils) | 0.20 to 0.375 fl oz (0.004 to 0.008 lb ai) To deliver 0.0117 - 0.0220 mg ai per seed, apply 0.125 - 0.234 fl oz of product per 100,000 seeds (Based on average of 1,600 seeds per lb) | <i>Pythium</i> seed rot and damping-off | |
| Sunflowers | 0.10 to 0.375 fl oz (0.004 to 0.008 lb ai) To deliver 0.0017 - 0.0064 mg ai per seed, apply 0.018 - 0.068 fl oz of product per 100,000 seeds (Based on average of 5,500 seeds per lb) | <i>Pythium</i> seed rot and damping-off | |
| Use Restriction The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb ai/A | | | |

SEED TREATMENT DIRECTIONS - FOR EXPORT USE ONLY:

For treatment of the following seed for export only to countries requiring treatment with metalaxyl before importation. The treater is responsible for confirming the acceptability of his treated seed in the country to which he is exporting.

| Crop | Application Rate Per 100 lbs. of seed | Diseases Controlled |
|--|--|----------------------------|
| Brassica (Cole) Head, Stem & Leafy Vegetables | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| Canola, (cultivars, varieties and/or hybrids, including Brassica napus, Brassica rapa, and Brassica juncea) | 0.5 – 1.70 fl. oz. (0.01 to 0.036 lb ai) (32 to 110 ml per 100 kg of seed) | <i>Pythium</i> damping-off |
| Corn | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |

| Crop | Application Rate Per 100 lbs. of seed | Diseases Controlled |
|---|--|---|
| | 3.0 fl. oz. (0.063 lb ai) | Systemic downy mildew and <i>Pythium</i> damping-off |
| Cucurbit Vegetables | 0.75 fl. oz. (0.016 lb ai) | <i>Pythium</i> damping-off |
| Fruiting Vegetables | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| Leafy Vegetables | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| Millet | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| | 2.5 fl. oz. (0.052 lb ai) | Systemic downy mildew and <i>Pythium</i> damping-off |
| Onions | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| Peas | 3.5 fl. oz. (0.074 lb ai) | <i>Pythium</i> damping-off and early season <i>Phytophthora</i> |
| Root and Tuber Vegetables (except potatoes) | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| Sorghum | 0.75 – 1.5 fl. oz. (0.016 to 0.032 lb ai) | <i>Pythium</i> damping-off |
| | 5.25 fl. oz. (0.11 lb ai) | Systemic downy mildew and <i>Pythium</i> damping-off |
| Sunflowers | 9.0 fl. oz. (0.19 lb ai) | Systemic downy mildew |
| Use Restriction The maximum annual application rate for metalaxyl or mefenoxam must not exceed 12.3 lb ai/A | | |

TREATED SEED LABELING:

Seed treated in accordance with the Export Use pattern must be labeled as follows: "FOR EXPORT ONLY – NOT FOR DOMESTIC SALES OR USE".

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool dry location. **DO NOT** store in direct hot sunlight. **DO NOT** store above 90°F for prolonged periods of time. After prolonged storage in intense cold, allow product to return to approximately 75°F before using.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Non-refillable container

Plastic Containers: **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: For containers with capacity equal to or less than 5 gallons or 50 pounds: Empty the remaining

contents into application equipment or a mix-tank and drain for 10 seconds after the flow begins to drip. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume - and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix-tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

For containers with capacities greater than 5 gallons or 50 pounds: Empty the remaining contents into application equipment or a mix-tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to ¼ of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix-tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

Recycling: Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrecycle.org.

Refillable container (>250 gallon)

Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents into application equipment or mix-tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to 1/4 of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

Warranty and Disclaimer Statement

The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Such risks may arise from weather conditions, soil factors, off-target movement, unconventional farming techniques, the presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of UPL NA Inc. ("UPL"), and can cause crop injury, injury to non-target crops or plants, ineffectiveness of the product, or other unintended consequences. All such risks shall be assumed by the user or buyer.

UPL warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions. This warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to UPL, and is subject to the inherent risks described above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL DISCLAIMS ALL OTHER WARRANTIES, EXPRESSED OR IMPLIED, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL, MANUFACTURER, AND SELLER DISCLAIM AND SHALL NOT BE LIABLE FOR ANY SPECIAL, INCIDENTAL, INDIRECT, OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE, HANDLING, APPLICATION, STORAGE, OR DISPOSAL OF THIS PRODUCT OR FOR DAMAGES IN THE NATURE OF PENALTIES, AND THE USER AND BUYER WAIVE ANY RIGHT THAT THEY MAY HAVE TO SUCH DAMAGES. NO AGENT, REPRESENTATIVE OR EMPLOYEE OF UPL IS AUTHORIZED TO MAKE ANY WARRANTY, GUARANTEE OR REPRESENTATION BEYOND THOSE CONTAINED HEREIN OR TO MODIFY THE WARRANTIES CONTAINED HEREIN.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE TOTAL LIABILITY OF UPL, MANUFACTURER, AND SELLER, SHALL BE LIMITED TO THE PURCHASE PRICE PAID, OR AT UPL'S ELECTION, THE REPLACEMENT OF THE PRODUCT.

IMPORTANT: READ BEFORE USE

Treatment of highly mechanically damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond control of UPL, UPL makes no claims of guarantees as to germination of carry-over seed.

BELMONT is a trademark and VITAVAX and RANCONA are registered trademarks of a UPL Corporation Limited Group Company.

©2025 UPL Corporation Limited Group Company.