



## OFFICE OF CHEMICAL SAFETY AND POLLUTION PREVENTION

WASHINGTON, D.C. 20460

September 11, 2025

Arianna Shorey  
Regulatory Consultant for UPL NA, Inc.  
c/o Pyxis Regulatory Consulting Inc.  
535 Dock St, Ste 211  
Tacoma, WA 98402

Subject: Label Amendment - Registration Review Mitigation for Metalaxyl and Ipconazole  
Product Name: Rancona CTS Seed Protectant  
EPA Registration Number: 70506-601  
Application Date: January 19, 2022  
Case Number: 478562

Dear Arianna Shorey:

The Agency, in accordance with the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, has completed reviewing all the information submitted with your application to support the Registration Review of the above referenced product in connection with the Metalaxyl and Ipconazole Interim Decisions, and has concluded that your submission is acceptable. The label referred to above, submitted in connection with registration under FIFRA, as amended, is acceptable.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide, Fungicide, and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling and must be used at your next label printing. You must

submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 12 months from the date of this letter. After 12 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

If you have any questions about this letter, please contact Concepción Rodríguez by phone at 202-566-0820, or via email at [rodriguez.concepcion@epa.gov](mailto:rodriguez.concepcion@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read 'M. Perch', with a long, sweeping horizontal line extending to the right.

Maryam K. Muhammad-Perch, Team Lead  
Risk Management and Implementation Branch 4  
Pesticide Re-Evaluation Division  
Office of Pesticide Programs


ENCLOSURE: Stamped label

{Note to reviewer: [Text] in brackets denotes optional text. {Text} in braces denotes where in the final label text will appear and notes to reviewer.}

IPCONAZOLE	GROUP	3	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE

# RANCONA<sup>®</sup> CTS

## SEED PROTECTANT



Active Ingredients (% by weight):  
ipconazole (CAS No. 125225-28-7).....2.42%\*  
metalaxyl (CAS No. 57837-19-1).....1.94%\*  
Other Ingredients: .....95.64%  
Total:.....100.00%

\*Contains 0.209 lb. ipconazole and 0.167 lb. metalaxyl per gallon.

\*Contains 25 grams ipconazole and 20 grams metalaxyl per liter.

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
IF SWALLOWED:	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor.</li><li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li><li>• Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.</li></ul>
<b>EMERGENCY ASSISTANCE:</b> Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671.</b>	
<b>FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.</b>	
<b>For Product Use Information Call 1-800-438-6071</b>	

{Optional reference statements:} [See [front][back][side][inside] panel for First Aid Instructions and [leaflet][booklet] for complete Precautionary Statements and Directions for Use.] [Peel Down for Directions.] [See inside for First Aid, Precautionary Statements, and complete Directions for Use.] {Box/Case:} [See containers inside for additional Precautionary Statements and complete Directions for Use.] {Container:} [See attached booklet for additional Precautionary Statements and complete Directions for Use.]

### Net Contents:

EPA REG. NO. 70506-601

EPA EST. NO.

[xxxxxxV00x]

[Batch Code will be placed on the container.]

[Manufactured] [Produced] for:

**UPL NA Inc.**

630 Freedom Business Center, Suite 402

King of Prussia, PA 19406

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION**

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and Other Handlers Must Wear:** Long-sleeved shirt and long pants; chemical-resistant gloves, made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, or Viton ≥14 mils; and shoes plus socks. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing.
- As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water. Do not contaminate water when disposing of equipment washwater or rinsate.

Ipconazole may persist in soil, and long-term use in the same field may result in an accumulation of active ingredient.

Groundwater Advisory: Metalaxyl is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

## **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**PRODUCT INFORMATION**

RANCONA® CTS Seed Protectant is a systemic and contact broad-spectrum fungicide for seed treatment that protects against a wide variety of seed borne and soil borne diseases as listed in the Application Instructions table for the specific crops.

RANCONA CTS may be diluted with water to achieve good uniformity of treatment, or mixed with other registered seed treatment products before use in order achieve broader spectrum control. The actual final slurry volume will depend on the type of treater used, other seed treatment products that are used in combination, the seed type being treated and/or the treating facility temperatures. Mix RANCONA CTS and water or other seed treatment products thoroughly into a slurry before treating seed. Use a minimum total slurry application volume of 3 fl oz per 100 lb of seed. Recalibrate treating equipment to apply the targeted slurry volume, taking into account the total treatment and dilution water volumes. Contact your local UPL NA Inc. representative or supplier for specific recommendations.

Do not use RANCONA CTS in combination with other seed treatment products unless their compatibility has been verified. Always read and carefully follow all label directions and precautions of each combination product. When using combinations of products, the most restrictive of label limitations and precautions must be followed. Do not tank mix with any pesticide that has a prohibition against tank mixing. Recalibrate treating equipment to compensate for the addition of the other seed treatment products.

NOTE: **This product is undyed.** The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately colored with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

## **ALWAYS SHAKE, STIR OR OTHERWISE MIX THIS PRODUCT WELL BEFORE USE**

### **RESISTANCE-MANAGEMENT**

For resistance management, RANCONA CTS Seed Protectant contains Group 3 and 4 fungicides. Any fungal/bacterial population may contain individuals naturally resistant to RANCONA CTS Seed Protectant and other Group 3 and 4 fungicides/bactericides. A gradual or total loss of pest control may occur over time if these fungicides/bactericides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide/bactericide resistance, take the following steps:

- Rotate the use of RANCONA CTS Seed Protectant or other Group 3 and 4 fungicides/bactericides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicide/bactericides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide/bactericide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide/bactericide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal/bacterial populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact your UPL NA Inc. retailer, representative or call 1-800-438-6071. You can also contact your pesticide distributor or university extension specialist to report resistance.

### **SEED TREATMENT EQUIPMENT**

Apply RANCONA CTS Seed Protectant using mechanical, slurry, or mist-type seed treating equipment, provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease protection and optimum performance. Seed should be sound and well cured before treatment. Refer to the label rates below.

**Equipment Cleaning:** Following applications, thoroughly rinse seed treatment equipment with water. Do not contaminate water by disposal of equipment washwater or rinsate.

### **USE RESTRICTIONS AND PRECAUTIONS**

- Do not use treated seed for food, feed or oil purposes.
- Care must be exercised in the handling of treated seed. Do not use augers used for handling treated seed to move seed for feed, food or oil processing. Do not re-use packaging from treated seed to handle food or feed products. Dispose of seed packaging or containers in accordance with local requirements.
- IN CALIFORNIA: ONLY for use in commercial seed-treatment establishments. NOT allowed for "on-farm" use.
- Do not use in hopper boxes, planter boxes, or seed boxes.
- For soybeans, beans (dried shelled, edible podded, succulent shelled), and peas (dried shelled, edible podded, succulent shelled) the maximum annual rate must not exceed 12.3 lbs metalaxyl/A.
- Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: Do not apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.

NOTE regarding maximum usage when applying multiple metalaxyl and/or mefenoxam containing products to the same crop within the same season: Do not apply more than the maximum seasonal total for the active ingredient

as stated on the label containing the lowest seasonal total on that crop.

**This product is undyed.** The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately colored with a suitable colorant to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any colorant added to treated seed must be cleared for use under 40CFR, Part 153.155.

#### APPLICATION INSTRUCTIONS

Crop	Application Rate	Diseases Controlled	Diseases Partially Controlled*
Soybeans	1.53 fl oz per 100 lb seed or To deliver 0.0073 mg ai per seed, apply 0.76 fl oz of product per 140,000 seeds (Based on average of 2,800 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> ) Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Phomopsis (Diaporthe)</i> . Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i>	Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)
Beans dried shelled beans edible podded beans succulent shelled beans	1.53 fl oz per 100 lb seed or To deliver 0.0127 mg ai per seed, apply 0.96 fl oz of product per 100,000 seeds (Based on average of 1,600 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> ). Seed rot, damping off and seedling blight, (seed and/or soil borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i> . Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i> Seed borne seed rot, damping off and seedling blight ( <i>Ascochyta</i> spp.)	Seed-borne Anthracnose ( <i>Colletotrichum lindemuthianum</i> )Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)
Peas dried shelled peas edible podded peas succulent shelled peas	1.53 fl oz per 100 lb seed or To deliver 0.0064 mg ai per seed, apply 0.48 fl oz of product per 100,000 seeds (Based on average of 3,200 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> ). Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i> . Seed-borne <i>Botrytis</i> and <i>Sclerotinia</i> Seed borne seed rot, damping off and seedling blight ( <i>Ascochyta</i> spp.)	Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)

Crop	Application Rate	Diseases Controlled	Diseases Partially Controlled*
Wheat (Spring and Winter)	0.92 to 1.53** fl oz per 100 lb seed  or  To deliver 0.0009 - 0.0016 mg ai per seed, apply 0.071-0.118 fl oz of product per 100,000 seeds (Based on average of 13,000 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> )  Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i> )  Loose Smut ( <i>Ustilago tritici</i> )  Common Bunt ( <i>Tilletia</i> <i>caries</i> , <i>T. foetida</i> )	Common root rot ( <i>Cochliobolus sativus</i> )  Crown and foot Rot ( <i>Fusarium</i> spp.)  Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)*
Barley	0.92 to 1.53** fl oz per 100 lb seed  or  To deliver 0.0009 - 0.0016 mg ai per seed, apply 0.071-0.118 fl oz of product per 100,000 seeds (Based on average of 13,000 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> )  Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i> )  True Loose Smut ( <i>Ustilago</i> <i>nuda</i> )  Covered Smut ( <i>U. hordei</i> )  False Loose Smut ( <i>U. nigra</i> )	Common root rot ( <i>Cochliobolus sativus</i> )  Crown and foot Rot ( <i>Fusarium</i> spp.)  Leaf Stripe ( <i>Pyrenophora</i> <i>graminea</i> )  Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)*
Oats	0.92 to 1.53** fl oz per 100 lb seed  or  To deliver 0.0009 - 0.0016 mg ai per seed, apply 0.071-0.118 fl oz of product per 100,000 seeds (Based on average of 13,000 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> )  Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i> )  Loose Smut ( <i>Ustilago</i> <i>avenae</i> )  Covered Smut ( <i>U. segetum</i> var. <i>hordei</i> )	Common root rot ( <i>Cochliobolus sativus</i> )  Crown and foot Rot ( <i>Fusarium</i> spp.)  Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)*
Rye, Triticale	0.92 to 1.53** fl oz per 100 lb seed  or  To deliver 0.0006 - 0.0011 mg ai per seed, apply 0.049-0.081 fl oz of product per 100,000 seeds (Based on average of 18,850 seeds per lb)	Seed rots (seed and soil borne <i>Penicillium</i> and <i>Aspergillus</i> )  Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp., <i>Rhizoctonia solani</i> and <i>Cochliobolus sativus</i> )	Common root rot ( <i>Cochliobolus sativus</i> )  Crown and foot Rot ( <i>Fusarium</i> spp.)  Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)*



Crop	Application Rate	Diseases Controlled	Diseases Partially Controlled*
Sorghum	1.53 fl oz per 100 lb seed or To deliver 0.0011 mgai per seed, apply 0.085 fl oz of product per 100,000 seeds (Based on average of 18,000 seeds per lb)	Seed rot, damping off and seedling blight (seed and/or soil borne <i>Fusarium</i> spp. and <i>Rhizoctonia solani</i> )	Root rot ( <i>Fusarium</i> spp.) Seed rot, damping off and seedling blight ( <i>Pythium</i> spp.)

\*Partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial disease control

\*\*The rate of 1.53 fl oz per 100 lb of seed is needed to provide control of *Pythium* seed rot, damping-off and seedling blight on wheat, barley, oats, rye and triticale.

### TREATED SEED LABELING

Seeds treated with this product that are then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

"This package or bag contains seed which has been treated with ipconazole and metalaxyl.

- Do not use for food, feed or oil purposes. Store away from feeds and foodstuffs.
- Persons opening this bag or package or loading/pouring the treated seed must wear a long-sleeved shirt; long pants; chemical resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, or Viton ≥14 mils; shoes and socks.
- After seeds have been planted, do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. *Exception:* Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil subsurface or treated seed. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: long-sleeved shirt and long pants; chemical-resistant gloves made of barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, or Viton ≥14 mils; shoes plus socks. Treated seeds exposed on soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting. Dispose of all excess treated seed by burying seed away from bodies of water.
- Do not contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements."

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container only. After partial use, replace lid and close tightly. Store in a secure place that is temperate, dry and out of direct sunlight. Avoid excess heat. Do not freeze. **Always shake, stir or otherwise mix this product well before use.**

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. Cover or incorporate spilled treated seeds.

**CONTAINER DISPOSAL:**

#### Nonrefillable Container

**Plastic containers:** Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer container for recycling if available, reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Triple rinse** as follows:

*For containers with capacity equal to or less than 5 gallons:* Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Add water - at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume - and recap. Shake for 30 seconds. Pour rinsate into application



equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

*For containers with capacities greater than 5 gallons:* Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water - at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition, with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

#### **Refillable container**

Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

**To clean the container before final disposal**, empty the remaining contents into application equipment or mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition, with a maximum of 10% of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

### **IMPORTANT INFORMATION**

#### **READ BEFORE USING PRODUCT**

#### **CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, UPL NA Inc or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

Treatment of highly mechanically damaged seed, or seed known to be of low vigor and poor quality, except for the purpose of curative control of existing disease pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond the control of UPL NA Inc., UPL NA Inc. makes no claims or guarantees as to germination of carry-over seed.

UPL NA Inc. and UPL logo are trademarks of UPL Corporation Limited Group Company.  
[RANCONA®] [and] [BELMONT®] [is][are][a] registered trademark[s] of an UPL Corporation Limited Group Company.]

© 2021 UPL Corporation Limited Group Company.

[EPA APPROVAL DATE]