



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505T)
1200 Pennsylvania Ave., N.W.
Washington, D.C. 20460

EPA Reg. Number:

70506-640

Date of Issuance:

11/13/25

NOTICE OF PESTICIDE:

☒ Registration
☐ Reregistration
(under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Rancona VPL Peanut Red

Name and Address of Registrant (include ZIP Code):

UPL NA Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406 USA

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA).

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration/reregistration/registration review of your product when the Agency requires all registrants of similar products to submit such data.
2. The data requirements for storage stability and corrosion characteristics (Guidelines 830.6317 and 830.6320) are not satisfied. You have 18 months from the date of registration to provide these data.

Continues page 2

Signature of Approving Official:

Manjula Unnikrishnan, Product Manager 21
Fungicide Branch, Registration Division (7505T)

Date:

11/13/25

3. Make the following label changes before you release the product for shipment:
 - Revise the EPA Registration Number to read, "EPA Reg. No. 70506-640."
4. Submit one copy of the final printed label for the record before you release the product for shipment.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) lists examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.


The record for this product currently contains the following CSF(s):

- Basic CSF dated 6/15/2023

If you have any questions, please contact Carmen Swinger or at swinger.carmen@epa.gov.

Enclosure

[Text in brackets is optional]



IPCONAZOLE	GROUP	3	FUNGICIDE
CARBOXIN	GROUP	7	FUNGICIDE
METALAXYL	GROUP	4	FUNGICIDE
THIOPHANATE-METHYL	GROUP	1	FUNGICIDE

RANCONA[®] VPL Peanut Red

A Seed Treatment Fungicide For Use On Peanuts

ACTIVE INGREDIENTS:	% BY WT.
Ipconazole (CAS No. 125225-28-7).....	1.03%*
Carboxin (CAS No. 5234-68-4).....	16.58%**
Metalaxyl (CAS No. 57837-19-1).....	0.88%***
Thiophanate-methyl (CAS No. 23564-05-8).....	4.36%****
OTHER INGREDIENTS:	77.15%
TOTAL:	100.00%

*Contains 0.10 lbs ipconazole per gallon (12 grams per liter)

**Contains 1.57 lbs carboxin per gallon (188 grams per liter)

***Contains 0.08 lbs metalaxyl per gallon (10 grams per liter)

****Contains 0.41 lbs thiophanate-methyl per gallon (49 grams/liter)

KEEP OUT OF REACH OF CHILDREN

CAUTION / PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you DO NOT understand the label, find someone to explain it to you in detail.)

See [front][back][side][inside] panel for First Aid Instructions and [Leaflet][Booklet] for complete Precautionary Statements and Directions for Use.

FIRST AID	
IF ON SKIN OR CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center, doctor, or going for treatment. FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671 FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300.	

For Product Use Information: Call 1-800-438-6071

EPA Reg. No. 70506-xxx

EPA Est. No.:
NET CONTENTS:

UPL NA Inc.
630 Freedom Business Center, Suite 402
King of Prussia, PA 19406

PRECAUTIONARY STATEMENTS

CAUTION: Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below.

All Mixers, Loaders, Applicators and other handlers must wear:

- Long-sleeved shirt and long pants;
- Socks plus shoes;
- Chemical-resistant apron when mixing or loading;
- Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, polyvinyl chloride (PVC) \geq 14 mils, and/or viton \geq 14 mils) except when bagging or sewing bags of treated seed; and
- Protective eyewear.

When handlers use closed systems in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4)], the handler PPE may be reduced or modified as specified in the WPS.

User Safety Requirements: Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Treated seed exposed on soil surface may be hazardous to birds. Cover or incorporate spilled treated seeds. **DO NOT** contaminate water when disposing of equipment washwater or rinsate or by disposal of wastes.

Ipconazole may persist in soil, and long-term use in the same field may result in an accumulation of ipconazole.

Carboxin and its degradates are known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having high potential for reaching surface water via runoff for several months or more after application.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of carboxin and carboxin sulfoxide from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DO NOT plant any other crop not listed on this label in the treated area for at least twelve months after treated seeds are planted.

Groundwater Advisory: Metalaxyl in this product is known to leach through soil into groundwater under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge.

DO NOT discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional office of the EPA.

This bag contains seed treated with carboxin. This product may have effects on federally listed threatened or endangered species or their critical habitat in some counties. It is a violation of federal law to kill, harm or harass listed animal species without authorization. To limit the potential for such impacts when using this product, consult and follow the instructions provided in the EPA Endangered Species Bulletin for the County or Parish in which you are applying the seed. To determine whether your County or Parish has a Bulletin consult <http://www.epa.gov/espp> before each season's use of this product.

Physical-Chemical Hazards: **DO NOT** mix or allow coming in contact with Oxidizing agents. Hazardous Chemical reaction may occur.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read all label directions carefully before use.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. **For dry applications, DO NOT enter or allow others to enter the treated area until dusts have settled.** For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

When using formulations that **DO NOT** contain dye or colorant, to comply with 21 CFR Part 2.25 and 40 CFR Part 153.155, treated seed must be colored with an approved dye or colorant (40 CFR Part 180.920) to distinguish and prevent subsequent and inadvertent use as a food for man or feed for animals.

DO NOT plant treated seeds by broadcasting to the soil surface. Treated seed exposed on soil surfaces may be hazardous to birds or mammals. Cover or collect seed spilled during loading. If seeds are not thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the Restricted Entry Interval (REI) of **48 hours**.

Exception: Once the seeds are planted in soil, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area without restriction if there will be no worker contact with the soil subsurface or treated seed.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plants, soil, or water, is:

coveralls and long pants; Chemical-resistant gloves made of any waterproof material (barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride (PVC) ≥ 14 mils, and/or viton ≥ 14 mils) except when bagging or sewing bags of treated seed; protective eyewear; shoes plus socks.

PRODUCT INFORMATION

RANCONA® VPL PEANUT RED is a broad-spectrum, ready to use seed treatment fungicide containing ipconazole, carboxin, metalaxyl and thiophanate-methyl for the control of the listed labeled diseases on peanuts. Seed must be sound and well cured before treatment.

SEED TREATMENT EQUIPMENT

RANCONA VPL PEANUT RED is formulated as a ready to use product to be applied as a liquid, and utilizing equipment specifically designed to apply a liquid seed treatment product to peanut seed provided that the equipment can be calibrated to accurately and uniformly apply the product to seed. Uniform application to seed is necessary to ensure best disease protection and optimum performance. Refer to the label rates in **APPLICATION INSTRUCTIONS**. Contact your local representative or supplier for specific application instructions.

Equipment Cleaning: Following application, thoroughly rinse seed treatment equipment with water.

DO NOT contaminate water by disposal of equipment washwater or rinsate.

APPLICATION INSTRUCTIONS

RANCONA VPL PEANUT RED contains the systemic fungicides ipconazole, carboxin, metalaxyl and thiophanate-methyl for the control or suppression of the diseases listed in the table below.

USE RATE OF RANCONA® VPL Peanut RED						
Use rate based on 600 seeds/lb (30,000 seeds/unit)						
Fluid Ounces or ml of Product				Active Ingredient	Grams of ai/100 kg of seed	mg of ai/seed
100 lbs Seed	1000 Seeds	30000 Seeds	100 kg Seed			
3.2 (94.6 ml)	0.053 (1.57 ml)	1.6 (47.3 ml)	7.05 (208.6 ml)	Carboxin	39.12	0.2957
				Ipconazole	2.45	0.0185
				Thiophanate-methyl	9.86	0.0745
				Metalaxyl	2.07	0.0156

Crop	AI	Disease/Pest Controlled	Diseases/Pest Partially Controlled*
Peanut	ipconazole	Seed rot caused by seed-borne <i>Penicillium</i> , <i>Aspergillus</i> and <i>Rhizopus</i> . Damping-off and seedling blight caused by seed- and/or soil-borne <i>Aspergillus</i> , <i>Rhizoctonia</i> , <i>Fusarium</i> , <i>Sclerotinia</i> , and <i>Pythium</i> . Seedling stage Crown Rot caused by <i>Aspergillus niger</i> .	Early season Stem Rot/ White Mold caused by soil-borne <i>Sclerotium rolfsii</i> , and Black Rot caused by soil-borne <i>Cylindrocladium parasiticum</i> .
	metalaxyl	<i>Pythium</i> seed rot and damping-off control	
	carboxin	<i>Rhizoctonia solani</i> seed rots (seed- and / or soil-borne), damping off and seedling	

		blight	
	thiophanate-methyl	Seed Decay (<i>Aspergillus</i> spp.) Seedling Blight (suppression of <i>Fusarium</i> spp. and <i>Rhizoctonia</i> spp.) Seed borne <i>Sclerotinia</i> spp.	

* Partial control can either mean erratic control from good to poor, or consistent control at a level below that is generally considered acceptable for commercial disease control.

USE RESTRICTIONS

DO NOT apply more than 24 oz. of (0.02 lb ai ipconazole, 0.02 lb ai metalaxyl, 0.29 lb ai carboxin, 0.08 lb ai thiophanate-methyl) **RANCONA VPL PEANUT RED** product per acre per year.

DO NOT graze or feed livestock on treated areas or hay grown from treated peanuts.

DO NOT use treated seed for food, feed or oil purposes. Care must be exercised in the handling of treated seed. Augers used for handling treated seed must not be used to move seed for feed, food or oil processing.

Laboratory and field studies have shown that **RANCONA VPL PEANUT RED** will not adversely affect seed germination. Due to seed quality conditions beyond the control of the manufacturer, no claims are made to guarantee germination of carry-over seed as an untreated seed lot may decline in germination with storage irrespective of seed treatment application. Treatment of highly mechanically damaged seed or seed of known low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. If seed lot quality is unknown, conduct a germination test on a small portion of seed before committing the total seed lot to a selected chemical treatment.

TANK MIX INSTRUCTIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels when using a combination of products. Users must follow the most restrictive directions for use and precautionary statements of each product in the combination mixture.

RESISTANCE MANAGEMENT

For resistance management, please note that **RANCONA VPL PEANUT RED** contains a Group 3 (ipconazole), a Group 7 (carboxin), a Group 4 (metalaxyl) fungicides and a Group 1 (thiophanate-methyl) fungicide. Any fungal population may contain individuals naturally resistant to **RANCONA VPL PEANUT RED** and other Group 3, Group 7, Group 4, or Group 1 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies must be followed.

To delay fungicide resistance, take one or more of the following steps:

- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM instructions for specific crops and pathogens.
- For further information or to report suspected resistance contact UPL NA Inc. at 1-800-438-6071. You can also contact your pesticide distributor or university extension specialist to report resistance.

TREATED SEED LABELING

Seed that has been treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag:

Seed Bag Label Requirements

The Federal Seed Act requires that containers containing treated seeds shall be labeled with the following statements:

- This seed has been treated with **RANCONA VPL PEANUT RED**, a fungicide containing ipconazole,

carboxin, metalaxyl and thiophanate-methyl.

- **DO NOT** use treated seed for feed, food, or oil purposes.

The U.S. Environmental Protection Agency requires the following statements on containers containing seed treated with ipconazole, carboxin, metalaxyl and thiophanate-methyl:

- Store treated seed away from food and feedstuffs.
- **DO NOT** allow children, pets or livestock to have access to treated seeds.
- Wear long pants, long-sleeved shirt and protective gloves when handling treated seed.
- **DO NOT** plant treated seeds by broadcasting to the soil surface. Treated seed exposed on soil surfaces may be hazardous to wildlife. Cover or collect seed spilled during loading. If seeds are not thoroughly incorporated by the planter during planting, additional incorporation may be required to thoroughly cover exposed seeds.
- Treated seeds exposed on the soil surface may be hazardous to wildlife. Cover or collect treated seeds spilled during loading and planting (such as in row ends).
- Manage excess treated seeds (e.g., spilled, unused, or expired treated seeds) by one or more of the following methods:
 - Collect excess treated seeds for reuse for planting.
 - Bury excess treated seeds (only allowed if totaling 1 pound or less) at least 50 feet away from bodies of water at a depth of 6 inches or double the planting depth, whichever is greater.
 - Dispose of excess treated seed by placing them in a landfill in accordance with applicable laws in your state.
 - Excess treated seeds may be returned to the supplier if permitted by the state.
- Dispose of all excess treated seed by burying seed away from bodies of water.
- **DO NOT** contaminate bodies of water when disposing of planting equipment wash water.
- Dispose of seed packaging or containers in accordance with local requirements.
- Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same season: **DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop.
- Excess treated seed may be used for ethanol production if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice.
- Carboxin and its degradates are known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.
- This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water.
- This product is classified as having high potential for reaching surface water via runoff for several months or more after application.
- A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features including ponds, streams, and springs will reduce the potential loading of carboxin and carboxin sulfoxide from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.
- **DO NOT** plant any other crop not listed on this label in the treated area for at least twelvemonths after treated seeds are planted.

DO NOT graze or feed livestock on treated areas or hay grown from treated peanuts.

This product may have effects on federally listed threatened or endangered species or their critical habitat in some counties. It is a violation of federal law to kill, harm or harass listed animal species without authorization. To limit the potential for such impacts when using this product, consult and follow the instructions provided in the EPA Endangered Species Bulletin for the County or Parish in which you are applying the seed. To determine whether your County or Parish has a bulletin, consult

<http://www.epa.gov/espp> before each season's use of this product. Bulletins also may be available from local pesticide dealers, extension offices, or State pesticide agencies.

STORAGE AND DISPOSAL

DO NOT contaminate water, food or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container only. After partial use, replace lid and close tightly. Store in a secure place that is temperate, dry and out of direct sunlight. Due to its viscosity, store this product above 50°F prior to and during application. **DO NOT** store **RANCONA VPL Peanut Red** at low temperatures. PROTECT FROM FREEZING. **DO NOT** apply direct heat to the container. ALWAYS SHAKE, STIR OR OTHERWISE MIX THIS PRODUCT WELL BEFORE USING.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Nonrefillable container. Plastic Containers: **DO NOT** reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Then offer container for recycling, reconditioning, or puncture and dispose of in a sanitary landfill, by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse as follows: For containers with capacity equal to or less than 5 gallons: Empty the remaining contents into application equipment or a mix-tank and drain for 10 seconds after the flow begins to drip. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to 1/4 of the container volume - and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix-tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

For containers with capacities greater than 5 gallons: Empty the remaining contents into application equipment or a mix-tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix-tank or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

Refillable container

Refill this container with pesticide only. **DO NOT** reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents into application equipment or mix-tank. Add water - at least 2% of the container volume, and (if the treating mixture composition permits it) up to 1/4 of the container volume. Replace and tighten closure. Agitate vigorously or recirculate the rinsate with a pump for at least 2 minutes, ensuring that the rinsate rinses the walls of the container. Empty the rinsate into application equipment or rinsate collection system, for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water.

Recycling

Once cleaned, some agricultural plastic pesticide containers can be taken to a container collection site or picked up for recycling. To find the nearest site, contact your chemical dealer or manufacturer, or contact the Ag Container Recycling Council (ACRC) at 1-877-952-2272 (toll free) or www.acrc.org.

**IMPORTANT INFORMATION
READ BEFORE USING PRODUCT**

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

Treatment of highly mechanically damaged seed or seed known to be of low vigor and poor quality may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small test sample of seed before treating commercial quantities with a selected chemical treatment. Due to seed quality and seed storage conditions beyond the control of UPL NA Inc., UPL NA Inc. makes no claims or guarantees as to germination of carry-over seed.

RANCONA is a trademark of a UPL Corporation Limited Group Company
©2023, UPL Corporation Limited Group Company.

20230811 Original Registration Application