

Please read instructions on reverse before completing form.

Approved, OMB No. 2070-0060, Approval expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 70515-3	2. EPA Product Manager Carol Frazer	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) SianaFresh 10 EC	PM# Biochemical Pesticides BR./BPPD	
5. Name and Address of Applicant (Include ZIP Code) Nutra-Park Inc. 3225 Deming Way, Suite 140 Middleton, WI 53562 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application. NOTIFICATION
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below. Date Reviewed: <u>9/23/04</u>

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Reviewed By: apillard

Notification of Additional Brand Name "MT 405".
This notification is consistent with the provisions of PR Notice 98-10 and EPA regulation at 4 CFR 153.46, and no other changes have been made to the labeling or the confidential statement of formula for this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III


1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input type="checkbox"/> Metal	<input checked="" type="checkbox"/> Plastic
* Certification must be submitted	If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container	4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/>			
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Amy Plato Roberts	Title Regulatory Consultant	Telephone No. (Include Area Code) (202) 828-8964
2. Signature 		6. Date Application Received (Stamp)
3. Title Senior Consultant		
4. Typed Name Katherine A. Buteau	5. Date June 23, 2004	
<p>I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.</p>		

91

2 8 15



Technology Sciences Group Inc.
 Sacramento: State Affairs Division
 712 Fifth Street, Suite A
 Davis, California 95616
 (530) 757-1050
 FAX (530) 757-1299
 E-Mail: kbuteau@tsgusa.com

Katherine A. Buteau
 Senior Consultant
 State Affairs Division

June 23, 2004

Dr. Carol Frazer
 Regulatory Action Leader
 Biopesticides and Pollution Prevention Division (7511C)
 Office of Pesticide Programs, EPA
 1921 Jefferson Davis Highway
 Arlington, VA 22202

RE: SignaFresh® 10 EC
EPA Reg. No. 70515-3

Notification of additional brand name

Dear Dr. Frazer:


Enclosed with this letter you will find a notification for an additional brand name for SignaFresh® 10EC.

- 1) Notification form;
- 2) One (1) copy of the revised label with the additional brand name added to the front panel.

This notification is for minor label revisions that are:

- a. Consistent with 40 CFR Part 156; and
- b. Involve no change in the ingredients statement, signal word, use classification, precautionary statements, First Aid, physical / chemical / biological properties, storage and disposal or directions for use.

If you have any questions or comments, please do not hesitate to contact me or Ms. Amy Roberts at our DC office at (202) 828-8964.

Sincerely,


Katherine A. Buteau
 Regulatory Consultant for Nutra-Park, Inc.
 Direct dial (530) 757-1050; email kbuteau@tsgusa.com

cc: Tim Peoples, Nutra-Park, Inc.
 A. Roberts, TSG
 Files

3 7 15

(Front Panel)

SignaFresh® 10EC

(Alternate Brand Names: MT 405™)

- SignaFresh® 10EC is a growth regulator that enhances fruit quality, ripening and shelf life of select fruits, flowers and vegetables.
- With SignaFresh® Technology.
- Quality Enhancer™
- Fresh Focus™
- MasterFresh™

NOTIFICATION

Date Reviewed: 9/23/04
 Reviewed By: C. J. [Signature]

Active Ingredient:

Lysophosphatidylethanolamine (LPE).....	10%
Other Ingredients:.....	90%
Total	100%

Contains 0.751 pounds LPE per gallon

KEEP OUT OF REACH OF CHILDREN
CAUTION

See (side) (back) panel for first aid and additional precautionary statements.

EPA Reg. No.: 70515-3
 EPA Est. No.: XXXXX-XX-X

Manufactured for: Nutra-Park Inc.
 3225 Deming Way, Suite 140
 Middleton, WI 53562

Net Contents: XX

(Back Panel)

FIRST AID	
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-858-7378 for emergency medical treatment information.	

(Precautionary Statements for uses covered by the Worker Protection Standard)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Applicators and other handlers must wear coveralls worn over short-sleeved shirt and short pants, chemical resistant (such as nitrile or butyl) gloves, and shoes plus socks. Follow manufacturer's instructions for cleaning / maintaining PPE. Keep and wash PPE separately from other laundry.

User Safety Recommendations: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

(Precautionary Statements for uses not covered by the Worker Protection Standard)

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Remove contaminated clothing and wash clothing before reuse.

Environmental Hazards: Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean highwater mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

(Directions for Use for uses covered by the Worker Protection Standard)

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry intervals (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is coveralls, waterproof gloves and shoes plus socks.

(Directions for Use for uses not covered by the Worker Protection Standard)

DIRECTIONS FOR USE

It is a violation of Federal law to apply this product in a manner inconsistent with its labeling.

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are **not** within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried or dusts have settled. For post-harvest treated commodities, unprotected persons should not handle until dip has dried.

6 8 15

General:

SignaFresh® 10EC is a growth regulator that enhances fruit quality, ripening and shelf life of select fruits, flowers and vegetables. SignaFresh® 10EC application does not enhance respiration, like other ripening agents. SignaFresh® 10EC is also known to reduce senescence by inhibiting some of enzymes involved in membrane breakdown. For these reasons, SignaFresh® 10EC enhances and promotes shelf life. When applied to physiologically mature fruit (at breaker stage, i.e. early stage of ripening), it stimulates and promotes ripening.

READ ENTIRE LABEL BEFORE USING THIS PRODUCT.

DIRECTIONS FOR USE: SPECIFIED TREATMENTS

Mixing Instructions:

Shake well (1-3 minutes) before mixing. See Pre Harvest Application table below for the application rate (amount of SignaFresh® 10EC fl. oz./acre) specific to your crop. Dilute product with the amount of water necessary to treat the intended area. Add half the water to tank, then add SignaFresh® 10EC. Add remaining water and mix thoroughly.

Pre-Harvest Application Directions:

Apply SignaFresh® 10EC in water to wet fruit uniformly, using conventional ground or aerial application equipment. For better coverage, use a sticker deposition agent. Contact your Nutra-Park Representative for guidance on surfactant or adjuvant selection and use. Do not use with organo-silicone surfactants and use antifoam agents sparingly.

Apply SignaFresh® 10EC approximately two weeks prior to harvest, unless noted otherwise under "Special Instructions" on the following table.

Follow directions for specific crop listed below:

PRE HARVEST APPLICATION					
Crop			Rate in oz/acre	Use	Special Instructions
Fruiting Vegetables					
Eggplants			8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	
Tomatoes	Vine Ripe	Field	8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Uniform color development • Improved fruit firmness 	<ul style="list-style-type: none"> • Apply 10 to 14 days prior to desired use response. • Reapply at 14-day intervals until final harvest.
Tomatoes	Vine Ripe	Greenhouse	2 - 4		<ul style="list-style-type: none"> • Apply 7 to 10 days prior to desired use response. • Reapply at 7 to 14 day intervals until final harvest.
Tomatoes		Processing	15 - 45		<ul style="list-style-type: none"> • Apply approximately two weeks prior to harvest.
Peppers					
	Red Peppers Hot Peppers		27	<ul style="list-style-type: none"> • Accelerated fruit ripening • Increased early marketable yield • Uniform color development* • Improved fruit quality 	<ul style="list-style-type: none"> • Apply at 10 to 20% color break.
	Green Peppers		27		<ul style="list-style-type: none"> • Apply two to three weeks before harvest.

Pimentos	8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	
Cucurbits			
Melons Squash Pumpkins	15 - 45	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Improved fruit quality 	
Berries/Small Fruits			
Raspberries Boysenberries Blackberries Blueberries	8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	
Cranberries	8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	<ul style="list-style-type: none"> • Can be applied through irrigation systems. Consult your Nutra-Park representative for use guidance.
For tank-mixing with chlorothalonil fungicides	8 - 30	<ul style="list-style-type: none"> • Decrease the phytotoxic effect of chlorothalonil containing fungicides 	<ul style="list-style-type: none"> • Apply in tank mixture with the chlorothalonil fungicide. Consult the fungicide label for appropriate rate and timing of application.
Currants	8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	
Strawberries	6- 25	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	<ul style="list-style-type: none"> • Apply 7 to 10 days prior to desired use response. • Reapply at 7 to 14 day intervals until final harvest.

Tree Fruits			
Avocados	8 – 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	
Citrus Grapefruits, Mineolas, Lemons, Limes, Mandarins, Oranges, Pummelos, Tangelos and Tangerines	27	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality • Improved fruit and rind quality 	<ul style="list-style-type: none"> • Apply two to three weeks before to harvest. • Apply four to six weeks before harvest
Pome Fruits Apples	27	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development* • Improved fruit quality 	<ul style="list-style-type: none"> • Apply five to six weeks before harvest.
Crabapples and Pears	27		<ul style="list-style-type: none"> • Apply two weeks before harvest.

Stone Fruits Apricots and Plums	27	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development* • Improved fruit quality 	<ul style="list-style-type: none"> • Apply three weeks before to harvest.
Peaches and Nectarines	14		<ul style="list-style-type: none"> • Apply at pit hardening, approximately three weeks before harvest
Red Cherries	27		<ul style="list-style-type: none"> • Apply three weeks before harvest
Bi-color Cherries	27		<ul style="list-style-type: none"> • Apply seven to 10 days before harvest
Tree Nuts Pistachios, Walnuts, Pecans	8 – 30	<ul style="list-style-type: none"> • Accelerated fruit ripening • Improved fruit quality 	
Vine Fruits			
Grapes Raisin, Wine and Juice, and Table Grapes (such as Flame Seedless, Crimson, Thompson Seedless and Red Globe)	30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development* • Improved fruit quality 	
Kiwi Fruit	8 – 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Improved fruit quality 	

Others - Crops			
Coffee Mangos Pomegranates Pineapples	8 - 30	<ul style="list-style-type: none"> • Increased early marketable yield • Accelerated fruit ripening • Uniform color development • Improved fruit quality 	
Mushrooms	8 - 30	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life 	
Cotton	15 - 45	<ul style="list-style-type: none"> • Acceleration of boll opening. 	<ul style="list-style-type: none"> • Apply when there are sufficient unopened mature bolls to produce desired yields.

* Not intended for uniform color development use on green peppers, pears, apricots or raisin grapes.

Post-Harvest Application Directions:

Apply SignaFresh® 10EC as a washwater dip or as a spray. Prior to sorting and packaging, dip fruit in well-agitated water containing SignaFresh® 10EC for up to 3 minutes or uniformly spray fruit with SignaFresh® 10EC to achieve full coverage.

See Post-Harvest Application Table for the application rate (parts per million) specific to the commodity to be treated and the Dilution Table to determine the amount of SignaFresh® 10EC needed to make solution. For dip applications add SignaFresh® 10EC to washwater, maintain water agitation and submerge fruit in washwater for up to 3 minutes. For use as a spray, dilute SignaFresh® 10EC in water and apply uniformly to cover fruit prior to packing. Maintain chlorine levels and water pH at levels recommended for post-harvest treatments. Allow fruit to dry before packaging. Follow standard handling procedures after treatment.

Dilution Table

Parts Per Million (PPM)	Amount of SignaFresh® 10EC to make 100 gallons of spray solution
25	3.5 fl. oz.
50	7 fl. oz.
100	14 fl. oz.
200	28 fl. oz.
1000	1405 fl. oz.

Follow directions below for crop specific post harvest applications directions for use.

POST HARVEST APPLICATION		
Crop/Commodity	Rate (ppm) for dip and/or spray	Use
Fruiting Vegetables		
Eggplants	25-100 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life
Tomatoes Fresh Market	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Accelerated natural ripening
Peppers	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life

Berries/Small Fruits		
Blueberries Boysenberries Cranberries Currants	25-100 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life
Tree Fruits		
Avocados	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Accelerated natural ripening
Citrus Grapefruits, Mineolas, Lemons, Limes, Mandarins, Oranges, Pummelos, Tangelos and Tangerines	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Accelerated natural ripening
Pome Fruits Apple, Crabapple and Pear	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life
Stone Fruits Peaches, Plums, Cherries, Apricots	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life

Other Crops		
Bananas	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Accelerated natural ripening • Improved fruit quality • Extended shelf life
Mangos Pomegranates Pineapples	50-200 dip 500-1000 spray	<ul style="list-style-type: none"> • Improved fruit quality • Extended shelf life

Use on Cut Flowers: Use SignaFresh® 10EC on freshly cut flowers to extend shelf life, improve blossom life and enhance uniform bud opening. Dilute SignaFresh® 10EC in water at a rate of 1-25 ppm and submerge stems of freshly cut flowers in solution. Renew SignaFresh® 10EC solution if water becomes cloudy. Contact your Nutra-Park representative for use guidance for specific flower varieties.

Chemigation Directions for Use

General Requirements:

- 1) Apply this product only through a sprinkler including a center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set of hand move irrigation and systems. Do not apply this product through any other type of irrigation system.
- 2) Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.

- 3) If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.
- 4) Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- 5) A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make any necessary adjustments should the need arise.

Specific Requirements:

- 1) The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow.
- 2) The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back towards the injection pump.
- 3) The pesticide injection pipeline must contain a functional, normally dosed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- 4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5) The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6) Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being filled with a system interlock.
- 7) Do not apply when wind speed favors drift beyond the area intended for treatment.

Application Instructions:

- 1) Remove scale, pesticide residues, and other foreign matter from the chemical supply tank and entire injector system. Flush with clean water. Failure to provide a clean tank, void of scale or residues may cause product to lose effectiveness or strength.
- 2) Determine the treatment rates as indicated in the directions for use and make proper dilutions.
- 3) Prepare a solution in the chemical tank by filling the tank with the required water and then adding product as required. The product will immediately go into suspension without any required agitation.
- 4) The product is to be applied continuously for the duration of the water application.

<p>STORAGE AND DISPOSAL</p> <p>Do not contaminate water, food or feed by storage or disposal.</p> <p>Storage - Store container in cool place until used. Optimum storage is at refrigerated temperatures (i.e. 33°F to 50°F).</p> <p>Pesticide Disposal - Wastes resulting from use of this product may be disposed of at an approved waste disposal facility.</p> <p>Container Disposal- Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.</p>
--

(Limited Warranty)