

70627-10

1/15

MAR - 3 2009

Ms. Kresti A. Lyddon
Senior Registration Specialist
JohnsonDiversey, Inc.
8310 16th Street, MS 707
Sturtevant, WI 53177-0902

Subject: **Johnson's Forward Cleaner**
EPA Registration Number 70627-10
Your Application Dated: December 2, 2008
EPA Received Data: December 8, 2008

Dear Ms. Lyddon:

The following amendment, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended is acceptable with comments.

Proposed Labeling:

Label amendment in response to PR Notice 2007-4

Label Comments:

1. On page 1, the First Aid Statement should be on front panel.
2. On page 2, revise the second paragraph under (General uses) by deleting the parentheses around "food and non-food contact" and "food contact surfaces".
3. On page 2, revise the second paragraph under (Hospitals/Health Care Facilities) by deleting the phrase (and non-food). Also, delete parentheses around the statement "where disinfection is required".
4. On page 3, revise the second paragraph under (Non-Acid Bowl {& Bathroom} Disinfectant Cleaner:) by clarifying rims as "toilet bowl" rims.
5. On page 4, and 12, revise the "Bactericidal Stability of Use-Solution" claim by updating the nomenclature for the organism, Salmonella choleraesuis to read "~~Salmonella enterica~~".

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							

- 6. On page 5, revise the MOU statement or first paragraph under "Direction For Use" by delete the parentheses around it.
- 7. On page 5, revise the "Pre-Measured Packets Dispensing System" instructions by disclosing all the pre measure pickets sizes.
- 8. On page 6, revise the "For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet bowls from Concentrate" and "For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet bowls {and Urinals} from use Dilution" directions by adding the phrase. "Then Flush", to step.
- 9. On page 7, revise the "Personal Protection" statement under "HIV" directions by deleting the word or replace with "and".
- 10. On page 7, the "Disposal of Infectious Material" statement under the "HIV" directions must be brought into compliance with PR Notice 2005, Mandatory Labeling, by deleting the word, should, and replace with "must".
- 11. On page 7, the second statement under the "Use as a Deodorizer" directions must be revised to indicate what surfaces are to be deodorized by stating "Wipe surfaces that are listed on the label (and let air dry)."
- 12. On page 8, revise the, NONREFILLABLE SEALED CONTAINERS instructions by deleting "OUTPOST, Single Solutions, Smart Dose, Command Center" because the directions never references these types of containers.

Also, revise the SMALL NONREFILLABLE CONTAINERS: directions by deleting the following statements, "Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer container for recycling, if available."

- 13. On page 9, delete the direction for LARGE NONREFILLABLE CONTAINERS section because the language is not applicable to antimicrobial products. The rinsate cannot be placed in application equipment because the use dilution is critical in mitigating the organisms cited on the label.
- 14. The containers described in the refillable container of one gallon or less do not meet the definition of refillable containers as per definitions #9 and #10 at <http://www.epa.gov/pesticides/regulation/containerquestions.htm> and should be deleted.

In accordance with PR Notice 2007-4 the Agency defines refillable as: "Refillable container" means a container that is intended to be filled with pesticide more than once for sale or distribution. The registrant determines whether a container is intended to be refilled for sale or distribution (in which case it is a refillable container) or if it is not intended to be refilled for sale or

distribution (in which case it is a non refillable container) and implements its decision by labeling the container appropriately. This decision also would impact packaging design, packaging material, and distribution and marketing.


- 15. On page 9, The "PRECAUTIONARY STATEMENTS" must be brought into agreement with the Label Manual 3rd edition by revising the fifth statement to read "was thoroughly with soap and water, after handling and before eating drinking, chewing gum, using tobacco or using the toilet.

General Comments:

A stamped copy of the acceptable labeling is enclosed. Submit one (1) copy of your final printed labeling before distributing or selling the product bearing the revised labeling.

Should you have any questions or comments concerning this letter, please contact Drusilla Copeland at (703) 308-6224.

Sincerely,



Velma Noble
Product Manager (31)
Regulatory Management Branch I
Antimicrobials Division (7510P)

Enclosure: Stamped label

4/15

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

JohnsonDiversey



JOHNSON'S FORWARD® CLEANER (GENERAL PURPOSE) DISINFECTANT CLEANER

Bactericidal • Fungicidal • Deodorizing (Odor Counteractant) • *Virucidal

Disinfects in up to 500 ppm hard water and 10% serum load • Non-Acid Bowl (Cleaner) (Disinfectant) • Non-Acid Bathroom (Cleaner) (Disinfectant) • Non-Acid Shower Room (Cleaner) (Degreaser) (Disinfectant) • (Cleans the dirt and leaves the shine.) • (Cleans (and Deodorizes) in One Step) • (Leaves Toilet Bowls Sparkling Clean) • (Leaves the Bathroom Sparkling Clean) • (Cleans without Harmful Acid Fumes) • (Disinfects as it Cleans) • (Contains no abrasives (Non-Abrasive))

(Fragrance Free Formula), (Unscented),
((Fresh) [Lemon] [Pine] [Mint] [Spring Fresh] [Floral] [Citrus] [Powder] (Scent) (Fragrance))

For (Hospital,) (Foodservice,) (Commercial,) (Industrial) & (Institutional) Use (Only)

ACTIVE INGREDIENT:	
n-Alkyl (50% C ₁₄ , 40% C ₁₂ , 10% C ₁₆) dimethyl benzyl ammonium chloride	1.60%
INERT (OTHER) INGREDIENTS:	98.40%
TOTAL:	100.00%

**KEEP OUT OF REACH OF CHILDREN.
DANGER**

See additional precautionary statements on (back) (side) (left) (right) (panel) of label (below).

See reference sheet (enclosed in each case) for (a complete list of pathogenic organisms) (additional features, claims, direction for use) (eliminated by) (claimed by) (JOHNSON'S FORWARD® CLEANER) (for this product).

Net Contents:

(Product of USA)

ACCEPTED
with COMMENTS
in EPA Letter Dated:

MAR 3 2009

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 70627-10

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

(MARKETING CLAIMS): (Note to EPA - This text will only appear on the appropriate container.)

ACCUMIX™ CONTAINERS DISPENSING SYSTEM (32 oz. container) - ACCUMIX™. (Remove [Loosen] cap). Squeeze. (Squeeze bottle.) Measure. (Measure amount). Pour. (Pour contents). (Designed) For use with 5 gallon Buddy Jugs™ (the BIG BUDDY™ [ACCUTAINER™] System). This product can also be diluted into pre-cleaned and properly labeled 5-gallon BUDDY JUG™ (ACCUTAINERS™) (Dispensing Containers) for dispensing as needed.

SOLUTIONS CENTER™ DISPENSING SYSTEM (64 oz. containers) - For use with (JOHNSON WAX PROFESSIONAL™) (JohnsonDiversey) (SOLUTIONS CENTER™) Brand Dispensing Equipment. Eliminates Mixing. The Accurate Solution To Cleaner Dilution. Disinfectant Cleaning With The Convenience Of (SOLUTIONS CENTER™). ((SOLUTIONS CENTER™) (This packaging offers) Reduced Exposure To Concentrate Due To Closed Transfer System. Solutions To Go!

J-FILL™ DISPENSING SYSTEM (84.5 oz. containers) - For use with (JOHNSON WAX PROFESSIONAL™) (JohnsonDiversey) (J-FILL™) Brand Dispensing Equipment. Eliminates Mixing. The Accurate Solution To Cleaner Dilution. Disinfectant Cleaning With The Convenience Of J-FILL™. (J-FILL™ (This) packaging offers) Reduced (Reduces) Exposure To Concentrate Due To Closed Transfer System. Solutions To Go!

RTD™ DISPENSING SYSTEM (1.5L & 5L containers) - RTD. Very Simple. Very Smart. Ready-To-Dispense. Provides accuracy. No chemical connections, sealed bottle. Proven dilution control concentrate.

PRE-MEASURED PACKETS DISPENSING SYSTEM - Pre-Measured [Tear-Open] Packets.

(General Uses)

JOHNSON'S FORWARD® CLEANER cleans quickly by removing dirt, grime, body oils and other common soils. This product is for use in hospitals, nursing homes, schools and colleges, (day care centers), (medical) offices, funeral homes, veterinary clinics, pet shops, (equine farms), animal life science laboratories, hotels, motels, restrooms, food service establishments (restaurants).

It is designed for use on (Use daily on) the following hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, stainless steel, aluminum, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilets, urinals, sinks, shower rooms and locker rooms areas - (food and non-food contact) surface where disinfection is required. A potable water rinse is required for food contact surfaces. Do not use on glasses, dishes, and utensils.

(Hospitals/Health Care Facilities)

JOHNSON'S FORWARD® CLEANER is a one-step (hospital-use) germicidal (disinfectant) cleaner and deodorant (odor-counteractant) (odor neutralizer) designed for general cleaning, (and) disinfecting, (deodorizing) (of) hard, non-porous environmental surfaces. JOHNSON'S FORWARD® CLEANER cleans quickly by removing dirt, grime, food residue, body oils, dead skin, blood and other organic matter commonly found in hospitals (in health care facilities) (on medical surfaces). It (also) eliminates odors leaving (restroom) surfaces smelling clean and fresh. Use where odors are a problem.

JOHNSON'S FORWARD® CLEANER cleans, disinfects and deodorizes (hard, non-porous environmental hospital (medical) surfaces) in one step (with no rinsing required). Its non-abrasive formula is designed for use on (Use daily on) the following hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, stainless steel, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilets, urinals, sinks found in (health care facilities [hospitals],) patient rooms, operating rooms, ICU areas, shower rooms, and locker rooms. (It can also be used to pre-clean and disinfect hospital items: wheelchairs, [hospital] [patient] bed rails and linings, wash basins, bedpans, [and] - any washable (food and non-food contact) surface (where disinfection is required). A potable water rinse is required when disinfecting food contact surfaces. Do not use on glasses, dishes, and utensils.

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

(Food Service:)

JOHNSON'S FORWARD® CLEANER removes dirt, grime and food soils in food preparation and processing areas. Its non-abrasive formula will not harm (scratch) surfaces. It cleans, disinfects and eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

Its non-abrasive formula is designed for use on (Use daily on) the following hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, aluminum, stainless steel, brass, copper, laminated surfaces and baked enamel surfaces (associated with floors, walls, ceilings, tables, chairs, countertops, fixtures, glazed tile, toilets, (toilet bowls), urinals, sinks found in food establishments, (restaurants), (commercial kitchens) & restrooms. A potable water rinse is required for food contact surfaces. Do not use on glasses, dishes, and utensils.

(Animal Housing Facilities:)

JOHNSON'S FORWARD® CLEANER cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine or poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It (also) eliminates odors leaving surfaces smelling clean and fresh.

JOHNSON'S FORWARD® CLEANER cleans, disinfects and deodorizes (hard, non-porous environmental surfaces) in one step. Its non-abrasive formula is designed for use on (Use daily on) (Use daily to clean and disinfect) hard, non-porous surfaces: plated or stainless steel, aluminum, chrome, glazed porcelain, glazed tile, laminated surfaces (associated with floors, walls, countertops, cages, kennels, animal equipment) found in (barns, pens and stalls) animal housing facilities.

(Public Restrooms:)

JOHNSON'S FORWARD® CLEANER is a one-step disinfectant cleaner and deodorant (odor-counteractant) (odor-neutralizer) designed for general cleaning, (and) (disinfecting) (deodorizing) on hard, non-porous environmental surfaces.

JOHNSON'S FORWARD® CLEANER cleans, disinfects and deodorizes surfaces by killing odor-causing germs. Its non-abrasive formula is designed for use on (restroom surfaces): glazed ceramic (restroom) tile, glazed porcelain, chrome, stainless steel and plastic surfaces associated with floors, walls, fixtures, toilets, urinals, sinks, shower rooms and locker rooms.

(Non-Acid Bowl [& Bathroom] Disinfectant Cleaner:)

JOHNSON'S FORWARD® CLEANER is a (concentrated) non-acid (bowl and) bathroom cleaner which cleans, disinfects and deodorizes in one easy step. It cleans, disinfects and deodorizes toilet bowls, urinals, rims, sinks, sink basins, faucets, tubs, glazed tiles, glazed ceramic, glazed porcelain, chrome, stainless steel, and all hard non-porous, washable surfaces found in the bathroom (restroom) (in the presence of organic soil).

JOHNSON'S FORWARD® CLEANER eliminates odors leaving bathrooms (restrooms) smelling clean and fresh. Use where odors are a problem. It cleans, disinfects and deodorizes surfaces by killing odor-causing microorganisms. Its non-abrasive formula is designed for use (Use it daily) on ([hard, non-porous environmental] restroom surfaces): glazed ceramic (restroom) tile, glazed porcelain, chrome, stainless steel and plastic surfaces associated with floors, walls, fixtures, toilets, urinals, sinks, shower rooms and locker rooms.

(Refill)

(For Refilling Concentrate From Large Containers Into Smaller Containers:)

JOHNSON'S FORWARD® CLEANER may be used to fill and refill clean, properly labeled containers for dilution elsewhere within your facility. Make sure the small container has been cleaned, dried and properly labeled. Also make sure other items (such as funnels or hand pumps) are properly cleaned and dried. To refill, simply pour (or pump) product from the larger container directly into the smaller one being careful not to spill any product. Keep both containers sealed when not in use.

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

(CLAIMS:)

When used as directed at a 1:32 dilution (4 oz. per gallon of water), JOHNSON'S FORWARD® CLEANER is highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms (including bacteria, antibiotic resistant bacteria, viruses and fungi). (See reference sheet (enclosed in each case) for a complete list of organisms.) Using approved AOAC test methods (under Good Laboratory Practices, [GLP's]), in the presence of 500 ppm hard water, 10% serum load and 10 minutes contact time JOHNSON'S FORWARD® CLEANER kills the following on hard non-porous inanimate surfaces:

Bacteria (Bactericidal Activity)

- Pseudomonas aeruginosa*, (ATCC 15442)
- Staphylococcus aureus*, (ATCC 6538)
- Salmonella choleraesuis*, (ATCC 10708)
- Bordetella bronchiseptica*, (ATCC 10580)
- Burkholderia cepacia*, (ATCC 25416),
formerly known as *Pseudomonas cepacia*
- Campylobacter fetus*, (ATCC 27374)
- Chlamydia psittaci*, (VR-125)
- Citrobacter freundii*, (ATCC 8090)
- Enterobacter aerogenes*, (ATCC 13048)
- Enterobacter agglomerans*, (ATCC 27155)
- Enterobacter cloacae*, (ATCC 23355)
- Enterobacter gergoviae*, (ATCC 33028)
- Enterobacter liquefaciens*, (ATCC 14460)
- Enterococcus faecalis*, (ATCC 19433)
- Enterococcus hirae*, (ATCC 10541)
- Escherichia coli (E. coli)*, (ATCC 11229)
- Escherichia coli (E. coli) 0157:H7*, (ATCC 43890)
- Flavobacterium meningosepticum*, (ATCC 13253)
- Hafnia alvei*, (ATCC 13337)

- Klebsiella oxytoca*, (ATCC 13182)
- Klebsiella pneumoniae*, (ATCC 13883)
- Legionella pneumophila*, (ATCC 33153)
- Listeria monocytogenes*, (ATCC 15313)
- Micrococcus luteus*, (ATCC 4698)
- Micrococcus luteus*, (ATCC 14452)
- Micrococcus sedentarius*, (ATCC 27573)
- Morganella morganii*, (ATCC 25830)
- Neisseria gonorrhoeae*, (ATCC 43069)
- Pasteurella multocida*, (ATCC 43137)
- Proteus mirabilis*, (ATCC 9240)
- Proteus vulgaris*, (ATCC 13315)
- Pseudomonas diminuta*, (ATCC 11568)
- Pseudomonas fluorescens*, (ATCC 13525)
- Pseudomonas putida*, (ATCC 12633)
- Pseudomonas stutzeri*, (ATCC 17588)
- Salmonella enteritidis*, (ATCC 13076)
- Salmonella gallinarum*, (ATCC 9184)
- Salmonella schottmuelleri*, (ATCC 10719)
- Salmonella typhi*, (ATCC 6539)

- Salmonella typhimurium*, (ATCC 13311)
- Serratia grimesii*, (ATCC 14460)
- Serratia marcescens*, (ATCC 9103)
- Shigella dysenteriae*, (ATCC 29026)
- Shigella flexneri*, (ATCC 25875)
- Shigella sonnei*, (ATCC 25931)
- Staphylococcus aureus*, (ATCC 25923)
- Staphylococcus aureus* (Toxic Shock),
(ATCC 33586)
- Staphylococcus epidermidis*, (ATCC 14990)
- Staphylococcus haemolyticus*, (ATCC 29970)
- Staphylococcus saprophyticus*, (ATCC 15305)
- Staphylococcus species*, (ATCC 12715)
- Streptococcus agalactiae*, (ATCC 13813)
- Streptococcus mutans*, (ATCC 25175)
- Streptococcus pyogenes*, (ATCC 19615)
- Streptococcus pyogenes* ("Strep A" -
Flesh Eating Strain), (clinical isolate)
- Vibrio cholera*, (ATCC 11623)
- Yersinia enterocolitica*, (ATCC 9610)

Antibiotic-Resistant (Strains of) Bacteria (Antibiotic-Resistant Bactericidal Activity)

- E. coli*, (ATCC 55244);
(resistant to Kanamycin)
- E. coli*, (ATCC 47041);
(resistant to Tetracycline)
- Enterococcus faecalis*, (ATCC 51299);
(resistant to Vancomycin [VRE])

- Klebsiella oxytoca*, (ATCC 15764);
(resistant to Ampicillin,
Dihydrostreptomycin)
- Micrococcus sedentarius*, (ATCC 27573)
(resistant to Methicillin)
- Staphylococcus aureus*, (ATCC 14154);
(resistant to Erythromycin, Penicillin,
Streptomycin, Tetracycline)

- Staphylococcus aureus*, (ATCC 33592);
(resistant to Methicillin [MRSA],
Gentamicin [GRSA])
- Streptococcus pneumoniae*, (ATCC 51915)
resistant to Penicillin [PRSP]

***Viruses (*Virucidal Activity)**

- **Cytomegalovirus*, (VR-538)
- **Herpes simplex Type 1*, (VR-733)

- **Herpes simplex Type 2*, (VR-734)
- **Human Coronavirus*, (VR-740)
- **Influenza Type A₂ (Hong Kong)*, (VR-544)

- **Parainfluenza Type 3*, (VR-93)
- **Respiratory syncytial virus*, (VR-26)
- **Vaccinia virus (Smallpox Vaccine Virus)*,
(VR-119)

Kills HIV-1 (AIDS virus)(HTLV-III_B) when used as directed on hard non-porous inanimate surfaces with a 1 minute contact time.

(*Veterinary viruses:)

- **Avian Infectious bronchitis (IBV)*, (VR-22)

- **Avian Influenza*, (VR-2072)

- **Transmissible gastroenteritis virus (TGE)*,
(U of Minn. Strain)

Fungi/Yeast (Fungicidal and Yeast Activity)

- Geotrichum candidum*, (ATCC 18301)

- Trichophyton mentagrophytes* (athlete's foot
fungus), (ATCC 9533)

- Zygosaccharomyces bailii*, (ATCC 56075)

Bactericidal Stability of Use-Solution - Tests show JOHNSON'S FORWARD® CLEANER, when diluted in 400 ppm hard water in the presence of 10% serum, remains effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* for up to 30 days in storage as long as it remains sealed. If the product becomes visibly dirty or contaminated, the use-solution must be discarded and fresh product prepared. Always use clean, dry containers when diluting this product.

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

When used as directed at a 1:43 dilution (3 oz. per gallon of water) in the presence of 200 ppm hard water, 10% serum load, JOHNSON'S FORWARD® CLEANER, kills HIV-1 (AIDS virus)(HTLV-III_B) in 1 minute on hard non-porous inanimate surfaces.

When used as directed at a 1:43 dilution (3 oz. per gallon of water) in the presence of 200 ppm hard water, 10% serum load, JOHNSON'S FORWARD® CLEANER, kills *Trichophyton mentagrophytes* (athlete's foot fungus)(ATCC 9533) in 10 minutes when applied to areas as: shower and locker room floors, benches, and floor mats and other hard, non-porous environmental surfaces where *Trichophyton mentagrophytes*, (athlete's foot fungus) may be found.

Malodor(s) (Activity) (Counteractancy) – eliminates (destroys) odors and odor-causing bacteria in restroom areas; behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors).

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

(This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.)

JOHNSON'S FORWARD® CLEANER can be applied by mop, sponge, cloth, paper towel, (hand pump) trigger sprayer, auto-scrubber or foam gun (device). Change cloth, sponges or towels frequently to avoid redeposition of soil. For disinfection, all surfaces must remain wet for 10 minutes.

For 1:32 Dilution

To Prepare Use Solution: Add the product at 4 oz. per gallon of water (1:32).

(Note to EPA: The directions that appear for specific container sizes will be substituted for the sentence above only on that container size.)

ACCUMIX™ CONTAINERS DISPENSING SYSTEM (32 oz. container) - Add the product at 4 oz. per gallon of water (1:32).

SOLUTIONS CENTER™ DISPENSING SYSTEM (64 oz. containers) - Remove cap and insert cartridge into dispenser. Note: See dispenser instructions for proper cartridge placement. Once cartridge is in place, (squeeze the handle) (or) (press the button) to dispense a 1:32 solution into a bucket, bottle, scrubber or other container.

J-FILL™ DISPENSING SYSTEM (84.5 oz. containers) - Remove cap and insert cartridge into dispenser. Note: See dispenser instructions for proper cartridge placement. Once cartridge is in place, (squeeze the handle) (or) (press the button) to dispense a 1:32 solution into a bucket, bottle, scrubber or other container.

RTD™ DISPENSING SYSTEM (1.5L and/or 5L containers) -Turn off water to connect unit. Attach to water source. Rotate control knob to fill bottle or bucket. Squeeze handle to dispense a 1:32 solution into a bottle (bucket) (or other container). See device instruction manual for more information.

PRE-MEASURED PACKETS DISPENSING SYSTEM - (Simply) open and pour xx fl. oz. of product (contents) into xx fl. oz. of water (for a 1:32 dilution). (Pour contents of packet into xx fl. oz. (gallons) of water for a 1:32 dilution.) Keep packets in box until ready to use.

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

For Use as a (One-Step) Cleaner/Disinfectant:

1. Pre-clean heavily soiled areas.
2. Apply Use Solution to hard, non-porous environmental surfaces.
3. All surfaces must remain wet for ten (10) minutes.
4. Wipe surfaces (and let air dry).

Note - Rinsing is not necessary unless floors are to be coated with finish or restorer. All food contact surfaces such as appliances and kitchen countertops must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

For food processing plants/federally inspected meat and poultry establishments: Before using this product, food products and packaging materials must be removed from the room or carefully protected. After use, all surfaces in the area must be thoroughly rinsed with potable water.

To Kill Fungi:

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for ten (10) minutes. Wipe surfaces (and let air dry).

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls from Concentrate:

1. Pre-clean heavily soiled areas.
2. Pour 4 oz. of product (1 packet) into toilet bowl.
3. Swab entire surface area especially under the rim.
4. Allow entire surface to remain wet for ten (10) minutes.
5. Flush toilet and rinse swab applicator thoroughly.

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls [and Urinals] from Use-Dilution:

1. Pre-clean heavily soiled areas.
2. Empty toilet bowls by forcing water through the trap. Apply Use Solution to exposed surfaces in toilet bowls and urinals.
3. Swab entire surface area especially under the rim.
4. Allow entire surface to remain wet for ten (10) minutes.
5. Flush toilet or urinal and rinse swab applicator thoroughly.

For Use To Clean and Disinfect Shower Rooms, Locker Rooms and Other Large, Open Areas with Floor Drains:

1. Pre-clean heavily soiled areas.
2. Apply Use Solution to floors, walls and ceilings making sure not to over spray. To disinfect, all surfaces must remain wet for ten (10) minutes.
Special instructions for foam guns: Pour concentrate into foam gun bottle and attach bottle to spray nozzle and ensure gun is attached to hose. Note: See foam gun instructions for more information. Make sure setting is set for a 1:32 dilution. Once in place, squeeze the handle to dispense foam solution. To disinfect, all surfaces must remain wet for ten (10) minutes.
3. Scrub using a deck brush or other coarse material as necessary.
4. Rinse surfaces thoroughly and let air dry.

For Use In Treatment of Animal Housing Facilities:

1. Remove all animals and feed from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and rinse with water.
5. Apply fresh use solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for ten (10) minutes.
6. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until product has dried.
7. For feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand ten (10) minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

10/15

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

For 1:43 Dilution

To Prepare Use Solution: Add the product at 3 oz. per gallon of water (1:43).

(Note to EPA: The directions that appear for specific container sizes will be substituted for the sentence above only on that container size.)

ACCUMIX™ CONTAINERS DISPENSING SYSTEM (32 oz. container) - Add the product at 3 oz. per gallon of water (1:43).

SOLUTIONS CENTER™ DISPENSING SYSTEM (64 oz. containers) - Remove cap and insert cartridge into dispenser. Note: See dispenser instructions for proper cartridge placement. Once cartridge is in place, (squeeze the handle) (or) (press the button) to dispense a 1:43 solution into a bucket, bottle, scrubber or other container.

PRE-MEASURED PACKETS DISPENSING SYSTEM (84.5 oz. containers) - Remove cap and insert cartridge into dispenser. Note: See dispenser instructions for proper cartridge placement. Once cartridge is in place, (squeeze the handle) (or) (press the button) to dispense a 1:43 solution into a bucket, bottle, scrubber or other container.

To Kill Fungi:

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for ten (10) minutes. Wipe surfaces (and let air dry).

***(JOHNSON'S FORWARD® CLEANER) Kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of JOHNSON'S FORWARD® CLEANER.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1 and for 10 minutes to kill all other organisms cited on the label.

Disposal of Infectious Material: Blood and other body fluids should be autoclaved and disposed of according to Federal, State, and local regulations for infectious waste disposal.

Light Duty Cleaning: For cleaning of most soils on most surfaces: Use a 1:128 dilution (1 oz. per gallon of water). Apply to surfaces. Wipe surfaces (and let air dry).

Normal Cleaning: Use a 1:43 dilution (3 oz. per gallon of water). Apply to surfaces. Wipe surfaces (and let air dry).

Heavy Duty Cleaning: Use up to a 1:32 dilution (4 oz. per gallon of water). Apply to surfaces. Wipe surfaces (and let air dry).

For Use as a Deodorizer: Apply Use Solution to surfaces. Wipe surfaces (and let air dry).

11/15

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE:

Do not reuse empty container (unless refilling from a larger container of the same product according to the refilling directions outlined previously.) (Keep from freezing.)

PESTICIDE DISPOSAL:

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL Note to reviewer – One or more of the following paragraphs for Container Disposal will be selected, depending on packaging type:

(If not being refilled with this product,) triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NONREFILLABLE SEALED CONTAINERS: Note to EPA: Several of our packaging options (including ACCUMIX Containers™, Solutions Center™, J-Fill™, OUTPOST™, Single Solutions™, Smart Dose, Command Center™, or RTD™) are sealed containers or bottles designed to reduce worker exposure to the concentrate. None of these can be triple rinsed because they are closed, sonically welded, sealed containers. The following text will be used on these sealed container types:

Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash.

NONREFILLABLE NON-RIGID CONTAINERS: Note to EPA: Several of our packaging options are Bag-in-Box containers (a plastic bag liner supported inside a box) or are flexible bag-type containers (such as ACCUPACK™ Containers). These flexible containers are exempt from the triple rinsing requirements. The following text will be used on these container types:

Nonrefillable container. Do not reuse or refill this container. Wrap empty container and put in trash.

SMALL NONREFILLABLE CONTAINERS: Note to EPA: The following text will be used on rigid, nonrefillable containers small enough to shake (5 gallons or smaller):

Nonrefillable container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer container for recycling, if available.

12/15

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

LARGE NONREFILLABLE CONTAINERS: Note to EPA: One of the following paragraphs will be used on labels for rigid, nonrefillable containers too large to shake (larger than 5 gallons):

Nonrefillable container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for at least 30 seconds. Stand the container on its end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Offer container for reconditioning, if appropriate.

OR

Nonrefillable container. Clean container promptly after emptying. Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure-rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip. Offer for reconditioning, if appropriate.

REFILLABLE CONTAINERS: Note to EPA: One of the following paragraphs will be used on labels for refillable containers:

Refillable container. Refill this container with (this brand or brand name) pesticide only. Do not reuse this container for any other purpose.

Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

OR

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container prior to final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

ENVIRONMENTAL HAZARDS (for containers of 5 gallons or more)

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or clothing. Wear chemical splash-proof goggles or face shield, chemical resistant gloves and protective clothing. Harmful if absorbed through skin or swallowed. Wash thoroughly with soap and water after handling. Remove and wash contaminated clothing before reuse.

13/15

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE.

1-800-851-7145

Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage.

EPA Reg. No. 70627-10

EPA Est. No.

(Lot code letters indicate establishment number.)

MSDS No. XXXXXXX

© 2008 JohnsonDiversey, Inc., 8310 16th Street, Sturtevant, WI 53177, U.S.A. All Rights Reserved.

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

JohnsonDiversey



JOHNSON'S FORWARD® CLEANER REFERENCE SHEET

When used as directed at a 1:32 dilution (4 oz. per gallon of water), JOHNSON'S FORWARD® CLEANER is highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms. (See reference sheet (enclosed in each case) for a complete list of organisms.) Using approved AOAC test methods (under Good Laboratory Practices, [GLP's]), in the presence of 500 ppm hard water, 10% serum load and 10 minutes contact time JOHNSON'S FORWARD® CLEANER kills the following on hard non-porous inanimate surfaces:

Bacteria (Bactericidal Activity)

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella choleraesuis, (ATCC 10708)
Bordetella bronchiseptica, (ATCC 10580)
Burkholderia cepacia, (ATCC 25416),
 formerly known as *Pseudomonas cepacia*
Campylobacter fetus, (ATCC 27374)
Chlamydia psittaci, (VR-125)
Citrobacter freundii, (ATCC 8090)
Enterobacter aerogenes, (ATCC 13048)
Enterobacter agglomerans, (ATCC 27155)
Enterobacter cloacae, (ATCC 23355)
Enterobacter gergoviae, (ATCC 33028)
Enterobacter liquefaciens, (ATCC 14460)
Enterococcus faecalis, (ATCC 19433)
Enterococcus hirae, (ATCC 10541)
Escherichia coli (*E. coli*), (ATCC 11229)
Escherichia coli (*E. coli*) 0157:H7, (ATCC 43890)
Flavobacterium meningosepticum, (ATCC 13253)
Hafnia alvei, (ATCC 13337)

Klebsiella oxytoca, (ATCC 13182)
Klebsiella pneumoniae, (ATCC 13883)
Legionella pneumophila, (ATCC 33153)
Listeria monocytogenes, (ATCC 15313)
Micrococcus luteus, (ATCC 4698)
Micrococcus luteus, (ATCC 14452)
Micrococcus sedentarius, (ATCC 27573)
Morganella morganii, (ATCC 25830)
Neisseria gonorrhoeae, (ATCC 43069)
Pasteurella multocida, (ATCC 43137)
Proteus mirabilis, (ATCC 9240)
Proteus vulgaris, (ATCC 13315)
Pseudomonas diminuta, (ATCC 11568)
Pseudomonas fluorescens, (ATCC 13525)
Pseudomonas putida, (ATCC 12633)
Pseudomonas stutzeri, (ATCC 17588)
Salmonella enteritidis, (ATCC 13076)
Salmonella gallinarum, (ATCC 9184)
Salmonella schottmuelleri, (ATCC 10719)
Salmonella typhi, (ATCC 6539)

Salmonella typhimurium, (ATCC 13311)
Serratia grimesii, (ATCC 14460)
Serratia marcescens, (ATCC 9103)
Shigella dysenteriae, (ATCC 29026)
Shigella flexneri, (ATCC 25875)
Shigella sonnei, (ATCC 25931)
Staphylococcus aureus, (ATCC 25923)
Staphylococcus aureus (Toxic Shock),
 (ATCC 33586)
Staphylococcus epidermidis, (ATCC 14990)
Staphylococcus haemolyticus, (ATCC 29970)
Staphylococcus saprophyticus, (ATCC 15305)
Staphylococcus species, (ATCC 12715)
Streptococcus agalactiae, (ATCC 13813)
Streptococcus mutans, (ATCC 25175)
Streptococcus pyogenes, (ATCC 19615)
Streptococcus pyogenes ("Strep A" -
 Flesh Eating Strain), (clinical isolate)
Vibrio cholera, (ATCC 11623)
Yersinia enterocolitica, (ATCC 9610)

Antibiotic-Resistant (Strains of) Bacteria (Antibiotic-Resistant Bactericidal Activity)

E. coli, (ATCC 55244);
 (resistant to Kanamycin)

E. coli, (ATCC 47041);
 (resistant to Tetracycline)

Enterococcus faecalis, (ATCC 51299);
 (resistant to Vancomycin [VRE])

Klebsiella oxytoca, (ATCC 15764);
 (resistant to Ampicillin,
 Dihydrostreptomycin)

Micrococcus sedentarius, (ATCC 27573)
 (resistant to Methicillin)

Staphylococcus aureus, (ATCC 14154);
 (resistant to Erythromycin, Penicillin,
 Streptomycin, Tetracycline)

Staphylococcus aureus, (ATCC 33592);
 (resistant to Methicillin [MRSA],
 Gentamicin [GRSA])

Streptococcus pneumoniae, (ATCC 51915)
 resistant to Penicillin [PRSP]

*Viruses (*Virucidal Activity)

*Cytomegalovirus, (VR-538)
 *Herpes simplex Type 1, (VR-733)

*Herpes simplex Type 2, (VR-734)
 *Human Coronavirus, (VR-740)
 *Influenza Type A₂ (Hong Kong), (VR-544)

*Parainfluenza Type 3, (VR-93)
 *Respiratory syncytial virus, (VR-26)
 *Vaccinia virus (Smallpox Vaccine Virus),
 (VR-119)

Kills HIV-1 (AIDS virus)/(HTLV-III_B) when used as directed on hard non-porous inanimate surfaces with a 1 minute contact time.

(*Veterinary viruses:)

*Avian Infectious bronchitis (IBV), (VR-22)

*Avian Influenza, (VR-2072)

*Transmissible gastroenteritis virus (TGE),
 (U of Minn. Strain)

Fungi/Yeast (Fungicidal and Yeast Activity)

Geotrichum candidum, (ATCC 18301)

Trichophyton mentagrophytes (athlete's foot
 fungus), (ATCC 9533)

Zygosaccharomyces bailii, (ATCC 56075)

15/15

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

Bactericidal Stability of Use-Solution - Tests show JOHNSON'S FORWARD® CLEANER, when diluted in 400 ppm hard water in the presence of 10% serum, remains effective against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella choleraesuis* for up to 30 days in storage as long as it remains sealed. If the product becomes visibly dirty or contaminated, the use-solution must be discarded and fresh product prepared. Always use clean, dry containers when diluting this product.

When used as directed at a 1:43 dilution (3 oz. per gallon of water) in the presence of 200 ppm hard water, 10% serum load, JOHNSON'S FORWARD® CLEANER, kills HIV-1 (AIDS virus)(HTLV-III_B) in 1 minute on hard non-porous inanimate surfaces.

When used as directed at a 1:43 dilution (3 oz. per gallon of water) in the presence of 200 ppm hard water, 10% serum load, JOHNSON'S FORWARD® CLEANER, kills *Trichophyton mentagrophytes* (athlete's foot fungus)(ATCC 9533) in 10 minutes when applied to areas as: shower and locker room floors, benches, and floor mats and other hard, non-porous environmental surfaces where *Trichophyton mentagrophytes*, (athlete's foot fungus) may be found.

Malodor(s) (Activity) (Counteractancy) – eliminates (destroys) odors and odor-causing bacteria in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors).

EPA Reg. No.70627-10

© 2008 JohnsonDiversey, Inc., 8310 16th Street, Sturtevant, WI 53177, U.S.A. All Rights Reserved.