

70627-21

04/02/2001

1/13

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

SC Johnson Commercial Markets, Inc.
8310 16th Street, MS 675
Sturtevant, WI 53177-0902

APR 02 2001

Subject: Virex II/128
EPA Registration No. 70627-21
Amendment Date: 12/18/00
EPA Receive Date: 1/03/01


Attention: Ms. Lisa Amadio

The amendment, referenced above, submitted in connection with registration under FIFRA, as amended, for conditional use of historic inhalation statements as well as updating the label per Agency letter dated March 6, 2000 is acceptable provided you address the following comments before you release this product for shipment.

- 1) The Agency will allow the use of the historic inhalation statements that were accepted on September 30, 1996 stamped label until a decision has been made regarding the submitted inhalation exposure and risk assessment data on 70627-24's formulation.
- 2) The First Aid Statements will need to be updated in accordance with PR Notice 2001-1.

A stamped label is enclosed for your records. Please submit (1) copy of the final printed label before you release this product for shipment. If you have any questions regarding this letter, please contact Jacqueline McFarlane at (703) 308-6416.

Sincerely,



Velma Noble
Product Manager (31)
Regulatory Management Branch I
Antimicrobials Division (7510C)

CONCURRENCES

SYMBOL	720C						
SURNAME	McFarlane						
DATE	3/28/01						

Johnson Wax PROFESSIONAL

VIREX™ II/ 128 One-Step Disinfectant Cleaner And Deodorant

Bactericidal • Virucidal • Fungicidal • Mildewcidal • (Mildewstatic) • Deodorizer (Odor Counteractant) (Odor Neutralizer)
• Non-Dulling To Floors (Floor Finishes)

(Fragrance Free Formula), (Unscented), (Fresh [Lemon], [Pine], [Mint], [Spring Fresh], [Floral], [Citrus], [Powder] Scent)

For (Hospital,) (Foodservice,) (Commercial,) Industrial & Institutional Use (Only)
(Suitable) For Use in Meat and Poultry Plants

ACTIVE INGREDIENTS:	
Didecyl dimethyl ammonium chloride.....	4.352%
n-Alkyl (50% C ₁₄ , 40% C ₁₂ , 10% C ₁₆) dimethyl benzyl ammonium chloride.....	4.095%
INERT (OTHER) INGREDIENTS:	91.553%
TOTAL:	100.000%

**KEEP OUT OF REACH OF CHILDREN
DANGER:**

See additional precautionary statements on back (side) (left) (right) (panel) (of) (label) (below).

See reference sheet (enclosed in each case) for (a complete list of pathogenic organisms) (additional features, claims, directions for use) (claimed for this product) (eliminated by VIREX™ II/ 128).

Net Contents:

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 70627-21

(Product of USA)

(MARKETING CLAIMS): (Note to EPA - This text will only appear on the appropriate container.)

ACCUMIX™ CONTAINERS (32 oz. container) - ACCUMIX™. (Remove [Loosen] cap). Squeeze. (Squeeze bottle.) Measure. (Measure amount). Pour. (Pour contents). (Designed) For use with 5 gallon Buddy Jugs™ (the BIG BUDDY™ [ACCUTAINER™] System). VIREX™ II/ 128 can also be diluted into pre-cleaned and properly labeled 5-gallon BUDDY JUG™ (ACCUTAINERS™) (Dispensing Containers) for dispensing as needed.

SOLUTIONS CENTER™ (64 oz. containers) - For use with (JOHNSON™) (PRISM™) (DRACKET™) (SOLUTIONS CENTER™) Brand Dispensing Equipment. Eliminates Mixing. The Accurate Solution To Cleaner Dilution. Disinfectant Cleaning With The Convenience Of (SOLUTIONS CENTER™). (SOLUTIONS CENTER™) (This) packaging offers) Reduced Exposure To Concentrate Due To Closed Transfer System. Solutions To Go!

J-FILL™ DISPENSING SYSTEMS (84.5 oz. containers) - For use with (JOHNSON™) (PRISM™) (DRACKET™) (J-FILL™) Brand Dispensing Equipment. Eliminates Mixing. The Accurate Solution To Cleaner Dilution. Disinfectant Cleaning With The Convenience Of J-FILL™. J-FILL™ (This) packaging offers) Reduced Exposure To Concentrate Due To Closed Transfer System. Solutions To Go!

ACCUPACK™ CONTAINERS (Pre-measured packets) - Pre-Measured [Tear-Open] Packets.

J-SPRAY™ DISPENSING SYSTEM - For use with (JOHNSON™) (PRISM™) (DRACKET™) (J-Spray™) (EASY PAKS 4-SHOT™) Brand Dispensing Equipment. Eliminates Mixing. The Accurate Solution To Cleaner Dilution. Disinfectant Cleaning With The Convenience Of (J-Spray™) (EASY PAKS 4-SHOT™). (J-Spray™) (EASY PAKS 4-SHOT™) (This) packaging offers) Reduced Exposure To Concentrate Due To Closed Transfer System. Solutions To Go! Concentrate. Makes xx Bottles. Each compartment of a J-Spray™ cartridge produces xx oz. (xx mL) ready-to-use product. Use only with the J-Spray™ Virex™ II/ 128 Use Solution Trigger Bottle. Read instructions before use. Four concentrate cartridges makes 16 refills for at total xx gallons (xx L) of ready-to-use Virex™ II/ 128.

(FEATURES, CLAIMS & USES:)

(General Uses)

VIREX™ II/ 128 is a one-step (hospital-use) germicidal (disinfectant) cleaner and deodorant (odor-counteractant) (odor neutralizer) designed for general cleaning, (and) disinfecting, (deodorizing) (and controlling mold and mildew on) (of) hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew, body oils and other common soils found in hospitals, nursing homes, schools and colleges, (day care centers), (medical) offices, funeral homes, veterinary clinics, pet shops, (equine farms), animal life science laboratories, hotels, motels, public areas and restrooms, foodservice establishments (restaurants) and federally inspected meat and poultry establishments (food [processing] plants).

It is designed for use on (Use daily on) (the following) (hard, non-porous environmental surfaces:) vinyl, painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, stainless steel, aluminum, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilets, urinals, sinks, shower rooms and locker rooms areas - any washable (food and non-food contact) surface where disinfection is required. VIREX™ II/128's non-dulling formula eliminates the time and labor normally required for rinsing. A potable water rinse is required for food contact surfaces. Do not use on glasses, dishes and utensils.

(Food Service:)

VIREX™ II/ 128 cleans by removing dirt, grime and food soils in food preparation and processing areas. Its non-abrasive formula will not harm (scratch) surfaces. It cleans, disinfects and eliminates odors leaving surfaces smelling clean and fresh. Use where odors are a problem.

Its non-abrasive formula is designed for use on (Use daily on) (the following) hard, non-porous environmental surfaces: vinyl, painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, aluminum, stainless steel, brass, copper, laminated surfaces and baked enamel surfaces (associated with floors, walls, ceilings, tables, chairs, countertops, fixtures, glazed tile, toilets, (toilet bowls), urinals, sinks found in food establishments, (restaurants), (commercial kitchens) & restrooms. A potable water rinse is required for food contact surfaces.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 10627-21

(Hospitals/Health Care Facilities:)

VIREX™ II/ 128 is a one-step (hospital-use) germicidal (disinfectant) cleaner and deodorant (odor-counteractant) (odor neutralizer) designed for general cleaning, (and) disinfecting, (deodorizing) (and controlling mold and mildew on) (of) a hard, non-porous environmental surfaces. VIREX™ II/ 128 cleans quickly by removing dirt, grime, mold, mildew, food residue, body oils, dead skin, blood and other organic matter commonly found in hospitals (in health care facilities) (on medical surfaces). It (also) eliminates odors leaving (restroom) surfaces smelling clean and fresh. Use where odors are a problem.

VIREX™ II/ 128 cleans, disinfects and deodorizes (hard, non-porous environmental hospital (medical) surfaces) in one step (with no rinsing required). Its non-abrasive formula is designed for use on (Use daily on) (the following) (hard, non-porous environmental surfaces): vinyl, painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, stainless steel, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilets, urinals, sinks found in (health care facilities [hospitals],) patient rooms, operating rooms, ICU areas, shower rooms, and locker rooms. (It can also be used to pre-clean and disinfect hospital items: wheelchairs, [hospital] [patient] bed rails and linings, wash basins, bed pans, medical equipment surfaces) - any washable (food and non-food contact) surface (where disinfection is required). A potable water rinse is required when disinfecting food contact surfaces. Do not use on glasses, dishes and utensils.

(Animal Housing Facilities:)

VIREX™ II/ 128 cleans by removing dirt, grime, mold, mildew, blood, urine, fecal matter and other common soils found in animal housing facilities, livestock, swine or poultry facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It (also) eliminates odors leaving surfaces smelling clean and fresh.

VIREX™ II/ 128 cleans, disinfects and deodorizes (hard, non-porous environmental surfaces) in one step. Its non-abrasive formula is designed for use on (Use daily on) (Use daily to clean and disinfect) hard, non-porous surfaces: plated or stainless steel, aluminum, chrome, glazed porcelain, glazed tile, laminated surfaces (associated with floors, walls, countertops, cages, kennels, animal equipment) found in (barns, pens and stalls) animal housing facilities.

(Public Restrooms:)

VIREX™ II/ 128 is a one-step disinfectant cleaner and deodorant (odor-counteractant) (odor-neutralizer) designed for general cleaning, (and) (disinfecting) (deodorizing) (and controlling mold and mildew) on (of) of hard, non-porous environmental surfaces.

VIREX™ II/ 128 cleans, disinfects and deodorizes surfaces by killing odor-causing germs and mold & mildew. Its non-abrasive formula is designed for use on (restroom surfaces): glazed ceramic (restroom) tile, glazed porcelain, chrome, stainless steel and plastic surfaces associated with floors, walls, fixtures, toilets, urinals, sinks, shower rooms and locker rooms.

(Non-Acid Bowl [& Bathroom] Disinfectant Cleaner:)

VIREX™ II/ 128 is a (concentrated) non-acid (bowl and) bathroom cleaner which cleans, disinfects and deodorizes in one easy step. It cleans, disinfects and deodorizes toilet bowls, urinals, bowl brushes (bowl mops), rims, sinks, sink basins, faucets, tubs, glazed tiles, glazed ceramic, glazed porcelain, chrome, stainless steel, and all hard non-porous, washable surfaces found in the bathroom (restroom) (in the presence of organic soil).

VIREX™ II/ 128 eliminates odors leaving bathrooms (restrooms) smelling clean and fresh. Use where odors are a problem. It cleans, disinfects and deodorizes surfaces by killing odor-causing microorganisms and mold & mildew. Its non-abrasive formula is designed for use (Use it daily) on ([hard, non-porous environmental] restroom surfaces): glazed ceramic (restroom) tile, glazed porcelain, chrome, stainless steel and plastic surfaces associated with floors, walls, fixtures, toilets, urinals, sinks, shower rooms and locker rooms.

(Refill)

To Refill Concentrate From Large Containers Into Smaller Containers:)

VIREX™ II/ 128 may be used to fill and refill clean, properly labeled containers for dilution elsewhere within your facility. Make sure the small container has been cleaned, dried and properly labeled. Also make sure other items (funnels or hand pumps) are properly cleaned and dried. To refill, simply pour (or pump product) from the larger container directly into the smaller one being careful not to spill any product. Keep both containers sealed when not in use. Refill in a well ventilated area.

ACCEPTED
with COMMENTS
in EPA Letter Dated:
APR 02 2001

(CLAIMS:)

When used as directed at a 1:128 dilution (1oz. per gallon of water) (8mL/L), VIREX™ II/ 128 contains 660 ppm of active quaternary germicide making it highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms (including bacteria, antibiotic resistant bacteria, viruses, fungi, mold and mildew. See reference sheet (enclosed in each case) for a complete list of organisms).

Using approved AOAC test methods (under Good Laboratory Practices, [GLP's]), in the presence of 400 ppm hard water, 5% serum load and 10 minutes contact time VIREX™ II/ 128 kills the following:

Viruses (*Virucidal Activity) – kills on hard non-porous inanimate surfaces.

Adenovirus Type 2, (VR-2)

Fungi/Yeast (Fungicidal and Yeast Activity) – kills on hard non-porous inanimate surfaces.

Fungi:

Aspergillus niger, (ATCC 6275)

Trichophyton mentagrophytes (athlete's foot fungus), (ATCC 9533)

Yeast:

Candida albicans, (ATCC 10231)

Mold/Mildew (Mildewcidal Activity) - kills the growth of mold and mildew: *Aspergillus niger* (ATCC 6275) (and the odors caused by them when applied to hard, non-porous environmental surfaces).

Using approved AOAC test methods (under Good Laboratory Practices, [GLP's]), in the presence of 400 ppm hard water, 10% serum load and 10 minutes contact time VIREX™ II/ 128 kills the following:

Bacteria (Bactericidal Activity) - kills on hard non-porous inanimate surfaces:

Pseudomonas aeruginosa, (ATCC 15442)
Staphylococcus aureus, (ATCC 6538)
Salmonella choleraesuis, (ATCC 10708)
Acinetobacter calcoaceticus, (ATCC 9957)
Bordetella bronchiseptica, (ATCC 10590)
Burkholderia cepacia, (ATCC 25416)
formerly known as *Pseudomonas cepacia*
Campylobacter fetus, (ATCC 27374)
Citrobacter freundii, (ATCC 8090)
Enterobacter agglomerans, (ATCC 27155)
Enterobacter cloacae, (ATCC 23355)
Enterobacter liquefaciens, (ATCC 14460)
Enterococcus faecalis, (ATCC 19433)
formerly known as *Streptococcus faecalis*
Enterococcus hirae, (ATCC 10541)
Escherichia coli, (ATCC 11229)
Escherichia coli 0157:H7, (ATCC 43890)
Flavobacterium meningosepticum, (ATCC 13253)
Haemophilus influenza, (ATCC 10211)

Hafnia alvei, (ATCC 13337)
Klebsiella oxytoca, (ATCC 13182)
Klebsiella pneumoniae, (ATCC 13883)
Legionella pneumophila, (ATCC 33153)
Listeria monocytogenes, (ATCC 15313)
Micrococcus luteus, (ATCC 4698)
Micrococcus luteus, (ATCC 14452)
Micrococcus sedentarius, (ATCC 27573)
Neisseria gonorrhoeae, (ATCC 43069)
Pasteurella multocida, (ATCC 43137)
Proteus mirabilis, (ATCC 9240)
Proteus vulgaris, (ATCC 13315)
Pseudomonas diminuta, (ATCC 11568)
Pseudomonas fluorescens, (ATCC 13525)
Pseudomonas putida, (ATCC 12633)
Pseudomonas stutzeri, (ATCC 17588)
Salmonella choleraesuis pullorum, (ATCC 19945)
Salmonella enteritidis, (ATCC 13076)
Salmonella gallinarum, (ATCC 9184)

Salmonella schottmuelleri, (ATCC 10719)
Salmonella typhi, (ATCC 6539)
Salmonella typhimurium, (ATCC 13311)
Serratia grimesii, (ATCC 14460)
Serratia marcescens, (ATCC 9103)
Shigella dysenteriae, (ATCC 29026)
Shigella flexneri, (ATCC 25875)
Shigella sonnei, (ATCC 25931)
Staphylococcus aureus, (ATCC 25923)
Staphylococcus aureus (Toxic Shock), (ATCC 33586)
Staphylococcus epidermidis, (ATCC 14990)
Staphylococcus haemolyticus, (ATCC 29970)
Streptococcus agalactiae, (ATCC 13813)
Streptococcus mutans, (ATCC 25175)
Streptococcus pyogenes, (ATCC 19615)
Streptococcus pyogenes ("Strep A" - Flesh Eating Strain), (clinical isolate)
Vibrio cholera, (ATCC 11623)
Yersinia enterocolitica, (ATCC 9610)

Antibiotic-Resistant (Strains of) Bacteria (Antibiotic-Resistant Bactericidal Activity) - kills on hard non-porous inanimate surfaces:

E. coli, (ATCC 55244);
(Resistant to Kanamycin)

Klebsiella oxytoca, (ATCC 15764);
(Resistant to Ampicillin, Dihydrostreptomycin)

Staphylococcus aureus, (ATCC 14154);
(Resistant to Erythromycin, Penicillin, Streptomycin, Tetracycline)

E. coli, (ATCC 47041);
(Resistant to Tetracycline)

Micrococcus sedentarius, (ATCC 27573);
(Resistant to Methicillin)

Staphylococcus aureus, (ATCC 33592)
(Resistant to Methicillin [MRSA], Gentamicin [GRSA])

Enterococcus faecalis, (ATCC 51299);
(Resistant to Vancomycin [VRE])

Staphylococcus aureus, (CDC HIP 5836);
(Resistant to Vancomycin [VRSA])

Streptococcus pneumoniae, (ATCC 51915);
(Resistant to Penicillin [PRSP])

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 70627-21

To Prepare Use Solution:

Add the product at 1 oz. per gallon of water (8mL/L) (1:128).

(Note to EPA: The directions that appear for specific container sizes will be substituted for the sentence above only on that container size.)

ACCUMIX™ CONTAINERS (32 oz. container) - Add the product at 1 oz. per gallon of water (8 mL/L) (1:128).

SOLUTIONS CENTER™ (64 oz. containers) - Remove cap and insert cartridge into dispenser. Note: See dispenser instructions for proper cartridge placement. Once cartridge is in place, squeeze the handle or press the button to dispense a 1:128 solution into a bucket, bottle, scrubber or other container.

J-FILL™ DISPENSING SYSTEMS (84.5 oz. containers) - Remove cap and insert cartridge into dispenser. Note: See dispenser instructions for proper cartridge placement. Once cartridge is in place, squeeze the handle or press the button to dispense a 1:128 solution into a bucket, bottle, scrubber or other container.

ACCUPACK™ CONTAINERS (Pre-measured packets) - (Simply) open and pour contents into xx gallons of water (for a 1:128 dilution). Keep packets in box until ready to use. (Pour contents of packet into xx gallons of water for a 1:128 dilution.)

(Special Instructions for ACCUPACK CONTAINERS.)

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls from Concentrate:

1. Pre-clean heavily soiled areas.
2. Pour packet contents into toilet bowl.
3. Swab entire surface area especially under the rim.
4. Allow entire surface to remain wet for ten (10) minutes.
5. Flush toilet and rinse swab applicator thoroughly.

J-SPRAY™ DISPENSING SYSTEM

(Cartridge Placement Instructions:)

1. Remove side cap, and fill bottle with water to line. Replace cap.
2. Open bottle top by removing (black) (gray) neck ring. Remove snap cap from cartridge and insert J-Spray™ Virex™ II/ 128 concentrate cartridge into neck of bottle. Replace (black) (gray) neck ring.
3. Insert trigger spray dip tube into center cartridge hole, and gently push plastic plunger down through one of the sealed foil compartments. Push until top and bottom seals are broken. **DO NOT PIERCE MORE THAN ONE COMPARTMENT AT A TIME.**
4. Screw trigger sprayer down tightly, and gently agitate to mix solution.
5. Affix label provided.

(Refill Instructions:)

When trigger spray bottle is empty, refill with water. Leaving (black) (gray) neck ring on, lift trigger sprayer and pierce next cartridge section following steps 3 & 4. Replace cartridge when all four sections have been used. Use only in bottles labeled: J-Spray™ Virex™ II/ 128 Use Solution.

For Use as a One-Step Cleaner/Disinfectant Product:

1. Pre-clean heavily soiled areas.
2. Apply Use Solution to hard, non-porous environmental surfaces.
3. All surfaces must remain wet for ten (10) minutes.
4. Wipe surfaces (and let air dry).

Note - Rinsing is not necessary unless floors are to be coated with finish or restorer. All food contact surfaces such as appliances and kitchen countertops must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 70627-21

For Use as a Cleaner and/or Deodorizer:

Apply Use Solution to surfaces. Wipe surfaces (and let air dry).

For Use as a Cleaner/Disinfectant in Food Processing Plants:

1. Before using this product in food processing areas, food products and packaging materials must be removed from the room or carefully protected.
2. Apply Use Solution evenly over surface. Be sure to wet all surfaces thoroughly.
3. Allow product remain on surface for ten (10) minutes.
4. Wipe with clean cloth, sponge or paper towel.
5. For heavily soiled areas, thoroughly clean surface prior to disinfecting.
6. When disinfecting food contact surfaces such as kitchen counters and tables used for food preparation, cutting boards, appliances and sinks, then thoroughly rinse surfaces with potable water. This product must not be used to clean the following surfaces: utensils, glassware, and dishes.

For Use To Clean and Disinfect Barber and Beauty/Manicure Instruments and Tools:

1. Pre-clean heavily soiled items.
2. Completely immerse pre-cleaned combs, brushes, scissors, clipper blades, razors, manicure implements and other non-porous instruments in the Use Solution so that surfaces remain wet for ten (10) minutes.
3. Rinse surfaces thoroughly and let air dry before reuse.
4. Change solution daily or when visibly dirty.

NOTE: Plastics may remain immersed until ready to use. Stainless steel shears and other metal instruments must be removed after 10 minutes, rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged soaking may cause damage to metal instruments.

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls from Concentrate:

1. Pre-clean heavily soiled areas.
2. Add 3/4 oz. into toilet bowl.
3. Swab entire surface area especially under the rim.
4. Allow entire surface to remain wet for ten (10) minutes.
5. Flush toilet and rinse swab applicator thoroughly.

For Use as a Non-Acid Bowl Cleaner/Disinfectant in Toilet Bowls [and Urinals] from Use-Dilution:

1. Pre-clean heavily soiled areas.
2. Empty toilet bowls by forcing water through the trap. Apply Use Solution to exposed surfaces in toilet bowls and urinals.
3. Swab entire surface area especially under the rim.
4. Allow entire surface to remain wet for ten (10) minutes.
5. Flush toilet or urinal and rinse swab applicator thoroughly.

For Use To Clean and Disinfect Shower Rooms, Locker Rooms and Other Large, Open Areas with Floor Drains:

1. Pre-clean heavily soiled areas.
1. Apply Use Solution to floors, walls and ceilings making sure not to over spray. To disinfect, all surfaces must remain wet for ten (10) minutes.
Special instructions for foam guns: Pour concentrate into foam gun bottle and attach bottle to spray nozzle and ensure gun is attached to hose. Note: See foam gun instructions for more information. Make sure setting is set for a 1:128 dilution. Once in place, squeeze the handle to dispense foam solution. To disinfect, all surfaces must remain wet for ten (10) minutes.
3. Scrub using a deck brush or other coarse material as necessary.
4. Rinse surfaces thoroughly and let air dry.

To Kill Mold and Mildew (in 5% soil load):

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surfaces to remain wet for ten (10) minutes. Wipe surfaces (and let air dry).

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 50627-21

1/4/01
Page 7 of 12
EPA Reg. No. 70627-21

To Control Mold and Mildew:

Apply Use Solution to hard, non-porous environmental surfaces. Allow to air dry. Repeat application weekly or when growth reappears.

To Kill Fungi:

Pre-clean heavily soiled areas. Apply Use Solution to hard, non-porous environmental surfaces. Allow surface to remain wet for ten (10) minutes. Wipe surfaces (and let air dry).

For Use For Treatment of Animal Housing Facilities:

1. Remove all animals and feed from areas being treated.
2. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes, and other facilities occupied or traversed by animals.
3. Empty or cover all troughs, racks and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap and rinse with water.
5. Apply fresh use solution to floors, walls, cages and other washable hard, non-porous environmental surfaces. For smaller surfaces, use a trigger spray bottle to spray all surfaces with solution until wet. To disinfect, all surfaces must remain wet for ten (10) minutes.
6. Do not house animals or employ equipment until product has dried.
7. For disinfection of feed racks, troughs, automatic feeders, fountains and watering appliances scrub with use-solution, let stand ten (10) minutes. Then thoroughly scrub all treated surfaces with soap or detergent and rinse with potable water before reuse.

***VIREX™ III/ 128 kills HIV-1 on pre-cleaned environmental surfaces/objects previously soiled with blood/body fluids in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS).**

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 ON SURFACES/ OBJECTS SOILED WITH BLOOD/BODY FLUIDS.

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood and body fluids must be thoroughly cleaned from surfaces and objects before application of VIREX™ III/ 128.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1 and for 10 minutes to kill all other organisms cited on the label.

Disposal of Infectious Material: Blood and other body fluids should be autoclaved and disposed of according to federal, State, and local regulations for infectious waste disposal.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Do not reuse empty container unless refilling from a larger container of the same product according to the refilling directions outlined previously. (Keep from freezing).

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 70627-21

Johnson Wax PROFESSIONAL

12/13

VIREX™ II/ 128 REFERENCE SHEET

When used as directed at a 1:128 dilution (1oz. per gallon of water) (8mL/L), VIREX™ II/ 128 contains 660 ppm of active quaternary germicide making it highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms (including bacteria, antibiotic resistant bacteria, viruses, fungi, mold and mildew. See reference sheet (enclosed in each case) for a complete list of organisms).

Using approved AOAC test methods (under Good Laboratory Practices, [GLP's]), in the presence of 400 ppm hard water, 10% serum load and 10 minutes contact time VIREX™ II/ 128 kills the following:

Bacteria (Bactericidal Activity) - kills on hard non-porous inanimate surfaces:

<i>Pseudomonas aeruginosa</i> , (ATCC 15442)	<i>Hafnia alvei</i> , (ATCC 13337)	<i>Salmonella schottmuelleri</i> , (ATCC 10719)
<i>Staphylococcus aureus</i> , (ATCC 6538)	<i>Klebsiella oxytoca</i> , (ATCC 13182)	<i>Salmonella typhi</i> , (ATCC 6539)
<i>Salmonella choleraesuis</i> , (ATCC 10708)	<i>Klebsiella pneumoniae</i> , (ATCC 13883)	<i>Salmonella typhimurium</i> , (ATCC 13311)
<i>Acinetobacter calcoaceticus</i> , (ATCC 9957)	<i>Legionella pneumophila</i> , (ATCC 33153)	<i>Serratia grimesii</i> , (ATCC 14460)
<i>Bordetella bronchiseptica</i> , (ATCC 10580)	<i>Listeria monocytogenes</i> , (ATCC 15313)	<i>Serratia marcescens</i> , (ATCC 9103)
<i>Burkholderia cepacia</i> , (ATCC 25416)	<i>Micrococcus luteus</i> , (ATCC 4698)	<i>Shigella dysenteriae</i> , (ATCC 29028)
formerly known as <i>Pseudomonas cepacia</i>	<i>Micrococcus luteus</i> , (ATCC 14452)	<i>Shigella flexneri</i> , (ATCC 25875)
<i>Campylobacter fetus</i> , (ATCC 27374)	<i>Micrococcus sedentarius</i> , (ATCC 27573)	<i>Shigella sonnei</i> , (ATCC 25931)
<i>Citrobacter freundii</i> , (ATCC 8090)	<i>Neisseria gonorrhoea</i> , (ATCC 43069)	<i>Staphylococcus aureus</i> , (ATCC 25923)
<i>Enterobacter agglomerans</i> , (ATCC 27155)	<i>Pasteurella multocida</i> , (ATCC 43137)	<i>Staphylococcus aureus</i> (Toxic Shock), (ATCC 33586)
<i>Enterobacter cloacae</i> , (ATCC 23355)	<i>Proteus mirabilis</i> , (ATCC 9240)	<i>Staphylococcus epidermidis</i> , (ATCC 14990)
<i>Enterobacter liquefaciens</i> , (ATCC 14460)	<i>Proteus vulgaris</i> , (ATCC 13315)	<i>Staphylococcus haemolyticus</i> , (ATCC 29970)
<i>Enterococcus faecalis</i> , (ATCC 19433)	<i>Pseudomonas diminuta</i> , (ATCC 11568)	<i>Streptococcus agalactiae</i> , (ATCC 13813)
formerly known as <i>Streptococcus faecalis</i>	<i>Pseudomonas fluorescens</i> , (ATCC 13525)	<i>Streptococcus mutans</i> , (ATCC 25175)
<i>Enterococcus hirae</i> , (ATCC 10541)	<i>Pseudomonas putida</i> , (ATCC 12633)	<i>Streptococcus pyogenes</i> , (ATCC 19615)
<i>Escherichia coli</i> , (ATCC 11229)	<i>Pseudomonas stutzeri</i> , (ATCC 17588)	<i>Streptococcus pyogenes</i> ("Strep A" - Flesh Eating Strain), (clinical isolate)
<i>Escherichia coli</i> 0157:H7, (ATCC 43890)	<i>Salmonella choleraesuis pullorum</i> , (ATCC 19945)	<i>Vibrio cholera</i> , (ATCC 11623)
<i>Flavobacterium meningosepticum</i> , (ATCC 13253)	<i>Salmonella enteritidis</i> , (ATCC 13076)	<i>Yersinia enterocolitica</i> , (ATCC 9610)
<i>Haemophilus influenza</i> , (ATCC 10211)	<i>Salmonella gallinarum</i> , (ATCC 9184)	

Antibiotic-Resistant (Strains of) Bacteria (Antibiotic-Resistant Bactericidal Activity) - kills on hard non-porous inanimate surfaces:

<i>E. coli</i> , (ATCC 55244); (Resistant to Kanamycin)	<i>Klebsiella oxytoca</i> , (ATCC 15764); (Resistant to Ampicillin, Dihydrostreptomycin)	<i>Staphylococcus aureus</i> , (ATCC 14154); (Resistant to Erythromycin, Penicillin, Streptomycin, Tetracycline)
<i>E. coli</i> , (ATCC 47041); (Resistant to Tetracycline)	<i>Micrococcus sedentarius</i> , (ATCC 27573); (Resistant to Methicillin)	<i>Staphylococcus aureus</i> , (ATCC 33592) (Resistant to Methicillin [MRSA], Gentamicin [GRSA])
<i>Enterococcus faecalis</i> , (ATCC 51299); (Resistant to Vancomycin [VRE])	<i>Staphylococcus aureus</i> , (CDC HIP 5836); (Resistant to Vancomycin [VRSA])	<i>Streptococcus pneumoniae</i> , (ATCC 51915); (Resistant to Penicillin [PRSP])

Viruses (*Virucidal Activity) - kills on hard non-porous inanimate surfaces:

<i>Cytomegalovirus</i> , (VR-538)	<i>Herpes simplex Type 2</i> , (VR-734)	<i>Parainfluenza Type 3</i> , (VR-93)
<i>Herpes simplex Type 1</i> , (VR-733)	<i>Influenza Type A₂</i> (Hong Kong), (VR-544)	<i>Respiratory syncytial virus</i> , (VR-25)
		<i>Vaccinia virus</i> , (VR-119)

Kills HIV-1 (AIDS virus) (HTLV-III_B) when used as directed on hard, non-porous inanimate surfaces with a 1 minute contact time.

(Veterinary viruses:)

<i>Avian infectious bronchitis</i> (IBV), (VR-22)	<i>Chlamydia psittaci</i> , (VR-125)	<i>New Castle disease</i> , (VR-108)
<i>Avian Influenza</i> , (VR-2072)	<i>Feline viral rhinotracheitis</i> , (VR-636)	<i>Pseudorabies</i> , (VR-105)
<i>Canine distemper</i> , (VR-128)	<i>Infectious bovine rhinotracheitis</i> , (VR-188)	<i>Transmissible gastroenteritis virus</i> (TGEV), (U of Minn. Strain)

ACCEPTED
with COMMENTS
in EPA Letter Dated:

APR 02 2001

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No. 70627-21

1/4/01
Page 11 of 12
EPA Reg. No. 70627-21

