



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:
70627-37

Date of
4/6/2001

Term of Issuance:

Conditional

Name of Pesticide Product:

Johnson Wax Professional
Cockroach Gel Bait Formula 3

NOTICE OF PESTICIDE:

- Registration
- Reregistration

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

SC Johnson Commercial Markets, Inc.
8310 16th St, PO Box 902, MS 675
Sturtevant, WI 53117-0902

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Make the following label changes before you release the product for shipment:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. 70627-37".
3. Submit two copies of the revised final printed label for the record before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A copy of the label stamped "accepted with comments" is enclosed for your records.

Signature of Approving Official:

Thomas C. Harris, Biologist
Insecticide-Rodenticide Branch
Registration Division (7504C)

Date:

APR 6 2001

Page 1 of 4 Carton Front Panel

Master Label consists of Carton Front Panel, Other Carton Panels, Applicator and Supplemental Labeling

Johnson wax
PROFESSIONAL

Cockroach
Gel Bait
FORMULA 3

Recommended for use by Commercial Applicators For Commercial/Industrial Use

Controls: Cockroaches

(including carbamate, organophosphate, organochlorine, and pyrethroid resistant strains).

FOR USE IN AND AROUND: Commercial, Industrial, and Residential Buildings including Apartments, Garages, Food Storage Areas, Homes, Hospitals and Nursing Homes, Hotels, Meat Packing and Food Processing Plants, Motels, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains, Planes) and Warehouses.

KEEP OUT OF REACH OF CHILDREN
CAUTION

ACTIVE INGREDIENT:

Abamectin B1 0.05%

INERT INGREDIENTS: 99.95%

May be used in Federally Inspected Meat and Poultry Plants.

See side panels for precautionary statements.


See supplemental labeling for complete

Directions for Use.

Total Net Wt.: 90g,

Net Wt. Each: 30g per unit

EPA Reg. No. 70627-GT
EPA Est. No. 499-MO-1



70627-37

ACCEPTED
with COMMENTS
In EPA Letter Dated:
APR 6 2001
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.

Page 2 of 4 Other Carton Panels

Master Label consists of Carton Front Panel, Other Carton Panels, Applicator and Supplemental Labeling

Cockroach Gel Bait

Johnson Wax PROFESSIONAL

Cockroach Gel Bait FORMULA 3

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See supplemental labeling for complete Directions for Use.

GENERAL USE DIRECTIONS: Johnson Wax Professional Cockroach Gel Bait Formula 3 is for use in controlling cockroaches indoors and outdoors. Inject into cracks and crevices, void spaces and other locations where insects may be harboring, living and breeding. Concentrate the bait placements as individual spots or a bead at insect activity sites.

APPLICATION RATES - ALL USES: For light infestations of Cockroaches, 4 - 6 bait points are recommended per 100 sq. ft. of treatment area. For heavy infestations of Cockroaches, 12 - 24 bait points are recommended per 100 sq. ft. of treatment area. Each bait point should equal about 0.25 to 0.50 grams of product. A spot approximately 3 mm or 1/8 inch in diameter equals 0.25 grams of product, while a spot approximately 6 mm or 1/4 inch in diameter equals 0.50 grams of product.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool, dry area.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL:

APPLICATOR: Completely empty applicator by using the product according to the label. Replace cap, wrap the container, and discard in the trash, sanitary landfill, or by other approved state and local procedures.

Sold by:

© 2001 S.C. Johnson Commercial Markets, Inc., Sturtevant, WI 53177-0902 U.S.A.
All Rights Reserved.



Johnson Wax PROFESSIONAL

Cockroach Gel Bait

FORMULA 3

Controls: Cockroaches



PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash contaminated clothing before re-use.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PRECAUTIONS

- Do not use on or contaminate fruit, vegetables or other food or feed crops.
- Do not apply to humans, animals, clothing or bedding.
- Do not contaminate feed or food products or food preparation surfaces, dishes, kitchen utensils and food containers.
- To prevent staining, exposed bait should be cleaned up immediately with a damp cloth or sponge.

PHYSICAL OR CHEMICAL HAZARDS

Do not treat directly into any electronic equipment such as radios, televisions, computers, etc. Do not treat where electrical short circuits might result.

Johnson Wax PROFESSIONAL

Cockroach Gel Bait

FORMULA 3

Page 3 of 4 Applicator

Master Label consists of Carton Front Panel, Other Carton Panels, Applicator and Supplemental Labeling

Johnson wax
PROFESSIONAL

Cockroach
Gel Bait

FORMULA 3

Applicator

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

See supplemental label for complete Directions for Use.

EPA Reg. No. 70627-GT
EPA Est. No. 499-MO-1

Sold by:
©2001 S.C. Johnson Commercial Markets, Inc.,
Sturtevant, WI 53177-0902 U.S.A.
All Rights Reserved.

Recommended for use by
Commercial Applicators For
Commercial/Industrial Use



KEEP OUT OF REACH OF CHILDREN

CAUTION

ACTIVE INGREDIENT:

Abamectin B1 0.05%

INERT INGREDIENTS: 99.95%

May be used in Federally Inspected
Meat and Poultry Plants.

See supplemental label for additional
precautionary statements.

Net Wt. 30 grams



SUPPLEMENTAL LABELING



Cockroach Gel Bait Formula 3

Johnson Wax Professional Cockroach Gel Bait Formula 3 is a slow acting insect gel bait. Control of an ant colony may take several weeks. For best results placement of the bait should be into or near the insect colony.

CONTROLS: Cockroaches (including carbamate, organophosphate, organochlorine and pyrethroid resistant strains) and Ants (including, but not limited to, acrobat, allegheny, argentine, bigheaded, carpenter, corn field, crazy, fire, ghost, harvester, little black, odorous house, pavement, pharaoh, pyramid).

FOR USE IN AND AROUND: Commercial, Industrial and Residential Buildings including Apartments, Garages, Food Storage Areas, Homes, Hospitals and Nursing Homes, Hotels, Meat Packing and Food Processing Plants, Motels, Restaurants and other Food Handling Establishments, Schools, Supermarkets, Transportation Equipment (Buses, Boats, Ships, Trains and Planes) and Warehouses.

May be used in Federally Inspected Meat and Poultry Plants.

KEEP OUT OF REACH OF CHILDREN

CAUTION

ACTIVE INGREDIENT:

Abamectin B10.05%

OTHER INGREDIENTS:99.95%

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Wash hands and exposed skin before eating, drinking or smoking and after handling. Wash all contaminated clothing thoroughly before re-use.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

PHYSICAL OR CHEMICAL HAZARDS

Do not treat directly into any electronic equipment such as radios, televisions, computers, etc. Do not treat where electrical short circuits might result.

EPA Reg. No. 70627-GT EPA Est. No. 499-MO-1

Questions? Comments: 1-800-558-2332

Sold by: ©2001 S.C. Johnson Commercial Markets, Inc., Sturtevant, WI 53177-0902 U.S.A.

All Rights Reserved

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PRECAUTIONS:

- Do not use on or contaminate fruit, vegetables or other food or feed crops.
- Do not apply to humans, animals, clothing or bedding.
- Do not contaminate feed or food products or food preparation surfaces, dishes, kitchen utensils and food containers.
- To prevent staining, exposed bait should be cleaned up immediately with a damp cloth or sponge.

GENERAL USE DIRECTIONS: Johnson Wax Professional Cockroach Gel Bait Formula 3 is for use in controlling cockroaches indoors and outdoors. Inject into cracks and crevices, void spaces and other locations where insects may be harboring, living and breeding. Concentrate the bait placements as individual spots or a bead at insect activity sites.

APPLICATION RATES - ALL USES: For light infestations of Cockroaches, 4 - 6 bait points are recommended per 100 sq. ft. of treatment area. For heavy infestations of Cockroaches, 12 - 24 bait points are recommended per 100 sq. ft. of treatment

Page 4 of 4 SUPPLEMENTAL LABELING

Master Label consists of Carton Front Panel, Other Carton Panels, Applicator and Supplemental Labeling

area. Each bait point should equal about 0.25 to 0.50 grams of product. A spot approximately 3 mm or 1/8 inch in diameter equals 0.25 grams of product, while a spot approximately 6 mm or 1/4 inch in diameter equals 0.50 grams of product.

FOR ANTS: Place several bait points as close as possible to the site of actual or suspected activity.

APPLICATION IN ALL EXCEPT FOOD/FEED HANDLING AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

For treatment in residential and the non-food/non-feed areas of institutional, warehousing and commercial establishments, place bait into cracks and crevices; holes; pipe chases; undersides of furniture; under drain plates; in or under trash containers; hidden surfaces around sinks and storage areas; behind baseboards; around doors and windows; inside, behind and under cabinets, drawers and shelving; under and behind appliances such as stoves and refrigerators; and in attics and crawl spaces. Also apply in points between different elements of construction, between equipment and floors, openings leading to voids and hollow spaces in walls, equipment legs and bases and crawl spaces where insects hide. During follow-up visits, inspect bait placements and re-apply when necessary. Care should be taken to avoid depositing the product onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface. This product may also be used in food/feed areas of food/feed handling establishments as outlined below.

Non-food/feed areas are areas such as garbage rooms, lavatories, floor drains (to sewers), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage areas (after canning or bottling).

APPLICATIONS IN FOOD/FEED AREAS OF FOOD/FEED HANDLING ESTABLISHMENTS

Application within food/feed areas of food/feed handling establishments is limited to Crack and Crevice treatment only. Apply Johnson Wax Professional Cockroach Gel Bait Formula 3 directly into cracks and crevices. Apply small amounts of gel (up to 0.50 gram) per point. Do not apply bait to areas where food utensils or food processing surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food/feed preparation and prepared food holding surfaces as bait may be removed by washing. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface. Food/feed areas include areas for receiving, storage, packing (canning, bottling, wrapping, boxing), preparing edible waste storage, and enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas are also considered food/feed areas when food is exposed and facility is in operation. Apply bait in small amounts between different elements of construction, between equipment and floors, openings leading into voids and hollow spaces in floors, walls, ceilings, equipment legs and bases, around plumbing pipes, doors and windows, behind and under refrigerators, cabinets, sinks, closets, stoves and other equipment where cockroaches hide. During follow-up visits, inspect bait placements and re-apply when necessary.

Application of this product in food/feed areas of food/feed handling establishments other than as a crack and crevice treatment is not permitted. Do not treat food preparation surfaces.

APPLICATIONS IN AIRCRAFT

Application within passenger areas of aircraft is limited to Crack and Crevice treatment only. Apply Johnson Wax Professional Cockroach Gel Bait Formula 3 in areas between elements of construction (such as areas under sinks, openings around pipes, under shelving, where shelving meets other shelves or the wall, area between and around microwaves, where support braces attach to the main structure and where seats are attached to the floor). Use 12-24 bait points per 100 sq. ft. Do not apply bait to areas where food utensils or food preparation surfaces may become contaminated. Do not apply bait to areas that are routinely washed such as cracks and crevices in tops of tables, food preparation, and food holding surfaces as bait may be removed by washing. Care should be taken to avoid depositing gel onto exposed surfaces. If gel contacts an exposed surface, remove gel and wash exposed surface. During follow-up visits, inspect bait placements and re-apply when necessary. Do not apply any bait in cockpit area.

OUTDOOR USE

Use for control of cockroaches. Inject into cracks and crevices around windows and doors, porches, screens, eaves, patios, garages, under stairways and in open crawl spaces and other areas where pests may be nesting or trailing, such as tree holes and cracks in fence

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a cool dry area.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal:

Applicator - Completely empty applicator by using the product according to the label. Replace cap, wrap the container, and discard in the trash, sanitary landfill, or by other approved state and local procedures.