



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:

70627-56

Date of Issuance:

DEC 20 2005

NOTICE OF PESTICIDE:

Registration
 Reregistration

(under FIFRA, as amended)

Term of Issuance: Conditional

Name of Pesticide Product:

OXIVIR TB

Name and Address of Registrant (include ZIP Code):

JohnsonDiversey
8310 16th Street, MS 486
Sturtevant, WI 53177-0902

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following label changes:
 - a. Revise the EPA Registration Number to read, "EPA Reg. No. "70627-56".

Signature of Approving Official:

Marshall Swindell, PM 33
Regulatory Management Branch 1
Antimicrobial Division (7510C)

Date:

DEC 20 2005

2 8 11

Page 2


EPA Registration No. 70627-56

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

If you have any questions concerning this letter, please contact Martha Terry at (703) 308-6217.

Sincerely,



Marshall Swindell

Product Manager 33

Regulatory Management Branch 1


Antimicrobial Division (7510C)

Enclosure

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

3 8 11

JohnsonDiversey



OXIVIR TB
(Ready to Use)
One-Step Disinfectant and Deodorizer
General *Virucide, Bactericide, Tuberculocide, Fungicide, Sanitizer
Cleaner Disinfectant Deodorizer
Intermediate Disinfectant

(Hydrogen Peroxide)

For (Commercial) (Industrial) (&) (Institutional) Use (Only)

ACTIVE INGREDIENTS:	
Hydrogen Peroxide	0.5%
OTHER (INERT) INGREDIENTS	<u>99.5%</u>
TOTAL:	100.0%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See additional precautionary statements on (back) (side) (left) (right) (panel) (of) (label) (below).

See reference sheet (enclosed in each case) for (a completed list of organisms) (additional (features) (claims) (directions for use)) (for this product).

ACCEPTED
with COMMENTS
EPA Letter Dated:

DEC 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56

Net Contents:

Product of U.S.A.

4 8 11

(MARKETING CLAIMS):

(FEATURES, CLAIMS & USES:)

For Use in Health Care, Veterinary, Animal Housing, Dental, (Life Science) Laboratory, Medical, Industrial & Institutional Facilities, Food Processing Establishments, and Hospitals

(General Uses)

OXIVIR is a one-step (hospital-use) germicidal (disinfectant) cleaner and deodorant (odor-counteractant) (odor neutralizer) designed for general cleaning, (and) disinfecting (deodorizing) (of) hard, non-porous environmental surfaces. It cleans quickly by removing dirt, grime, mold, mildew and other common soils found in hospitals, nursing homes, schools & colleges, (medical) (dental) offices, veterinary clinics, pet shops, equine farms, animal & life science laboratories, transportation terminals (airports, train stations, bus stations, truck stops, rest (stops) areas), hotels, motels, casinos, cruise ships, other institutional/commercial buildings, homes cleaned by (home cleaning services,)(professional cleaning services,)(medical assistance services,) and office buildings.

OXIVIR's non-dulling formula eliminates the time and labor normally required for rinsing. It is designed for use on (Use daily on) (the following) (hard, non-porous environmental surfaces): (vinyl), (Naugahyde), painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, stainless steel, aluminum, laminated surfaces and baked enamel surfaces associated with floors, walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, sinks – almost any washable (food and non-food contact) surface where disinfection is required. A potable water rinse is required for food contact surfaces. Do not use on glasses, dishes and utensils.

(Hospitals/Health Care Facilities/Dental Offices/Life Science Laboratories)

OXIVIR is a one-step (hospital-use) (laboratory-use) germicidal (disinfectant) cleaner and deodorant (odor-counteractant) (odor neutralizer) designed for general cleaning, (and) disinfecting, (deodorizing) (of) hard, non-porous environmental surfaces. OXIVIR cleans quickly by removing dirt, grime, food residue, dead skin, blood and other organic matter commonly found in Hospitals, (in Health Care Facilities), (laboratory settings). It also eliminates odors, leaving (restroom) surfaces smelling clean and fresh. Use where odors are a problem.

OXIVIR cleans, disinfects and deodorizes hard, non-porous environmental (hospital) (medical) (dental) (laboratory) surfaces in one step (with no rinsing required). Its non-abrasive formula is designed for use on (Use daily on) (the following) (hard, non-porous environmental surfaces:) (vinyl), (Naugahyde), painted surfaces, plastic (surfaces), glazed ceramic, glazed porcelain, chrome, stainless steel, laminated surfaces (and baked enamel surfaces associated with floors), walls, ceilings, tables, chairs, countertops, telephones, fixtures, glazed tile, toilet exteriors, urinal exteriors, sinks found in (health care facilities), (hospitals), (dental offices), (patient rooms), (operating rooms), (ICU areas), (shower rooms), and (locker rooms) (fume hoods and animal cages [found in laboratory settings]). (It can be used to pre-clean and disinfect hospital items: wheelchairs, (hospital) (patient) bedrails and linings (patient bed linings) (vinyl mattress covers), wash basins, bed pans, non-critical medical equipment surfaces – almost any washable (food and non-food contact) surface (where disinfection is required). A potable water rinse is required when disinfecting food contact surfaces. Do not use on glasses, dishes and utensils.

(Animal Housing Facilities)

OXIVIR cleans by removing dirt, grime, blood, urine, fecal matter and other common soils found in animal housing facilities, grooming facilities, farms, kennels, pet stores, veterinary clinics, laboratories or other small animal facilities. It (also) eliminates odors leaving surfaces smelling clean and fresh.

ACCEPTED
with COMMENTS

EPA Letter Dated:

DEC 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56

5 8 11

This product cleans, disinfects and deodorizes hard, non-porous environmental surfaces in one step. Its non-abrasive formula is designed for use on (Use daily on) (Use daily to clean and disinfect) (hard, non-porous surfaces:) plated or stainless steel, chrome, glazed porcelain, glazed tile, laminated surfaces, (associated with floors, walls, countertops, cages, kennels, animal equipment) found in (barns, pens and stalls) animal housing facilities.

(CLAIMS:)

(When used as directed, this product (is) highly effective against a wide variety (broad-spectrum) of pathogenic microorganisms (including bacteria, antibiotic resistant bacteria, viruses, fungi, mold and mildew. See reference sheet (enclosed in each case) for a complete list of organisms).

HOSPITAL DISINFECTANT: BACTERICIDAL in the presence of 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces. (Hospital Disinfectant: (Bactericidal in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces. Bactericidal activity was determined by AOAC Use Dilution Test Method):

Pseudomonas aeruginosa (ATCC 15442)
Escherichia coli O157:H7 (ATCC 35150)
Acinetobacter baumannii (ATCC 19606)

Staphylococcus aureus (ATCC 6538)
Staphylococcus aureus MRSA (ATCC 33592)

Salmonella choleraesuis (ATCC 10708)
Enterococcus faecalis VRE (ATCC 51575)

Bactericidal activity was determined by the AOAC Use Dilution Test Method.

(***VIRUCIDAL** in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces.)

Viruses (*Virucidal Activity) - kills on hard non-porous inanimate surfaces:

Poliovirus Type 1, Strain Brunhilde (ATCC VR-1000)

HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1), (Acquired from Adv. Biotechnologies, Inc.)
Influenza A/Hong Kong (ATCC VR-544)

Feline Calicivirus, Strain F9 (ATCC VR-782)

Human Coronavirus (ATCC VR-740)

Herpes Simplex Virus, Type 1 (HSV-1) (ATCC VR-733)

Herpes Simplex Virus, Type 2 (HSV-2) (ATCC VR-734)

Rhinovirus Type 37, Strain 151-1 (ATCC VR-1147)

Rotavirus WA (Acquired from University of Ottawa)

Hepatitis C Virus (HCV) (Acquired from National Veterinary Services Laboratories)

Hepatitis B Virus (HBV) (Acquired from Hepadnavirus Testing)

TUBERCULOCIDAL in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Mycobacterium bovis (Tb) (OT 451C150)

Tuberculocidal activity was determined by the EPA Quantitative Tuberculocidal Activity Test Method.

FUNGICIDAL This product is fungicidal against the pathogenic fungi, *Tricophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% serum load and 10 minute contact time at 20°C (68°F) when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

Tricophyton mentagrophytes (ATCC 9533)

Fungicidal Activity was determined by the AOAC Fungicidal Activity of Disinfectants Method.

ACCEPTED
with COMMENTS
EPA Letter Dated:

DEC 20 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act, amended, for the pesticide, registered under EPA Reg. No. 70627-XX
10/25/2005
Page 3 of 9

70627-56

6 8 11

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

Non-Food Contact Surface Sanitizer for hard non-porous surfaces – in the presence of a 5% serum load and 30 second contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Klebsiella pneumoniae (ATCC 4352)
Staphylococcus aureus (ATCC 6538)
Escherichia coli 0157:H7 (ATCC 35150)

Salmonella choleraesuis (ATCC 10708)
Staphylococcus aureus MRSA (ATCC 33592)

Pseudomonas aeruginosa (ATCC 15442)
Enterococcus faecalis VRE (ATCC 51575)

Sanitizing activity was determined by the EPA Sanitizer Test for Inanimate, Non-Food Contact Surfaces and ASTM E1153 Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces.

Malodor(s) (Activity) (Counteractancy) – eliminates (destroys) odors and odor-causing bacteria in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors).

DIRECTIONS FOR USE:

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is not to be used as a terminal sterilant/high level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the blood stream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. This product may be used to pre-clean or decontaminate critical or semi-critical medical devices prior to sterilization or high level disinfection.

For Use as a One-Step Cleaner/Disinfectant Product:

1. Pre-clean heavily soiled areas.
2. Apply Solution by spray, cloth, disposable wipe or mop to hard, non-porous environmental surfaces.
3. All surfaces must remain wet for 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
4. Wipe surfaces dry (or rinse) (or allow to air dry).

Note: Rinsing is not necessary unless floors are to be coated with finish or restorer. All food contact surfaces must be rinsed with potable water. Do not use on glassware, utensils, or dishes.

For Use as a Cleaner and/or Deodorizer:

Apply to surfaces by spray, cloth, sponge, or mop. Wipe surfaces dry (or rinse) (or allow to air dry).

For Use to Clean and Disinfect Life Science Laboratory Surfaces:

1. Pre-clean heavily soiled areas.
2. Apply Solution with by spray, cloth, disposable wipe (paper towel) or mop to hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
4. Wipe surfaces dry (or rinse) (or allow to air dry).
5. Change solution after each use (application).

ACCEPTED
with COMMENTS
EPA Letter Dated:

DEC 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

7 7 11

For Use as a Tuberculocide:

1. Pre-clean heavily soiled areas.
2. Apply Solution with cloth, disposable wipe or mop to hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 5 minutes.
4. Wipe surfaces dry (or rinse) (or allow to air dry).
5. Change solution after each use (application).

ACCEPTED
with COMMENTS
EPA Letter Dated:

To Kill Fungi:

1. Pre-clean heavily soiled areas.
2. Apply Solution to hard, non-porous inanimate surfaces.
3. Allow surface to remain wet for 10 minutes.
4. Wipe surfaces dry (or rinse) (or allow to air dry).

DEC 20 2005
Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56

***KILLS HIV AND HCV and HBV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) and other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood/body fluids, and in which the surfaces/objects likely to be soiled with blood/body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) Hepatitis B Virus (HBV), or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATING AGAINST HIV-1 (AIDS Virus), HBV OR HCV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS:

Personal Protection: Disposable latex or vinyl gloves, gowns, face masks, or eye coverings as appropriate, must be worn during all cleaning of body fluids, blood, and decontamination procedures.

Cleaning Procedures: Blood/Body Fluids must be thoroughly cleaned from surfaces/objects before application of this product.

Contact Time: Allow surface to remain wet for 1 minute to kill HIV-1, HBV and HCV. Use a 5 minute contact time for Tb and a 10 minute contact for fungi.

Disposal of Infectious Material: Blood/Body Fluids should be autoclaved and disposed according to Federal, State and local regulations for infectious waste disposal.

Animal Housing Facilities (Including Veterinary Clinics, (Equine) Farms, Livestock, Swine and Poultry Houses): For cleaning and disinfecting the following hard, non-porous surfaces: feeding and watering equipment, utensils, instruments, cages, kennels, stables, catteries, etc.

1. Remove all animals and feeds from premises, animal transportation vehicles, crates, etc.
2. Remove all litter, droppings and manure from floors, walls and surfaces of facilities (including pens, chutes, and barns) occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding or watering appliances.
4. Thoroughly clean all surfaces with soap or detergent and rinse with potable water. Saturate surfaces by wiping or spraying with product for a period of 1 minute. Use a 5 minute contact time for TB and a 10 minute contact time for fungi.
5. Immerse handling and restraining equipment such as leashes, muzzles, halters or ropes, as well as forks, shovels, and scrapers used for removing litter and manure.
6. Ventilate buildings and other closed spaces.
7. Do not house animals or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub treated feed racks, troughs, mangers, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

8 2 11

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

For use as a Non-Food Contact Sanitizer:

1. Pre-clean heavily soiled areas.
2. Apply Solution with cloth, disposable wipe or mop to hard, non-porous environmental surfaces.
3. Allow the surface to remain wet for 30 seconds.
4. Wipe surfaces dry (or rinse) (or allow to air dry).
5. Change solution after each use (application).

(Refill)

To Transfer Oxivir TB Into Smaller Containers:

[Oxivir TB may be transferred into clean, properly labeled containers for use elsewhere within your facility.] The small container and transfer items (funnels or hand pumps) must be clean and dry. Pour (or pump) product directly into a properly labeled smaller container being careful not to spill any product. Keep containers sealed when not in use. Refill in a well ventilated area.

(For sizes 1 gallon or less)

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to small children. Keep securely closed. Do not reuse empty container. Rinse thoroughly. (Wrap and) Discard in trash.

(For sizes over 1 gallon)

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE

Store in a dry, well-ventilated area away from chemicals, direct light, heat or open flame. Do not mix with other cleaning or disinfecting products.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. For more information on the disposal of unused, unwanted product and the clean up of spills, contact the EPA Regional Office of the Manufacturer for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

ENVIRONMENTAL HAZARDS (for labels with veterinary use patterns)

This product is toxic to birds, fish and aquatic invertebrates. Caution should be used when applying indoors because pets may be at risk.

ENVIRONMENTAL HAZARDS (for containers of 5 gallons or 50 lbs. or more)

Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting agency has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

**ACCEPTED
with COMMENTS
EPA Letter Dated:**

DEC 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56

9 7 11

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

**IN CASE OF EMERGENCY, CALL A POISON CONTROL CENTER OR DOCTOR FOR TREATMENT ADVICE.
1-800-851-7145**

Have the product container or label with you when calling a Poison Control Center or doctor or going in for treatment.

Si no puede leer en ingles, pregunte a su supervisor sobre las instrucciones de uso apropiadas antes de trabajar con este producto.

EPA Reg. No. 70627-XX

EPA Est. No.

Lot code indicates site of manufacture.

© 2005 JohnsonDiversey, Inc., Sturtevant, WI 53177-0902 U.S.A. All Rights Reserved.

ACCEPTED
with COMMENTS
EPA Letter Dated:

DEC 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

10 7 11
ACCEPTED
with COMMENTS
EPA Letter Dated:

JohnsonDiversey



DEC 20 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide, registered under EPA Reg. No. 70627-56

OXIVIR TB REFERENCE SHEET

HOSPITAL DISINFECTANT: BACTERICIDAL in the presence of 5% serum load and 1 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces. (Hospital Disinfectant: (Bactericidal in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces. Bactericidal activity was determined by AOAC Use Dilution Test Method):

Pseudomonas aeruginosa (ATCC 15442)
Escherichia coli 0157:H7 (ATCC 35150)
Acinetobacter baumannii (ATCC 19606)

Staphylococcus aureus (ATCC 6538)
Staphylococcus aureus MRSA (ATCC 33592)

Salmonella choleraesuis (ATCC 10708)
Enterococcus faecalis VRE (ATCC 51575)

Bactericidal activity was determined by the AOAC Use Dilution Test Method.

(***VIRUCIDAL** in the presence of 5% serum load and 1 minute contact time at 20°C (68° F) on hard, non-porous environmental surfaces.)

Viruses (*Virucidal Activity) - kills on hard non-porous inanimate surfaces:

Poliovirus Type 1, Strain Brunhilde (ATCC VR-1000)

HIV-1 (AIDS Virus), Strain HTLV-III_B (HIV-1),
(Acquired from Adv. Biotechnologies, Inc.)
Influenza A/Hong Kong (ATCC VR-544)

Feline Calicivirus, Strain F9 (ATCC VR-782)

Human Coronavirus (ATCC VR-740)

Rhinovirus Type 37, Strain 151-1 (ATCC VR-1147)

Herpes Simplex Virus, Type 1 (HSV-1)
(ATCC VR-733)

Herpes Simplex Virus, Type 2 (HSV-2) (ATCC VR-734)

Hepatitis B Virus (HBV) (Acquired from
Hepadnavirus Testing)

Rotavirus WA (Acquired from University of
Ottawa)

Hepatitis C Virus (HCV) (Acquired from National Veterinary
Services Laboratories)

TUBERCULOCIDAL in the presence of a 5% serum load and 5 minute contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Mycobacterium bovis (Tb) (OT 451C150)

Tuberculocidal activity was determined by the EPA Quantitative Tuberculocidal Activity Test Method.

FUNGICIDAL This product is fungicidal against the pathogenic fungi, *Tricophyton mentagrophytes* (ATCC 9533) (the fungus which causes Athlete's Foot) in the presence of a 5% serum load and 10 minute contact time at 20°C (68°F) when used as directed on hard surfaces found in bathrooms, shower stalls, locker rooms, or other clean, non-porous, hard surfaces commonly contacted by bare feet.

Tricophyton mentagrophytes (ATCC 9533)

Fungicidal Activity was determined by the AOAC Fungicidal Activity of Disinfectants Method.

11 7 11

(Note to Agency: Text appearing in parenthesis or brackets is done to show optional text.)

Non-Food Contact Surface Sanitizer for hard non-porous surfaces – in the presence of a 5% serum load and 30 second contact time at 20°C (68°F) on hard, non-porous environmental surfaces.

Klebsiella pneumoniae (ATCC 4352)
Staphylococcus aureus (ATCC 6538)
Escherichia coli 0157:H7 (ATCC 35150)

Salmonella choleraesuis (ATCC 10708)
Staphylococcus aureus MRSA (ATCC 33592)

Pseudomonas aeruginosa (ATCC 15442)
Enterococcus faecalis VRE (ATCC 51575)

Sanitizing activity was determined by the EPA Sanitizer Test for Inanimate, Non-Food Contact Surfaces and ASTM E1153 Standard Test Method for Efficacy of Sanitizers Recommended for Inanimate Non-Food Contact Surfaces.

Malodor(s) (Activity) (Counteractancy) – eliminates (destroys) odors and odor-causing bacteria in restroom areas, behind and under sinks and counters, and storage areas (and other places where bacterial growth can cause malodors).

**ACCEPTED
With COMMENTS
EPA Letter Dated:**

DEC 20 2005

Under the Federal Insecticide,
Fungicide, and Rodenticide Act as
amended, for the pesticide,
registered under EPA Reg. No.

70627-56