

PATENTED PRODUCTION PROCESS

121101 Ekspunge-ALL first aid

EKSPUNGE™

Fungicide, Plant Nutrient & Tank Buffering Agent

For the Control of Powdery Mildew on Apples, Grapes, Cucurbits; (Cucumbers, Melons, Squash, Watermelons),
Mangoes, Stone Fruits; (Peaches, Nectarines, Plums and Cherries), Peppers, Tomatoes and Roses.

Active Ingredient: *Potassium Dihydrogen Phosphate -- 100% *Commonly known as Monopotassium Phosphate

* EPA Residue Exempt 40 CFR 180.1001 and 180.1193 * EPA REG. NO. 70644-1

SEE BACK PANEL FOR ADDITIONAL APPLICATION GUIDELINES

CROP PROTECTION WITH EKSPUNGE

ACCEPTED
MAR 11 2002
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 70644-1

*Best copy
available*

70644-1

03/11/2002

1/4

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

Precaucion al usario:
Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicada ampliamente.

**Precautionary Statement
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse

FIRST AID

If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for further treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear a long-sleeved shirt, long pants, and shoes plus socks.

User Safety Recommendations

Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Change into clean clothing.

Environmental Hazards

Do not contaminate water when disposing of equipment wash waters. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water-mark.

Directions For Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers (long sleeved shirt, long pants, shoes and socks) may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instruction and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of four (4) hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is coveralls, shoes and socks.

Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself provide all the nutrients normally required by agricultural crops.

Storage and Disposal

Storage: Store product in original container away from children and domestic animals. Do not contaminate water, food or feed by storage or disposal. **Product Disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **Container Disposal:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill via industry supported recycling programs. Incineration is not recommended.

See Side/Back Panel for Additional Precautionary Statements

Crop Protection with EksPunge

EksPunge is a soluble crystalline product to be mixed with water. Application rates vary according to the specific volumes of water applied to the crop. The lower range of spray volumes may be used as long as the crop can be completely covered with the spray solution. Surfactant use and sprayer efficiency can impact coverage. Select a water volume and corresponding rate of EksPunge necessary to thoroughly spray / mist all fruit and foliage surfaces. Gradually add the recommended amount of product to a half-filled sprayer tank and mix, then add the balance of required water while continuing to agitate the solution. Always add EksPunge to the tank mix first, then add other products after all EksPunge has been completely solubilized. **Add an approved/compatible spreader/sticker to the solution to assure complete spray coverage of plant surfaces.** Plant disease pressure can increase when plant surfaces are frequently wet and temperatures are warm. Under these severe disease conditions use the higher recommended spray rate and apply at the shorter spray interval.

NutrEcology's EksPunge suppresses existing mildew disease and inhibits further development of new mildew growth on plant tissue. It can be used alone, in alternating applications or in tank-mix spray programs with other compatible, EPA approved fungicides. It is rapidly absorbed by the plant and is mobile within the plant tissues, improving the potassium and phosphorous content in the plant. It therefore acts in a dual role as a bio-compatible fungicide for plant disease control and also as an essential plant food. EksPunge will also acidify/buffer your spray tank solution to help reduce alkaline hydrolysis of other compatible, tank mixed materials.

Best performance is attained by beginning EksPunge applications prior to the onset of disease, as a preventative disease control program. DO NOT MIX with copper fungicides or with any spray materials that warn against low pH (<5.5) applications. The active ingredient, potassium dihydrogen phosphate, when applied as directed to all crops and in accordance with good agricultural practices, is exempt from the requirement of residual tolerance as referenced in 40 CFR 180.1001 and 180.1193.

IMPORTANT

Resistant Powdery Mildew Fungus Strains May Be Present!

If treatment is not effective following use of conventional fungicides as recommended, a resistant strain of the fungus may be present. If this occurs then fungicides such as benzimidazole, thiophanate, or DMJ type may not give effective control. When resistant fungus strains are present, serious consideration should be given to the use of EksPunge for effective mildew control and crop protection. EksPunge controls mildew strains that are resistant to other fungicides and is a valuable "resistance management" tool. Best performance is attained by beginning EksPunge applications prior to the onset of disease, as a preventative disease control program. DO NOT MIX with copper fungicides or with any spray materials that warn against low pH (<5.5) solutions. The pH of a 1% aqueous solution of EksPunge is 4.5 ± 0.3.

CROP PROTECTION with eKsPunge lbs/Acre

Crop	Water Volume gal / acre					Crop	Water Volume gal / acre				
	50	100	150	200	250		50	100	150	200	250
Apples	8	8-16	17-20	21-32	33-40	Peppers	8	8-16	17-20	20	20
Cucurbits	10	10-20	17-20	20	20	Roses	5	6-8	9-10	10	10
Grapes	8	8-16	17-20	21-32	33-40	Stone Fruits	8	8-16	17-20	20	20
Mangoes	8	8-16	17-20	21-32	33-40	Tomatoes	8	8-16	17-20	20	20
Ornamentals	8-13	13-25	19-38	25-50	32-63	Turf	8-13	13-25	19-38	25-50	32-63

Conditions of Sale
1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than the officer of Sales, is authorized to make any warranty, guarantee or claim concerning this product.
2. Because the time, place, rate of application and other conditions of use are beyond Seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

NET WEIGHT 50 POUNDS (22.68 KG)



Distributed and Guaranteed by:
LidoChem, Inc.
20 Village Court - Hazlet, NJ 07730
Phone: 732.686.8000 - Fax: 732.264.2761

Product of Israel
LidoChem, Inc. Logo and EksPunge™ are trademarks of LidoChem, Inc.
Manufactured and Packaged for LidoChem, Inc.

0-50-32

- EPA REG NO 70644 I
- EPA EST NO. 70644 NJ I
- EPA EST NO. 67536-FL-001
- EPA EST NO. 14322-NY-1
- EPA EST NO. 2935-CA-1
- EPA EST NO. 66196-CA-1

Guaranteed Analysis: Available Phosphate (P₂O₅).....50 %
 Soluble Potash (K₂O).....32 %
 Derived from: Monopotassium Phosphate

Product Description

Ekspunge is manufactured specifically as a low salt, water soluble, foliar and special application fungicide and plant nutrient. Its use is suggested as a supplement to a growers standard practice fungicide and fertilizer programs. The target is reduced pesticide use and enhanced yield and quality. Ekspunge is a highly soluble, low salt index formulation developed to supplement standard fertility practices by providing a highly available source of phosphorous and potassium.

Research has shown that foliar-applied nutrients, in a pure and soluble form, are absorbed more efficiently by foliage than are those supplied in the soil. Nutrient translocation to all parts of the plant is generally more rapid when nutrients are applied foliarly.

A good tissue testing program may be helpful to monitor and maintain optimum plant growth and development. Adverse conditions such as moisture, stress, weather, salts, soil type, etc., may induce nutrient deficiency symptoms. Ekspunge application is a means of obtaining a quick response to needed nutrients when applied as directed.

Salt Index: 8.4 (.097 per 1% of plant nutrient)
 pH (1% solution): 4.5 ± 0.3

Mixing Directions

- Add approximately 1/2 water to tank before gradually adding Ekspunge. Agitate thoroughly while adding Ekspunge and the remaining water. When tank mixing add pesticide last.
- When temperatures are cold, allow extra time for this product to completely dissolve.
- Research has demonstrated enhanced uptake with the addition of a surfactant. Do not use with surfactants when plants are under severe stress conditions, such as heat or water stress. Applications should begin immediately after adverse stress conditions subside.
- Ekspunge should normally be mixed at 1lb. or less per gallon of water. Concentrations up to 1.9 lbs. per gallon are possible, but consideration must be given to the pH of the solution and that storability of high concentration mixes in temperatures less than 60 degrees Fahrenheit is not recommended.

Compatibility with Product Combinations

Ekspunge™ is compatible with most pesticides and liquid fertilizers. Ekspunge™ may be applied in an alternating tank mix program. **WARNING:** Ekspunge™ acts as a buffer resulting in an acidic spray solution. Combination with copper fungicides or with any spray materials that warn against low pH (<5.5) solutions is not recommended. Always test tank mixes for compatibility, via a jar test, before mixing large batches. Ekspunge™ combined with labeled rates of Prudent, or Prudent Plus liquid fertilizers, may be used with local crop protection practices.

EKSPUNGE Guidelines for Nutritional Application

Crop	Rate	Timing
Alfalfa	Foliar: 5-8 lbs/A per application. Can be applied after cuttings at a maximum concentration of 1.9 lbs/A if label mixing directions are followed. Fertigation: Apply 40 to 50 lbs/A	Foliar: Apply at first re-growth - when alfalfa is 6-8" tall; apply after each cutting. Fertigation: Apply one week after every cut through irrigation.
Almonds	Foliar: Pre-bloom: 5-10 lbs/A; Finish: 5-15 lbs/A Use a maximum of 1.5 lbs of product per 10 gallons of spray solution by ground rig or a maximum of 4 lbs of product per 10 gallons of spray solution by air.	Apply pre-bloom 1 week before to 1 week after peak bloom. Apply as a finish spray 1-2 times from 3 weeks to 10 days before harvest.
Apples	10 - 20 lbs/A per application. Use a maximum of 1.5 lbs of product per 10 gallons of spray solution by ground rig or a maximum of 4 lbs of product per 10 gallons of spray solution by air.	Mid-Season Sprays: Apply during June/July, up to 4 successive sprays 7 to 10 days apart. Finish Spray: Apply at the color break period. Post Harvest Spray: Apply immediately after harvest.
Avocado	25-40 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply 2-3 times from fruit set until 30 days before harvest every 30 days.
Banana	20-30 lbs/A per application. Use a maximum of 2 lbs of product per 10 gallons of spray solution.	Apply 1-2 times - 15 and 21 days after shooting. Apply 1 time 21-30 days before bloom.
Beans - Dry, Succulent, Limas	5-8 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply at first flower and 2 additional applications during the main filling stage of pod development 7 to 10 days apart.
Berries: Bush Type	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	2-4 foliar applications starting at first flower at 14 - 21 days intervals.
Citrus	20-25 lbs/A per application. Use a maximum of 4 lbs of product per 10 gallons of spray solution.	Apply up to 3 times: Pre-bloom, late June (after June drop), and in early September.
Cool Season Turf grass	2-4 ozs. per 1000 sq. ft. (6-11 lbs/A) per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution	Apply every 7 to 14 days throughout the season. Higher rates can be used in fertigation systems where EKSPUNGE is the primary source of P & K.
Corn, Sweet	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	2 applications: Apply 2 weeks prior to tasseling and again between tasseling and silking.
Cotton	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution by ground rig and a maximum of 10 lbs of product per 10 gallons of spray solution by air.	Make applications at 30 (square development), 60 (first flowering), and 90 (boll set) days after emergence.
Deciduous Fruits - apples, pears, cherries, apricots, peaches, plums, and nectarines	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply as a pre-bloom and post-bloom spray
Grapes	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply starting at the 3 to 5 cm shoot stage through veraison every 2 to 4 weeks.
Hops	5-10 lbs/A per application. Use sufficient water for complete coverage by ground sprayers. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Begin applications at early season training and continue through end of bloom period as often as every 7 days.
Legumes	5-8 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply at first flower with 2 additional applications during the mid filling stage of pod development 7 to 10 days apart.
Lemons	8-10 lbs/A per application. Use a maximum of 4 lbs of product per 10 gallons of spray solution.	Make 2 applications - 1 after fruit setting and the second 1 month later.
Mango	15 - 20 lbs/A per application. Use a maximum of 4 lbs of product per 10 gallons of spray solution.	Apply up to 3 times after Panicle development every 14 days.
Melons, Pumpkins, Cucurbits	8 - 12 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply 2-4 sprays beginning at fruit set on a 7 to 14 day interval.
Onion	8-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	2-4 applications beginning at transplanting. Repeat applications every 30 days.
Ornamental	Use 1 lb in 10 gallons of water and spray to wet. Transplants - Use 1-2 lbs of product per 10 gallons and apply approximately 1/2 gallon per transplanted shrub/tree.	Apply at bloom, spring shoot push or shortly after transplant and repeat in 14-21 days. Use any time new growth is pushing or in conjunction with pesticide applications.
Peanuts	5-8 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	3 applications - 1 at early bloom with 2 additional sprays at 60 days after planting and 10 days later.
Peppers and Tomatoes	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply 2-6 sprays every 14 days starting at first bloom.
Potato	5-10 lbs/A per application. Use a maximum of 5 lbs of product per 10 gallons of spray solution.	Apply at early initial tuber formation. Subsequent sprays can be made with fungicide applications.
Produce/Lettuce Cole Crops	2-4 lbs/A per application. Use a maximum of 1 lbs of product per 10 gallons of spray solution.	Use multiple low rate applications 10-14 days apart starting just after transplant. Use as a pre harvest application from 3-14 days before harvest to improve color.
Rice	3-6 lbs/A per application. Use a maximum of 1 lb of product per 10 gallons of spray solution by ground rig or a maximum of 5 lbs of product per 10 gallons of spray solution by air.	Spray two times at the end of tillering and at Panicle initiation.
Root Crops	2-8 lbs/A per application. Use a maximum of 2 lbs of product per 10 gallons of spray solution.	Apply at increasing rates every 14-21 days from early root swell until 2 weeks before harvest.
Small Grains	8-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution by ground rig or a maximum of 8 lbs of product per 10 gallons of spray solution by air.	Apply at late anthesis stage.
Soybean	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution by ground rig or a maximum of 8 lbs of product per 10 gallons of spray solution by air.	2 applications - one at the early bloom stage and second at the main pod filling stage.
Strawberry	5-10 lbs/A per application. Use a maximum of 3 lbs per 10 gallons of spray solution.	2-4 applications during the harvest period on a 7 to 14 day schedule. Can also be applied as needed through fertigation.
Sugarbeet	5-10 lbs/A per application. Use a maximum of 3 lbs of product per 10 gallons of spray solution by ground rig or a maximum of 8 lbs of product per 10 gallons of spray solution by air.	Apply when leaves are 10" across and again 3-4 weeks later. Apply again 4 weeks before harvest.
Warm Season Grass	2-4 ozs per 1000 sq. ft. (6-11 lbs/A) per application. If used as a starter, apply 4-6 ozs per 1000 sq. ft. (11-16 lbs/A) Use a maximum of 3 lbs of product per 10 gallons of spray solution.	Apply every 7 to 14 days throughout the season. Can be used as a starter fertilizer during transition periods to cool season grasses as a starter. Higher rates can be used in fertigation systems where EKSPUNGE is the primary source of P & K.

Ekspunge Crop Protection Fungicide Application Guidelines

APPLES

For control of powdery mildew (*Podosphaera leucotricha*) on Apples use 8 to 40 lbs of EKSPUNGE per acre. Start spraying at tight cluster and continue spraying every 7 to 10 days until terminal shoots cease their vegetative growth. The rate of product per acre will vary depending upon the tree size (canopy development) and the volume of water.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 40 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

CUCURBITS

(CUCUMBER, MELONS, KORMARH, WATERMELONS)

For control of powdery mildew (*Sphaerotheca fuliginea*) on the above listed cucurbits use 10 to 20 lbs of EKSPUNGE per acre. Start spraying when plants begin to run or when disease pressure is anticipated. Repeat at 7 to 14 day intervals as needed. Under conditions of severe disease pressure, use the higher rate and apply at 7 day intervals. For best results, avoid application when temperatures are over 85°F and humidity is high. For greenhouse use, shading may be required.

MI: 10 lbs / 50 gallons spray solution per acre.
 Max: 20 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 20 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

GRAPES

For control of powdery mildew (*Uncinula necator*) on grapes use 8 to 40 lbs of EKSPUNGE per acre. Start spraying in the spring when shoots are 3 to 5 inches in length and when disease pressure is anticipated. Repeat every 10 to 14 days. The low per acre rates are intended for early in the season when disease pressure is low. The per acre rate should be increased as disease pressure increases. For improved appearance on table grapes, use lower application rates.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 40 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

MANGOES

For control of powdery mildew (*Oidium mangiferae*) on mango use 8 to 40 lbs of EKSPUNGE per acre. Start spraying at first appearance of bloom panicles (approximately 2 inches long) and repeat at 7 to 14 day intervals until all fruit are set. If additional sprays are required, continue at 2 to 3 week intervals until shoot growth ceases - which should be about 6 sprays.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 40 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

ORNAMENTALS

For control of powdery mildew, including but not limited to *Microspheerium alni* and *Erysiphe cichoracearum* on woody and herbaceous ornamentals, use 8 to 40 lbs of EKSPUNGE per acre. Start spraying in early Spring when conditions become favorable for disease development (i.e. cool, humid, cloudy periods) and continue spraying on a 7 to 14 day schedule for the entire season.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 40 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

PEPPERS

For control of powdery mildew (*Leveillula taurica*) on peppers.

Greenhouse Grown: Mix 10 lbs per 100 gallons and apply 1.5 gallons of finished spray per 1,000 sq. ft. at 5 to 7 day intervals. Shading may be needed to reduce temperatures during spraying.

Field Grown: Use 8 to 20 lbs of EKSPUNGE per acre when disease pressure begins to increase. Repeat at 7 to 10 day intervals.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 20 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

ROSES

For control of powdery mildew (*Sphaerotheca pannosa* var. *roseae*), use 5 to 10 lbs EKSPUNGE per acre. Apply at 5 to 7 day intervals as needed. Best performance can be achieved with full wetting of leaves without runoff.

MI: 5 lbs / 50 gallons spray solution per acre.
 Max: 10 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 8 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

STONE FRUITS

(PEACHES, NECTARINES, PLUMS, CHERRIES)

For control of powdery mildew (*Sphaerotheca pannosa* var. *persicae* and *Podosphaera oxycanthae*) on stone fruits as listed use 8 to 20 lbs of EKSPUNGE per acre. Follow local recommendations for powdery mildew control timings or apply when disease pressure is anticipated and repeat every 7 to 14 days.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 20 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

TOMATOES

For control of powdery mildew (*Leveillula taurica*) on tomatoes.

Greenhouse Grown: Use 10 lbs per 100 gallons and apply 1.5 gallons of finished spray per 1,000 sq. ft. at 5 to 7 day intervals. Shading may be needed to reduce temperatures during spraying.

Field Grown: Use 8 to 20 lbs of EKSPUNGE per acre when disease pressure begins to increase. Repeat at 7 to 10 day intervals.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 20 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.

TURFGRASS

For control of powdery mildew (*Erysiphe graminis* D.C.), use 8 to 40 lbs of Ekspunge per acre. Start spraying in early Spring when conditions become favorable for disease development (i.e. cool, humid, cloudy periods) and continue spraying on a 7 to 14 day schedule for the entire season.

MI: 8 lbs / 50 gallons spray solution per acre.
 Max: 20 lbs / 250 gallons spray solution per acre.
 DO NOT exceed 16 lbs per 100 gallons of finished spray solution.
 * See Table for additional rates per water volume.



Distributed and Guaranteed by:
 LidoChem, Inc.
 20 Village Court - Hazlet, NJ 07730
 Phone 212.898.8000 / Fax 212.264.2751

LidoChem, Inc. Logo and Ekspunge™ are trademarks of LidoChem, Inc.
 Manufactured and Packaged for LidoChem, Inc.

121101 Etekspunge-ALL final.d

