


70810-1

09/13/1999

111

### Supplemental Label: AuxiGro™ WP Plant Metabolic Primer

Crops covered by this label: Onions



**FOR USE ONLY IN THE FOLLOWING STATES: Texas and Arizona.**

**Expiration date of this label: 12/31/99**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

EPA Reg. No. 70810-1

EPA Est. No. 33967-NJ-1, 39578-TX-1

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION - PRECAUCION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

Refer to AuxiGro package (bag) for ingredient information, precautionary statements and information concerning storage and disposal. Please see package insert for Agricultural Use Requirements.

**PRECAUTIONARY STATEMENTS, STATEMENTS OF PRACTICAL TREATMENT AND ENVIRONMENTAL HAZARDS**  
STATEMENTS are provided on the package label.

**MIXING INSTRUCTIONS** for tank mix combinations, compatibility testing and mixing of AuxiGro are provided on the product package insert for Agricultural Use Requirements.

**APPLICATION INSTRUCTIONS.** Applications of AuxiGro can effectively enhance plant development, growth of numerous plant species and induce plant resistance to control or suppress selected plant pathogens. Applications of AuxiGro may begin at seeding (or transplanting) and continue until harvest. For optimal yield/disease response, apply AuxiGro postemergence according to the application timings and rates provided in the table below. Multiple applications (with fungicides) may be made at a rate of 1 - 8 ounces per acre per application. Total seasonal applications should not exceed 24 ounces per acre per crop.

All rates of application for AuxiGro are given as broadcast rates and expressed as ounces of formulated product per acre per application. If crop growth will permit a band treatment, use proportionally less product per acre per application than the broadcast rate. AuxiGro should be applied with silicone-based surfactants unless tank mix combinations prohibit their use or crop injury is likely to occur.

Allow heavy dew to evaporate prior to AuxiGro applications. Do not apply AuxiGro if windy conditions exist or a strong threat of rain exists within 4 hours of application.

APPLICATION TIMING	Rate/Acre/ Application
<b>Onions.</b> <b>First application:</b> Apply 4 oz. of AuxiGro at the beginning of bulbing. <b>Second application:</b> Apply 4 oz. of AuxiGro at approximately 4 weeks prior to harvest, a minimum of 21 days after the initial application.	4 ounces

Auxein Corporation  
3125 Sovereign Dr., Ste. B  
Lansing, MI 49811-4240  
1-888-8AUXEIN (1-888-828-9346)