

Revision 021011


70810-1

6/19/2003

Page 1 of 14

AuxiGro® WP Plant Metabolic Primer
WETTABLE POWDER
FOR INCREASED YIELD AND QUALITY
MASTER LABEL

EPA Registration Number 70810 - 1
 EPA Establishment Number 33967-NJ-1, 39578-TX-1



Active Ingredients:

GABA: gamma aminobutyric acid -----	29.2 %
L-Glutamic acid -----	29.2 %
Other Ingredients -----	41.6 %
TOTAL -----	100.0 %

(1 pound contains 0.6 pounds of active ingredient)

(5 pounds contains 3 pounds of active ingredient)

KEEP OUT OF REACH OF CHILDREN

CAUTION!;PRECAUCIÓN!

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
 (If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS - CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush eyes with plenty of water. Call physician if irritation persists.

Personal Protective Equipment (PPE). Applicators and other handlers must wear:

- long sleeve shirt and long pants
- shoes plus socks
- protective eyewear
- waterproof gloves

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if product gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

DO NOT apply directly to water or areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Please see package insert on bottom of bag for complete Directions For Use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notifications and emergency assistance. It also contains specific instructions and exceptions pertaining to the restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Re-Entry Statement:

Do not enter or allow worker entry into treated areas without protective clothing during the restricted-entry interval (REI) of 4 hours.

For Early Entry

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated (plants, soil) wear:

- ◆ Long sleeve shirt and long pants
- ◆ Shoes plus socks
- ◆ Protective eyewear
- ◆ Coveralls

Observe all cautions and limitations on this label and on the labels of products used in combination with Auxigro. Do not use Auxigro other than in accordance with the instructions set forth on this label. The use of Auxigro not consistent with this label may result in injury to crops.

GENERAL INFORMATION

Auxigro can be applied on all labeled crops per label instructions (Directions for Use). When used as directed, Auxigro has been shown to increase yields of crops. Auxigro applications have also resulted in increased size and/or quality of treated commodities, early ripening in certain vegetables, increased root growth, early flowering and fruit set, faster seed germination and rooting, and induction of plant resistance to control or suppress certain plant pathogens.

MIXING INSTRUCTIONS

Mixing of Auxigro: Fill the spray tank one-half to three-quarters full with clean water. Use a calibrated device to weigh the required amount of Auxigro. Add Auxigro to the spray tank while agitating. Add other products, adjuvants or surfactants and then fill the remainder of the tank with water.

Tank mix combinations with Auxigro: When Auxigro is used in combination with other products, refer to the respective product labels for rates and methods of application; proper timing; disease, weeds or insects controlled; restrictions and precautions. Always use in accordance with the most restrictive label restrictions and precautions. Do not exceed any label dosage.

Spray solution pH: For best results, final spray solution should be near neutral or slightly acidic (pH 6.0 to 7.5).

Compatibility Testing: Auxigro is compatible with most commonly used agricultural fungicides, insecticides, micronutrients and spray adjuvants. If compatibility with another product is unknown, a jar test for physical compatibility should be conducted before field mixtures are made. Read and recognize all cautions and limitations on labeling of all products used in mixtures.

Adjuvants and Surfactants: Optimal results are obtained when a silicone-based surfactant is used with Auxigro. The surfactant should be used at the lowest label rate that is recommended for use with Plant Growth Regulators (PGRs) or foliar sprays (typically 4 to 6 ounces surfactant per 100 gallons). If Auxigro is tank mixed with other products which prohibit surfactant use, follow the tank mix product label and do not use a surfactant.

Mixing containers, hoses and spraying equipment should be cleaned or triple rinsed prior to mixing Auxigro for application.

APPLICATION INSTRUCTIONS

Applications of Auxigro can effectively enhance plant development, growth of numerous plant species and induce plant resistance to control or suppress selected plant pathogens. Applications of Auxigro may begin at seeding (or transplanting) and continue until harvest. For optimal yield/disease response, apply Auxigro postemergence according to the application timings and rates provided in the table below. Multiple applications (with fungicides) may be made at a rate of 2 - 4 ounces (per application). Total season application should not exceed 24 ounces per acre per crop.

All rates of application for Auxigro are given as broadcast rates and expressed as ounces of formulated product per acre per application. If crop growth will permit a band treatment, use proportionally less product than the broadcast rate. **Auxigro should be applied with silicone-based surfactants unless tank mix combinations prohibit their use or crop injury is likely to occur.**

Allow heavy dew to evaporate prior to Auxigro applications. Do not apply Auxigro if windy

conditions exist or a strong threat of rain exists within 4 hours of application.

Ground Application: Apply uniformly with properly calibrated ground equipment in 20 to 100 gallons of water per acre, except as noted in the Directions for Use. Application volume and pressure may be increased, depending on the size of plants being treated. Hollow cone nozzles are recommended; however, other types of nozzles that achieve uniform wetting of the entire plant canopy are acceptable.

Aerial Application: Thorough spray coverage is required for maximum benefit. To achieve good coverage, use proper droplet size, spray uniformly, and ensure swath width and spray volume are correct. Do not use less than 7 gallons total spray volume per acre.

Overhead Sprinkler Irrigation (For Potatoes Only. Use typical surfactants or additives that are recommended for overhead irrigation. Do not use a silicone based surfactant when applying Auxigro by overhead irrigation): Auxigro can be applied through common overhead sprinkler systems, such as center-point, lateral move, end tow, side (wheel) roll, traveler and others. All equipment must be properly calibrated for pesticide application before use.

Center-pivot, Lateral Move, End Tow, and Traveler Irrigation Equipment. Apply Auxigro with no more than 1/4 inches of water (6750 gallons of water) per acre when the system and injector are operated at normal pressures recommended by their manufacturer.

Solid Set, Side (wheel) Roll and Hand Move Irrigation Equipment. Inject Auxigro at the end of the irrigation cycle or as a separate application to maximize the foliar retention.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The chemical injection pipeline must also contain a functional, normally closed solenoid operated valve located on the intake side of the injection pump and connected to a system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the chemical injection pump when the water pump motor stops. The irrigation line or water pump must include functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where chemical distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with chemicals and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Lack of effectiveness may result from non-uniform distribution of treated water.

Allow sufficient time for chemical to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for chemical application to a public water system unless the label-prescribed safety devices for drawing water supplies are in place.

Crops & Application(s)	Rate / Acre / Application & Timing*
Almond	To increase total yield. First application: Apply 4 oz. of Auxigro when trees are in pink bud (< 20% bloom) Second application: Apply 4 oz. of Auxigro at full bloom Third application: Apply 4 oz. of Auxigro at petal fall
Broccoli and Cauliflower	To increase total yield. First application: Apply 4 oz. of Auxigro 3 weeks prior to "button" (head) formation Second application: Apply 4 oz. of Auxigro at button initiation.
Celery	To increase the size and/or total yield. First application: Apply 4 oz. of Auxigro when plants are 6 – 8 inches tall. Second application: Apply 4 oz. of Auxigro when plants are 14 – 16 inches tall, approximately 14 – 21 days after the initial application, no sooner than 12 days after the initial application.
Cotton	To increase total lint and/or seed yield. Apply 4 oz. of Auxigro at early boll formation.
Edible Dry Beans	To enhance yields One application: Apply 4 oz. of Auxigro to the crop at flower bud initiation (less than 5% bloom) OR at initial pod fill stage of growth (approximately 21 days after first bud formation).

<p>Cucumbers (fresh market).</p> <p>First application:</p> <p>Second application:</p> <p>Other applications:</p>	<p>To increase the grade and/or marketable yield.</p> <p>Apply 4 oz. of AuxIGro at early bloom, just prior to running.</p> <p>Apply 4 oz. of AuxIGro at 10-14 days after the initial application.</p> <p>Apply 4 oz. of AuxIGro at 14-16 day intervals when multiple harvests are taken. Total season application should not exceed 24 oz. per acre per crop.</p>
<p>Grapes (Disease Control).</p> <p>Powdery Mildew (<i>Uncinula sp.</i>) and Bunch Rot (<i>Botrytis cinerea</i>).</p> <p>Application timing:</p>	<p>AuxIGro will control Powdery Mildew and provide suppression of Bunch Rot. To do so it must be used as part of a disease management program. AuxIGro can be used alone if alternated with standard commercial fungicides on an established spray schedule, or can be applied in combination with commercial fungicides on an established spray schedule.</p> <p>Apply 2 to 4 oz. of AuxIGro beginning at pre-bloom (6 – 12 inch shoots) and do not extend applications beyond a 16 day interval. Total season application should not exceed 24 oz. per acre per crop.</p> <p>Use the higher application rates when AuxIGro is applied alone, on highly susceptible varieties or under heavy disease pressure.</p> <p>AuxIGro may be applied alone or in tank mix combinations with recommended rates of fungicide, insecticide, and/or miticide.</p> <p>Thorough coverage is essential for good disease control. AuxIGro should be uniformly applied to the crop canopy. For ground applications use a minimum of 50 gallons per acre for conventional spray equipment, and a minimum of 36 gallons per acre for electrostatic sprayers.</p>
<p>Grapes</p> <p>First Application:</p> <p>Second Application:</p>	<p>To increase Soluble Sugars as measured by degrees Brix:</p> <p>Apply 4 oz. of AuxIGro at full bloom (beginning of berry development).</p> <p>Apply 4 oz. of AuxIGro at veraison (berry softening) prior to grapes achieving 18 degrees Brix and within 60 days of anticipated harvest.</p>
<p>Grass grown for Seed (annual rye, perennial rye, Kentucky bluegrass, tall fescue)</p>	<p>To increase total yield</p> <p>Apply 4 oz. of AuxIGro at early seed head fill (post anthesis)</p>

<p>Options for: Tall fescue Perennial rye (established only)</p>	<p>Apply 4 oz. of Auxigro in the fall at the 2 to 4 leaf stage. Apply 4 oz. of Auxigro at beginning of spring re-growth (greening up)</p>
<p>Onions First Application Second Application:</p>	<p>To increase the marketable grade and/or yield Apply 4 oz. of Auxigro at bulb swell (typically 1 ¼ to 1 ½ inch diameter) Apply 4 oz. of Auxigro at approximately 4 weeks prior to harvest, a minimum of 21 days after the initial application.</p>
<p>Peppers (bell, green, jalapeño). First application: Second application:</p>	<p>To increase the marketable grade and/or total yield. Apply 4 oz. of Auxigro at flower bud initiation (pre-bloom of main canopy). Apply 4 oz. of Auxigro after first fruit set, no sooner than 14 days after initial application.</p>
<p>Head lettuce (iceberg). First application: Second application:</p>	<p>To increase the marketable yield and/or provide earlier maturity. Apply 8 oz. of Auxigro immediately after thinning (4 – 6 true leaves). Apply 8 oz. of Auxigro at first sign of head cupping, no sooner than 14 days after first application.</p>
<p>Leaf lettuce (romaine, butter). First application: Second application:</p>	<p>To increase the marketable yield and/or provide earlier maturity. Apply 8 oz. of Auxigro immediately after thinning (4 – 6 true leaves). Apply 8 oz. of Auxigro at first sign of head cupping, no sooner than 14 days after first application.</p>
<p>Peanuts. First application: Second application:</p>	<p>To increase the total yield. Apply 4 oz. of Auxigro at early bloom, 18 – 20 trifoliates (prior to running). Apply 4 oz. of Auxigro at full bloom, approximately 25–30 days after initial application.</p>

<p>Potatoes.</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the marketable yield and/or improve tuber grades.</p> <p>Apply 5 oz. of Auxigro when the first tubers are approximately the size of a small pea (plants 6 – 10 inches in height, typically 2 -3 weeks after emergence).</p> <p>Apply 5 oz. of Auxigro 21-28 days after first application, OR at 30 days prior to harvest.</p>
<p>Potato Seed Pieces</p>	<p>To enhance emergence, vigor and/or increased tuber set.</p> <p>Apply 0.1 oz Auxigro per 100 pounds (hundredweight) of seed.</p>
<p>Snap Beans (<i>Phaseolus spp</i>) (Southeast U.S. recommendation only).</p> <p>One application:</p>	<p>To increase the total yield.</p> <p>Apply 8 oz. of Auxigro at pre-bloom (bud formation).</p>
<p>Stone Fruit (including almonds, apricots, cherries, plums, nectarines, peaches and prunes)</p> <p>First application:</p> <p>Second application:</p> <p>Third application:</p>	<p>For the control of brown rot (<i>Monilinia laxa</i>) and suppression of shot hole (<i>Stigmia carpophila</i>)</p> <p>Apply 4 ounces of Auxigro when trees are in pink bud (<20% bloom)</p> <p>Apply 4 ounces of Auxigro at full bloom</p> <p>Apply 4 ounces of Auxigro at petal fall</p>
<p>Strawberries.</p> <p>Ever Bearing Varieties.</p> <p>First application:</p> <p>Later applications:</p> <p>June Bearing Varieties.</p> <p>First Application:</p> <p>Second Application:</p>	<p>To increase the total yield and/or quality.</p> <p>(harvest continues several months)</p> <p>Apply 8 oz. of Auxigro at any time after first harvest.</p> <p>Additional 8 oz. applications of Auxigro should be made at 1-month intervals following the first application. Total season application should not exceed 24 oz. per acre per crop.</p> <p>(harvest duration is typically less than 3 months)</p> <p>Apply 8 oz. of Auxigro 5 - 7 days prior to first harvest.</p> <p>Apply 8 oz. of Auxigro 14 – 21 days after initial application.</p>

<p>Sugarbeets</p> <p>One application</p>	<p>To increase total sugar production.</p> <p>Apply 4 oz AuxiGro to crop 30 to 45 days before harvest</p>
<p>Tomatoes (processing).</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the marketable yield and to maintain or improve the soluble solids levels as measured by degrees Brix.</p> <p>Apply 4 oz. of AuxiGro at full bloom in the canopy (50-70% bloom).</p> <p>Apply 4 oz. of AuxiGro at first fruit set 10 –21 days after first application.</p>
<p>Tomatoes (fresh).</p> <p>First application:</p> <p>Second application:</p> <p>Multiple applications:</p>	<p>To increase the marketable yield and/or improve grades.</p> <p>Apply 4 oz. of AuxiGro at pre-bloom (bud formation for initial production set).</p> <p>Apply 4 oz. of AuxiGro at first fruit set, at least 21 days after first application.</p> <p>Additional 4 oz. applications may be made after the second application at intervals of at least 21 days, if multiple harvests over an extended period are taken. Total season application should not exceed 24 oz. per acre per crop.</p>
<p>Watermelons.</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the yield and/or quality.</p> <p>Apply 4 oz. of AuxiGro at early female bloom (less than 10%), just prior to running.</p> <p>Apply 4 oz. of AuxiGro at first fruit set, typically 14 – 21 days following the initial application.</p>
<p>Watermelons.</p> <p>Multiple applications:</p>	<p>For suppression of gummy stem blight (<i>Didymella bryoniae</i>).</p> <p>AuxiGro may be applied in combination with commercial fungicides at 2 oz. of AuxiGro from emergence until harvest on a regular basis (7-14 day intervals) for suppression of gummy stem blight. Total season application should not exceed 24 oz. per acre per crop.</p>

Crop Group 1: Root and Tubers (Rates and directions apply to root and tuber crops other than potatoes and carrots)	To increase the total yield and/or quality of selected root and tuber crops:
First application:	Apply 4-8 oz of Auxigro at first tuber initiation or the beginning of bulking up as appropriate.
Second application:	Apply 4-8 oz of Auxigro 21-28 days after initial application or approximately 30 days prior to harvest.
Crop Group 3: Bulb Vegetables	To increase the size and/or total yield of onions and garlic:
First application:	Apply 4-8 oz of Auxigro at initial bulb swell (bulbing).
Second application:	Apply 4-8 oz of Auxigro at 4 weeks after initial application, at least 30 days prior to harvest."
Crop Group 4: Leafy Vegetables (Rates apply to Leafy Vegetables crops other than head lettuce, leaf lettuce and spinach)	To increase the size and/or total yield of celery and other leafy vegetables.
First application:	Apply 4-8 oz of Auxigro at 1-5 days after thinning or transplant.
Second application:	Apply 4-8 oz of Auxigro 14 – 30 days after first application
Or	
Multiple applications:	Apply 1-2 oz of Auxigro every 7 days.
Crop Group 5: Cole crops	To increase the total yield and/or marketable grade.
First application:	Apply 4 – 8 oz. Auxigro at the 4-8 leaf stage (3 to 4 weeks prior to head formation)
Second application:	Apply 4 – 8 oz. Auxigro just prior to head formation, no sooner than 14 days after first application.
Crop Group 6: Legume Vegetables	To increase the total yield and/or quality of various legume vegetables:
First application:	Apply 4 – 8 oz. Auxigro at flower bud initiation.
Second application:	Apply 4 – 8 oz Auxigro at at early pod formation.

<p>Crop Group 8: Fruiting Vegetables (Separate use instructions are provided for tomatoes and peppers.)</p> <p>First application: Second application:</p>	<p>To increase the yield of various fruiting vegetables:</p> <p>Apply 4 – 8 oz. Auxigro at bud set (initiation) in canopy. Apply 4 – 8 oz. Auxigro at full bloom, no sooner than 10 – 14 days after initial application.</p>
<p>Crop Group 9: Cucurbit vegetables (including melons)</p> <p>First application: Second application:</p>	<p>To increase the total yield and/or quality of various cucurbits:</p> <p>Apply 4 – 8 oz. Auxigro at early female bloom, prior to running. Apply 4-8 oz. Auxigro 14 – 21 days after first application.</p>
<p>Crop Group 10: Citrus Fruits</p> <p>First application: Second application: Third Application: Fourth Application:</p>	<p>To increase the growth and development or the total yield and/or quality.</p> <p>Apply 2-8 oz Auxigro per 100 gal. at bud formation Apply 2-8 oz Auxigro per 100 gal. at full bloom Apply 2-8 oz Auxigro per 100 gal. at petal fall Apply 2-8 oz Auxigro per 100 gal. as fruit begins to size (bulk) up</p>
<p>Crop Group 11: Pome Fruits</p> <p>Bloom time applications: Nutrient Combinations:</p>	<p>To increase the growth and development as well as the total yield and/or quality.</p> <p>Apply 2-8 oz /100 gal. Of Auxigro at each of the following growth stages - Pink bud, full bloom & petal fall.</p> <p>Apply 2-8 oz Auxigro per 100 gal. as a tankmix partner with foliar micro-nutrients during fruit development.</p>
<p>Crop Group 12: Stone Fruits</p> <p>Bloom time applications: Nutrient Combinations:</p>	<p>To increase the growth and development or the total yield and/or quality the fruits:</p> <p>Apply 2-8 oz /100 gal. Of Auxigro at each of the following growth stages - Pink bud, full bloom & petal fall.</p> <p>Apply 2-8 oz Auxigro per 100 gal. as a tankmix partner with foliar micro-nutrients during fruit development.</p>
<p>Crop Group 13: Berries Group</p> <p>First application: Second application:</p>	<p>To increase the total yield and/or quality of berries:</p> <p>Apply 2-4 oz /100 gal. of Auxigro at first flower or fruit set. Apply 2-4 oz /100 gal. of Auxigro 4 – 8 weeks after first application.</p>

<p>Crop Group 14: Tree Nuts Group</p>	<p>To increase total yield and/or quality and the Nut Trees:</p>
<p>Bloom time applications:</p>	<p>Apply 2-8 oz /100 gal. Of Auxigro at each of the following growth stages - Pink bud, full bloom & petal fall as well as bulking up.</p>
<p>Nutrient Combinations:</p>	<p>Apply 2-8 oz Auxigro per 100 gal. as a tankmix partner with foliar micro-nutrients during fruit development.</p>
<p>Crop Group 15: Cereal grains group</p>	<p>To increase the total yield and or quality.</p>
<p>Small Grains: (i.e., wheat, oats, rice, etc.) Corn:</p>	<p>Apply 2 – 4 oz. Auxigro at initial seed head fill. Apply 2 – 4 oz. Auxigro at 20 to 80% tasseling</p>
<p>Crop Group 16: Forage, fodder and straw of cereal grains group.</p>	<p>To increase the yield and or quality of various forage crops:</p>
<p>First application:</p>	<p>Apply 2-4 oz Auxigro prior to tillering on small grains or at V-4 to V-6 growth stage for corn and sorghum.</p>
<p>Second application:</p>	<p>Apply 2-4 oz Auxigro at boot stage for small grains and at V-10 to V12 growth stage for corn and sorghum.</p>
<p>Crop Group 17: Grass forage, fodder, and hay.</p>	<p>To increase the yield and or quality of various forage crops:</p>
<p>First application:</p>	<p>Apply 2-4 oz Auxigro when 2-4 inches of new growth appears.</p>
<p>Second application:</p>	<p>Apply 2-4 oz Auxigro 4 weeks after first application.</p>
<p>Crop Group 18: Non-grass animal feeds</p>	<p>To increase the yield and or quality of the forage.</p>
<p>First application:</p>	<p>Apply 2-4 oz Auxigro when 2-4 inches of new growth appears.</p>
<p>Second application:</p>	<p>Apply 2-4 oz Auxigro when 2-4 inches of re-growth appears after cutting.</p>
<p>Crop Group 19: Herbs and Spices</p>	<p>To increase the yield and/or quality of various herb and spice crops:</p>
<p>First application:</p>	<p>Apply 2-4 oz Auxigro when 2-4 inches of new vegetative growth appears.</p>
<p>Second application:</p>	<p>Apply 2-4 oz Auxigro 14-30 days after first application.</p>

Lawn and Turfgrasses (nonfood uses)	To increase the seedling establishment and growth of many lawn and turfgrass species:
First application:	Apply 2 – 8 oz of AuxiGro after 1-3" of new growth after seeding or re-growth appears for established turf.
Later applications:	Apply 2 – 8 oz of AuxiGro 14-30 days after the initial application. However, a 7 day application interval may be used for seedling turf.
<i>AuxiGro applications to turfgrass and ornamental plants must not exceed a total application rate of 128 oz. AuxiGro per acre per season.</i>	
Ornamentals (terrestrial nonfood uses)	To increase the growth, development and re-establishment of seedling, cuttings, or mature ornamental plant species.
First application:	Apply 2- 16 oz of AuxiGro as a soil drench or foliar spray.
Later applications:	Apply 2- 16 oz of AuxiGro as a soil drench or foliar spray at an interval of no less than 14 days from any previous application.
<i>AuxiGro applications to turfgrass and ornamental plants must not exceed a total application rate of 128 oz. AuxiGro per acre per season.</i>	

***AuxiGro should be applied with silicone-based surfactants unless tank mix combinations prohibit their use or crop injury is likely to occur.**

ROTATIONAL CROPS

All crops may be safely planted at any time following an AuxiGro application.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PRODUCT STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: DO NOT reuse any part of the packaging. Dispose of all containers and packaging in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of the smoke.

CONDITIONS OF SALE AND WARRANTY

The label instructions for the use of this product are based on research and field use. The instructions are believed to be reliable and must be followed carefully. However, it is not possible to eliminate all risks inherently associated with use of this product. Crop injury,

ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of Emerald Bio Corporation. All such risks shall be assumed by the user.

Emerald BioAgriculture Corporation (Emerald Bio) shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

Emerald Bio Corporation warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. EMERALD BIO CORPORATION DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND EMERALD BIO CORPORATIONS' EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF AUXIGRO. In no case shall Emerald Bio Corporation or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Buyers and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

Emerald BioAgriculture Corporation
3125 Sovereign Drive, Suite B
Lansing, MI 48911-4240
Toll Free Phone: 1-888-828-9346

® Trademark of Emerald Bio Corporation

