
 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Biopesticides and Pollution Prevention Division (7511C) 1200 Pennsylvania Avenue NW Washington, DC 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration (under FIFRA, as amended)</p>	EPA Reg. Number: 70810-5	Date of Issuance: AUG 16 2000
	Term of Issuance: Unconditional	
	Name of Pesticide Product: AuxiGro™ Soluble Granule Plant Metabolic Primer	
Name and Address of Registrant (include ZIP Code): Auxein Corporation 3125 Sovereign Dr., Ste. B Lansing, MI 48911-4240		
<p>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</p> <p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This registration does not eliminate the need for continual reassessment of the pesticide. If EPA determines at any time, that additional data are required to maintain in effect an existing registration, the Agency will require submission of such data under section 3(c)(2)(B) of FIFRA.</p> <p>This product is registered in accordance with FIFRA section 3(c)(5) and is subject to the following terms and conditions:</p> <ol style="list-style-type: none"> 1. Make the following modification to your label before you release your product for shipment: <ol style="list-style-type: none"> a. Revise the registration number to read 70810-5. 		
Signature of Approving Official: (See second page for signature)	Date: AUG 16 2000	

Registration number: 70810-5

2. Submit two (2) copies of the revised final printed labeling before you release the product for shipment.

A stamped copy of the label is enclosed for your records.

Sincerely,

A handwritten signature in cursive script, appearing to read "Janet L. Andersen".


Janet L. Andersen, Director
Biopesticides and Pollution Prevention Division

3/15

Aux01F24
8-3-00 (3)

AuxiGro™ Soluble Granule Plant Metabolic Primer

FOR INCREASED YIELD AND QUALITY



EPA Registration Number 70810-⁵~~X~~
EPA Establishment Number 33967-NJ-1, 39578-TX-1

Active Ingredients:

GABA: gamma aminobutyric acid -----	27.1 %
L-Glutamic acid -----	27.1 %
Other Ingredients -----	45.8 %
TOTAL -----	100.0 %

(1 pound contains 0.6 pounds of active ingredient)
(5 pounds contains 3 pounds of active ingredient)

**KEEP OUT OF REACH OF CHILDREN
CAUTION!/PRECAUTION!**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand this label, find someone to explain it to you in detail.)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS – CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

FIRST AID

If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15 or 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes • Remove contact lenses, if present, after 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

Personal Protective Equipment (PPE). Applicators and other handlers must wear:

- long sleeve shirt and long pants
- shoes plus socks
- protective eyewear

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if product gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

DO NOT apply directly to water or areas where surface water is present, or to intertidal areas below the mean high water mark.

DO NOT contaminate water when cleaning equipment or disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Please see package insert on bottom of bag for complete Directions For Use.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notifications and emergency assistance. It also contains specific instructions and exceptions pertaining to the restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Re-Entry Statement:

Do not enter or allow worker entry into treated areas without protective clothing during the restricted-entry interval (REI) of 4 hours.

For Early Entry

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated (plants, soil) wear:

- ◆ Long sleeve shirt and long pants
- ◆ Shoes plus socks
- ◆ Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas without protective clothing within 4 hours of application.

Observe all cautions and limitations on this label and on the labels of products used in combination with Auxigro. Do not use Auxigro other than in accordance with the instructions

Aux01F24
8-3-00 (3)

set forth on this label. The use of Auxigro not consistent with this label may result in injury to crops.

GENERAL INFORMATION

Auxigro can be applied on all labeled crops per label application instructions (Directions for Use). When used as directed, Auxigro has been shown to increase yields and/or quality of crops. Auxigro applications have also resulted in increased size and/or quality of treated commodities, early ripening in certain vegetables, increased root growth, early flowering and fruit set, faster seed germination and rooting.

MIXING INSTRUCTIONS

Mixing of Auxigro: Fill the spray tank one-half full to three-quarters full with clean water. Use a calibrated device to weigh the required amount of Auxigro. Add Auxigro to the spray tank while agitating. Add other products, adjuvants or surfactants and then fill the remainder of the tank with water. Mixing containers, hoses and spraying equipment should be cleaned or triple rinsed prior to mixing Auxigro for application

Tankmix combinations with Auxigro: When Auxigro is used in combination with other pesticide products, refer to the respective labels for instructions, such as rates, timing and methods of application, the disease or insects controlled and other labeled restrictions and precautions. No label dosage rate should be exceeded and the most restrictive label precautions and limitations should be followed.

Compatibility Testing: Auxigro is believed to be compatible with most agricultural fungicides, insecticides, micronutrients and spray adjuvants. If compatibility with another product is unknown, it is advisable to conduct a tank spray compatibility test when you plan to mix Auxigro with other products. To determine the physical compatibility of this product with other products, use a jar test. Using a quart jar, add the proportionate amounts of the products to approximately one quart of water with agitation. After thorough mixing, allow this mixture to stand for 5 minutes. If the combination remains mixed or can be readily remixed, it is physically compatible. Read and recognize all cautions and limitations of labeling of all products used in mixtures.

Adjuvants and Surfactants: Use a silicone-based surfactant at the lowest recommended label rate for Plant Growth Regulators (PGRs) or foliar sprays (typically 4 to 6 ounces per 100 gallons). If the other product limits the use of any type of surfactant, follow the other label recommendation for use of surfactants. If rain or overhead irrigation is expected from 4 to 24 hours of application, use an appropriate sticker to prevent spray deposit wash-off. DO NOT apply Auxigro if rain or overhead irrigation is expected within 4 hours.

APPLICATION INSTRUCTIONS

Application of Auxigro can effectively enhance plant development and growth of numerous plant species when properly applied. Applications of Auxigro may begin at seeding (or transplanting) and continue until harvest. For optimal yield, apply Auxigro postemergence according to the application timings and rates provided in the table below. Multiple applications (with fungicides and foliar nutritional products) may typically be made at a rate of 1 - 4 ounces per acre for each application. Total season application should not exceed 24 ounces per acre per crop.

All rates of application for Auxigro are given as broadcast rates and expressed as ounces of formulated product per acre per application. If crop growth stage will permit a band treatment, use the broadcast rate on proportionally less area to reduce the total amount of product applied. **Auxigro should be applied with silicone-based surfactants unless tankmix combinations prohibit their use or crop injury is likely to occur.**

Allow heavy dew to evaporate prior to Auxigro applications. Do not apply Auxigro if windy conditions exist or a strong threat of rain exists within 4 hours of application.

Ground Application: Apply uniformly with properly calibrated ground equipment in 20 to 400 gallons of water per acre for conventional dilute, air blast spray equipment, concentrate or electrostatic spray equipment. Use nozzles and spray delivery methods that insure thorough coverage of foliage and fruit. Application volume and pressure may be adjusted, based on plant size, foliage density and sprayer selected.

Aerial Application: Thorough spray coverage is required for maximum benefit. To achieve good coverage, use proper droplet size, spray uniformly, and ensure swath width and spray volume are correct. Do not use less than 5 gallons total spray volume per acre.

Overhead Sprinkler Irrigation (For Potatoes Only. Use typical surfactants or additives that are recommended for overhead irrigation. Do not use a silicone based surfactant when applying Auxigro by overhead irrigation): Auxigro can be applied through common overhead sprinkler systems, such as center-point, lateral move, end tow, side (wheel) roll, traveler and others. All equipment must be properly calibrated for chemical application before use.

Center-pivot, Lateral Move, End Tow, and Traveler Irrigation Equipment. Do not apply Auxigro with more than 1/4 inches of water (6750 gallons of water) per acre when the system and injector are operated at normal pressures recommended by their manufacturer.

Solid Set, Side (wheel) Roll and Hand Move Irrigation Equipment. Inject Auxigro at the end of the irrigation cycle or as a separate application to maximize the foliar retention.

The system must contain a functional check valve, vacuum relief valve and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. The chemical injection pipeline must also contain a functional, normally closed

7/15

Aux01F24
8-3-00 (3)

solenoid operated valve located on the intake side of the injection pump and connected to a system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the chemical injection pump when the water pump motor stops. The irrigation line or water pump must include functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where chemical distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with chemicals and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift, when system connections or fittings leak, when nozzles do not provide uniform distribution or when lines containing the product must be dismantled and drained.

Lack of effectiveness may result from non-uniform distribution of treated water or excessively high application volumes (more than 1/4 inch per acre).

Allow sufficient time for the chemical to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut the system down and make necessary adjustments should the need arise.

Do not connect an irrigation system (including greenhouse systems) used for chemical application to a public water system unless the label-prescribed safety devices for drawing water supplies are in place.

DIRECTIONS FOR USE

Crops & Application(s)	Rate / Acre / Application & Timing*
Celery.	To increase the size and/or total yield.
First application:	Apply 2 - 4 oz. of AuxGro when plants are 6 - 8 inches tall.
Second application:	Apply 2 - 4 oz. of AuxGro when plants are 14 - 16 inches tall, approximately 14 - 21 days after the initial application, no sooner than 12 days after the initial application.

Aux01F24
8-3-00 (3)

Cucumbers (fresh market).	To increase the grade and/or marketable yield.
First application:	Apply 2 - 4 oz. of AuxiGro at early bloom, just prior to running.
Second application:	Apply 2 - 4 oz. of AuxiGro at 10-14 days after the initial application.
Other applications:	Apply 2 - 4 oz. of AuxiGro at 14-16 day intervals when multiple harvests are taken. Total season application should not exceed 24 oz. per acre per crop.
Edible Dry Beans (navy and pinto varieties only)	To enhance yields
One application:	Apply 2 - 4 oz. of AuxiGro to the crop at flower bud initiation (less than 5% bloom) OR at initial pod fill stage of growth (approximately 21 days after first bud formation). Apply by ground application only.
Grapes (<i>Vitis vinifera</i> varieties only).	To increase Soluble Sugars as measured by degrees Brix.
First application:	Apply 2 - 4 oz. of AuxiGro at full bloom (beginning of berry development).
Second application:	Apply 2 - 4 oz. of AuxiGro at the onset of veraison (beginning of ripening).
Bulb Onions.	To increase the total yield and/or marketable grade.
First application:	Apply 2 - 4 oz. of AuxiGro at the initiation of bulbing (1 ¼ in. to 1 ½ in. in diameter).
Second application:	Apply 2 - 4 oz. of AuxiGro at approximately 4 weeks prior to harvest, a minimum of 21 days after the initial application.
Peppers (bell, green, jalapeño).	To increase the marketable grade and/or total yield.
First application:	Apply 2 - 4 oz. of AuxiGro at flower bud initiation (pre-bloom of main canopy).
Second application:	Apply 2 - 4 oz. of AuxiGro after first fruit set, no sooner than 14 days after initial application.

<p>Head lettuce (iceberg).</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the marketable yield and/or provide earlier maturity.</p> <p>Apply 2 - 8 oz. of AuxiGro immediately after thinning (4 - 6 true leaves).</p> <p>Apply 2 - 8 oz. of AuxiGro at first sign of head cupping, no sooner than 14 days after first application.</p>
<p>Leaf lettuce (romaine, butter).</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the marketable yield and/or provide earlier maturity.</p> <p>Apply 2 - 8 oz. of AuxiGro immediately after thinning (4 - 6 true leaves).</p> <p>Apply 2 - 8 oz. of AuxiGro at first sign of head cupping, no sooner than 14 days after first application.</p>
<p>Peanuts.</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the total yield.</p> <p>Apply 2 - 4 oz. of AuxiGro at early bloom, 18 - 20 trifoliates (prior to running).</p> <p>Apply 2 - 4 oz. of AuxiGro at full bloom, approximately 25 -30 days after initial application.</p>
<p>Potatoes.</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the marketable yield and/or improve tuber grades.</p> <p>Apply 3 - 5 oz. of AuxiGro at first tuber initiation when tubers are no larger than a small pea (plants 6 - 10 inches in height, typically 2 -3 weeks after emergence).</p> <p>Apply 3 - 5 oz. of AuxiGro 21-28 days after first application.</p>
<p>Snap Beans (<i>Phaseolus spp</i>) (Southeast U.S. recommendation only).</p> <p>One application:</p>	<p>To increase the total yield.</p> <p>Apply 4 -8 oz. of AuxiGro at pre-bloom (bud formation).</p>

10/15

Aux01F24
8-3-00 (3)

Strawberries.	To increase the total yield and/or quality.
Ever Bearing Varieties.	(harvest continues several months)
First application:	Apply 4 - 8 oz. of Auxigro at any time after first harvest.
Later applications:	Additional 4 - 8 oz. applications of Auxigro should be made at 1-month intervals following the first application. Total season application should not exceed 24 oz. per acre per crop.
June Bearing Varieties.	(harvest duration is typically less than 3months)
First Application:	Apply 4 -8 oz. Of Auxigro 5 - 7 days prior to first harvest.
Second Application:	Apply 4 - 8 oz. Of Auxigro 14 - 21 days after initial application.
Tomatoes (processing).	To increase the marketable yield and to maintain or improve the soluble sugar levels as measured by degrees Brix.
First application:	Apply 2 - 4 oz. of Auxigro at full bloom in the canopy (50-70% bloom).
Second application:	Apply 2 - 4 oz. of Auxigro at first fruit set, approximately 10 -21 days after first application.
Tomatoes (fresh).	To increase the marketable yield and/or improve grades.
First application:	Apply 2 - 4 oz. of Auxigro at pre-bloom (bud formation for initial production set).
Second application:	Apply 2 - 4 oz. of Auxigro at full bloom, at least 21 days after first application.
Multiple applications:	Additional 2 - 4 oz. applications may be made after the second application at intervals of at least 21 days, if multiple harvests over an extended period are taken. Total season application should not exceed 24 oz. per acre per crop.
Watermelons.	To increase the yield and/or quality.
First application:	Apply 2 - 4 oz. of Auxigro at early bloom (less than 10%), just prior to running.
Second application:	Apply 2 - 4 oz. of Auxigro 14 - 21 days following the initial application.

CROP GROUPINGS GENERAL USE DIRECTIONS

Crop Group 1: Root and Tubers (Rates and directions apply to all root and tuber crops except potatoes)	To increase the total yield and/or quality of selected root and tuber crops:
First application:	Apply 2 - 8 oz of AuxiGro at first tuber initiation or the beginning of bulking up as appropriate.
Second application:	Apply 2 - 8 oz of AuxiGro 21-28 days after initial application or approximately 30 days prior to harvest.
Crop Group 3: Bulb Vegetables (Rates and directions apply to all Bulb Vegetables except bulb onions).	To increase the size and/or total yield of onions and garlic:
First application:	Apply 2 - 8 oz of AuxiGro when the plant leaves are 8-12" tall.
Second application:	Apply 2 - 8 oz of AuxiGro at start of "bulking-up."
Crop Group 4: Leafy Vegetables (Rates apply to all Leafy Vegetables crops except celery, head lettuce and leaf lettuce)	To increase the size and/or total yield of celery and other leafy vegetables.
First application:	Apply 2 - 8 oz of AuxiGro at 1-5 days after thinning or transplant.
Second application:	Apply 2 - 8 oz of AuxiGro 14 - 30 days after first application
Crop Group 5: Cole crops	To increase the total yield and/or marketable grade.
First application:	Apply 2 - 8 oz. AuxiGro at the 4-8 leaf stage.
Second application:	Apply 2 - 8 oz. AuxiGro just prior to head formation, no sooner than 14 days after first application.
Crop Group 6: Legume Vegetables (Rates and timing of application apply to all Legume Vegetables except Snap Beans and Edible Dry Beans - navy and pinto).	To increase the total yield and/or quality of various legume vegetables:
First application:	Apply 2 - 8 oz. AuxiGro at 5 - 7 days after first pod formation.
Second application:	Apply 2 - 8 oz AuxiGro at 10 - 14 days after first application. <i>AuxiGro may also be applied at flower bud initiation, an earlier application timing, as a substitution for the first application on some legume species.</i>

<p>Grapes</p> <p>First application:</p> <p>Second application:</p> <p>Third application:</p>	<p>To increase Soluble Sugars as measured by degrees Brix.</p> <p>Apply 2 - 4 oz. of Auxigro at full bloom (beginning of berry development).</p> <p>Apply 2 - 4 oz. of Auxigro at the onset of veraison (beginning of ripening).</p> <p>Apply 2 - 4 oz. of Auxigro 2 to 3 weeks after the 2nd application.</p>
<p>Crop Group 8: Fruiting Vegetables (Separate use directions are provided for tomatoes and peppers.)</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the yield of various fruiting vegetables:</p> <p>Apply 2 - 8 oz. Auxigro at bud set (initiation) in canopy.</p> <p>Apply 2 - 8 oz. Auxigro at full bloom, no sooner than 10 - 14 days after initial application.</p>
<p>Crop Group 9: Cucurbit vegetables (For use on all Cucurbit vegetables except cucumber and watermelon).</p> <p>First application:</p> <p>Second application:</p>	<p>To increase the total yield and/or quality of various cucurbits:</p> <p>Apply 2 - 8 oz. Auxigro at bud formation pre-bloom, prior to running.</p> <p>Apply 2 - 8 oz. Auxigro 14 - 21 days after first application.</p>
<p>Crop Group 10: Citrus Fruits</p> <p>First application:</p> <p>Later applications:</p>	<p>To increase the growth and development or the total yield and/or quality.</p> <p>Apply 1 - 8 oz of Auxigro at first fruit set.</p> <p>Apply 1 - 8 oz of Auxigro every 4 - 8 weeks after first application.</p>
<p>Crop Group 11: Pome Fruits</p> <p>First application:</p> <p>Second application:</p> <p>Third application:</p> <p>Fourth application:</p>	<p>To enhance the total yield and/or quality of the fruit.</p> <p>Apply 2-8 oz of Auxigro at pink bud.</p> <p>Apply 2-8 oz of Auxigro at full bloom.</p> <p>Apply 2-8 oz of Auxigro at petal fall.</p> <p>Apply 2-8 oz of Auxigro two weeks after petal fall (if needed).</p>
<p>Crop Group 12: Stone Fruits</p> <p>First application:</p> <p>Second application:</p> <p>Third application:</p> <p>Fourth application:</p>	<p>To increase the total yield and/or quality the fruit:</p> <p>Apply 2-8 oz of Auxigro at pink bud .</p> <p>Apply 2-8 oz of Auxigro at full bloom.</p> <p>Apply 2-8 oz of Auxigro at petal fall.</p> <p>Apply 2-8 oz of Auxigro two weeks after petal fall (if needed).</p>
<p>Crop Group 13: Berries Group</p> <p>First application:</p> <p>Second application:</p> <p>Third application:</p> <p>Fourth application:</p>	<p>To increase the total yield and/or quality of berries:</p> <p>Apply 2-8 oz of Auxigro at pink bud.</p> <p>Apply 2-8 oz of Auxigro at full bloom.</p> <p>Apply 2-8 oz of Auxigro at petal fall.</p> <p>Apply 2-8 oz of Auxigro two weeks after petal fall (if needed).</p>

13/15

Aux01F24
8-3-00 (3)

Crop Group 14: Almonds	To increase the total yield and/or quality the nuts: First application: Apply 2-8 oz of Auxigro at pink bud Second application: Apply 2-8 oz of Auxigro at full bloom. Third application: Apply 2-8 oz of Auxigro at petal fall. Fourth application: Apply 2-8 oz of Auxigro two weeks after petal fall (if needed).
Crop Group 14: Tree Nuts Group (except almonds)	To Increase total yield and/or quality and the Nuts: First application: Apply 2-8 oz of Auxigro at pink bud Second application: Apply 2-8 oz of Auxigro at full bloom. Third application: Apply 2-8 oz of Auxigro at petal fall. Fourth application: Apply 2-8 oz of Auxigro two weeks after petal fall (if needed).
Crop Group 15: Cereal grains group	To increase the total yield and or quality. First application: On small grains (i.e., wheat, oats, rice, etc.) apply 2 – 4 oz. Auxigro prior to tillering. On corn and sorghum, apply 2 – 4 oz. Auxigro at V-4 to V-6 growth stage. Second application: On small grains, apply 2 – 4 oz. Auxigro at boot stage. On corn and sorghum, apply 2 – 4 oz. Auxigro at V-10 to V12 growth stage.
Crop Group 16: Forage, fodder and straw of cereal grains group.	To increase the yield and or quality of various forage crops: First application: Apply 2-4 oz Auxigro prior to tillering on small grains and forage grasses or at V-4 to V-6 growth stage for corn and sorghum. Second application: Apply 2-4 oz Auxigro at boot stage for small grains and forage grasses and at V-10 to V12 growth stage for corn and sorghum.
Crop Group 17: Grass forage, fodder, and hay.	To increase the yield and or quality of various forage crops: First application: Apply 2-4 oz Auxigro when 2-6 inches of new growth appears. Later applications: Apply 2-4 oz Auxigro every 4 weeks after first application.
Crop Group 18: Non-grass animal feeds	To increase the yield and or quality of the forage. First application: Apply 2-4 oz Auxigro when 2-4 inches of new growth appears. Second application: Apply 2-4 oz Auxigro when 2-4 inches of re-growth appears after cutting.
Crop Group 19: Herbs and Spices	To increase the yield and/or quality of various herb and spice crops: First application: Apply 2-4 oz Auxigro when 2-4 inches of new vegetative growth appears. Second application: Apply 2-4 oz Auxigro 14-30 days after first application.

Lawn and Turfgrasses (nonfood uses)	To increase the seedling establishment and growth of many lawn and turfgrass species:
First application:	Apply 2 – 8 oz of AuxiGro after 1-3" of new growth after seeding or re-growth appears for established turf.
Later applications:	Apply 2 – 8 oz of AuxiGro every 14-30 days after the initial application. However, a 7 day application interval may be used for seedling turf.
<i>AuxiGro applications to turfgrass and ornamental plants must not exceed a total application rate of 128 oz. AuxiGro per acre per season.</i>	
Ornamentals (terrestrial nonfood uses)	To increase the growth, development and re-establishment of seedling, cuttings, or mature ornamental plant species.
First application:	Apply 2- 16 oz of AuxiGro as a soil drench or foliar spray.
Later applications:	Apply 2- 16 oz of AuxiGro as a soil drench or foliar spray at an interval of no less than 14 days from any previous application.
<i>AuxiGro applications to turfgrass and ornamental plants must not exceed a total application rate of 128 oz. AuxiGro per acre per season.</i>	

AuxiGro should be applied with silicone-based surfactants unless tankmix combinations prohibit their use or crop injury is likely to occur.

ROTATIONAL CROPS

All crops may be safely planted at any time following an AuxiGro application.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

PRODUCT STORAGE: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of reach of children, preferably in a locked storage area.

PRODUCT DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: **DO NOT** reuse any part of the packaging. Dispose of all containers and packaging in a sanitary landfill, by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of the smoke.

15/15

Aux01F24
8-3-00 (3)

CONDITIONS OF SALE AND WARRANTY

The label instructions for the use of this product are based on research and field use. The instructions are believed to be reliable and must be followed carefully. However, it is not possible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the use of, or application of the product contrary to label instructions, all of which are beyond the control of Auxein Corporation. All such risks shall be assumed by the user.

Auxein Corporation shall not be responsible for losses or damages resulting from use of this product in any manner not set forth on this label. User assumes all risks associated with the use of this product in any manner not specifically set forth on this label.

Auxein Corporation warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. AUXEIN CORPORATION DOES NOT MAKE OR AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTIES, EXPRESS OR IMPLIED AND EXPRESSLY EXCLUDES AND DISCLAIMS ALL IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE.

BUYER'S EXCLUSIVE REMEDY AND AUXEIN CORPORATIONS' EXCLUSIVE LIABILITY; WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF AUXIGRO. In no case shall Auxein Corporation or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. Buyers and all users are deemed to have accepted the terms of this LIMIT OF WARRANTY AND LIABILITY which may not be varied by any verbal or written agreement.

**Auxein Corporation
3125 Sovereign Drive, Suite B
Lansing, MI 48911-4240
1-888-3AUXEIN (1-888-828-9346)**

™ Trademark of Auxein Corporation

(Revised August 4, 2000)