


70870-1

04/26/2005

1/26



Master Label

**Agricure**

FOR ORGANIC PRODUCTION

A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER  
DISEASES ON TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS,  
GREENHOUSE AND AGRICULTURAL CROPS, AND TURF

ACTIVE INGREDIENT:

Potassium Bicarbonate.....85.00%

OTHER INGREDIENTS:.....15.00%

TOTAL.....100.00%

A: SUBLABEL - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS  
SUBJECT TO WORKER PROTECTION STANDARD

B: SUBLABEL - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS  
SUBJECT TO WORKER PROTECTION STANDARD - WATER  
SOLUBLE PACKAGING LABEL

C: SUBLABEL - HOUSEHOLD PRODUCTS: FOR NON-COMMERCIAL  
APPLICATORS (NOT SUBJECT TO WORKER PROTECTION  
STANDARD)

EPA Reg. No. 70870-1

EPA Establishment No. XXXXX-XX-XXX

Produced For:

H & I AGRITECH, INC.  
95 BROWN ROAD, BOX 1030  
ITHACA, NY 14850-1257

2/26

**SUBLABEL A - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS**  
**SUBJECT TO WORKER PROTECTION STANDARD**

## **AGRICURE**

FOR ORGANIC PRODUCTION

A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER  
DISEASES ON TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS,  
GREENHOUSE AND AGRICULTURAL CROPS, AND TURF

This package can treat up to \_\_\_\_\_ acre(s) at the recommended maximum treatment rate.

**ACTIVE INGREDIENT:**

Potassium Bicarbonate.....85.00%

OTHER INGREDIENTS.....15.00%

TOTAL.....100.00%

**KEEP OUT OF REACH OF CHILDREN**

### **CAUTION**

See side (or back) panel for the First Aid Information.

Net weight:           5 lbs. (2.27 kg )  
                          10 lbs. (4.54 kg )  
                          25 lbs. (11.35 kg )  
                          50 lbs. (22.7 kg )  
                          100 lbs. (45.4 kg )

Produced For:

H & I AGRITECH, INC.  
95 BROWN ROAD BOX 1030  
ITHACA, NY 14850 - 1257

EPA REG. NO. 70870-1  
EPA EST. NO.

3/26  
1

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. Causes moderate-eye irritation. Avoid contact with eyes or clothing.

**FIRST AID**

- If swallowed:
- Call poison control center or doctor immediately for treatment advice.
  - Have person drink several glasses of water.
  - Do not induce vomiting.
  - Do not give anything by mouth to an unconscious person.
- If in eyes:
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have a product label or container with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858 7378 between the hours of 6:30 a.m. to 4:30 p.m. Pacific time, seven days a week for emergency medical information.

**PERSONAL PROTECTIVE EQUIPMENT**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters and rinsate.

**DIRECTIONS FOR USE**

4/20

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

#### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

There is a restricted entry of one (1) hour for this product.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeved shirt and short pants, waterproof gloves and shoes plus socks.

#### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

#### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a cool, dry place. Prevent exposure to moisture. Keep container tightly closed and out of reach of children. Avoid contamination with other pesticides or fertilizers.

**PESTICIDE SPILL PROCEDURE:** In case of a spill, carefully sweep up the material to avoid generating dust. Store in a labeled container for disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke

5/26  
1

### GENERAL INFORMATION

Agricure is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental plants, and on greenhouse, nursery and agricultural crops, and on turf. Agricure is applied using a sufficient volume of water to insure complete coverage of all stems and foliage.

**NOTICE TO USER:** Due to variability within species and in application techniques and equipment, and possible tank mix combinations, neither the manufacturer nor the seller has determined whether Agricure can be safely used on all varieties and species of ornamentals, greenhouse, turf and agricultural plants under all conditions. Therefore, the professional user should determine if Agricure can be used safely before broad usage by applying the prescribed use rate under the conditions expected to be encountered to a small test area. Any adverse effects should be visible within 14 days.

For curative disease control on sensitive crops apply at a rate of 2.5 pounds of Agricure per 100 gallons of water per week and increase the rate each week in 0.25 pound increments. Do not exceed a use rate of 5.0 pounds per acre per application .

**IMPORTANT:** Read the entire label before using Agricure. Consult your State Agricultural Experiment Station or Extension Service Specialist for additional information on timing of applications, rates, and any additional requirements or restrictions.

### MIXING INSTRUCTIONS

Agricure must be diluted with water before applying. Carefully measure and mix the product and water.

1. Fill tank with 1/2 to 3/4 of the selected volume of water prior to the addition of Agricure. Start mechanical or hydraulic agitation.
2. Slowly add the required amount of Agricure to the tank.
3. Continue to fill the tank with water to the desired volume while agitating to promote uniform distribution and prevent settling.

DO NOT MIX WITH OTHER SOLUBLE PESTICIDES OR FERTILIZERS NOT COMPATIBLE WITH MILD ALKALINE SOLUTIONS. THE ADDITION OF SURFACTANT OR WETTING AGENT TO SPRAY IS NOT REQUIRED. PRODUCT IS PRE-FORMULATED WITH THESE PRODUCTS FOR BEST PERFORMANCE. DO NOT ADJUST SPRAY SOLUTION pH AFTER MIXING OF PRODUCT. ACIDIFICATION OF SOLUTION WILL CAUSE REDUCED PRODUCT PERFORMANCE.

### SPRAY APPLICATION METHOD

Do not apply this product through any type of irrigation system. Do not apply directly to soil as a

6/20  
1

directed spray or drench. For best results, use a properly calibrated sprayer or similar equipment, and check equipment calibration frequently.

### **CLEANING SPRAY EQUIPMENT**

Clean application equipment thoroughly using a detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

### **FIELD APPLICATION**

Apply at a rate of 2.0 to 5 pounds of Agricure per acre. Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per acre of treatment can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 5 pounds of Agricure per acre, per application. Do not store unused spray solution. Dispose of unused solution on site.

### **GREENHOUSE APPLICATION**

Apply at a rate of 1.25 to 5.0 pounds per 100 gallons of water. Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 0.5 pounds of Agricure per 4350 sq.ft. or 1.15 pounds per 10,000 sq. ft., per application. Do not store unused spray solution. Dispose of unused solution on site.

See special instructions for use on poinsettia, impatiens and pansy.

### **Small Volume Application Chart**

Use the chart below to determine Agricure quantities for small area applications. Adjust to desired concentration for low disease pressure or as a curative treatment.

Chart Treatment Rate: 2.5 lbs. per 100 gallons

Amount of Agricure	Amount of Water	Square feet treated
1 Tbsp	1 gallon	500
2 Tbsp	2 gallons	1,000
4 Tbsp	4 gallons	2,000
1/8 cup	2 gallons	1,000
1/4 cup	4 gallons	2,000
1/2 cup	8 gallons	4,000

7/20

1 oz. (28 grams)	2.5 gallons	1,100
2 ozs. (57 grams)	5 gallons	2,200
4 ozs. (113 grams)	10 gallons	4,400

### **ELECTROSTATIC SPRAYERS**

Apply at a rate of 0.625 pounds (10 ounces) to 1.25 pounds (20 ounces) per 2 gallons of water per 10,000 sq. ft. Do not exceed a rate of 1.5 pounds (24 ounces) per 2 gallons of water.

### **THERMAL FOGGERS**

Apply at a rate of 0.38 pounds (6 ounces) to 0.75 pounds (12 ounces) per 1 gallon of water per 10,000 sq. ft. Use the 0.38 pounds (6 ounces) per 1 gallon of water rate for plants that are in flower. Do not exceed 0.75 pounds (12 ounces) per 1 gallon of water.

### **TURF APPLICATION**

Apply at a rate of 2.5 to 5 pounds of Agrisure per acre. Use a sufficient volume of spray solution to obtain complete coverage of foliage. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per acre of treatment can vary depending upon stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 5 pounds of Agrisure per acre, per application. Do not store unused spray solution. Dispose of unused solution on site.

### **APPLICATION TIMING**

Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, apply Agrisure as a post-harvest treatment to remaining foliage to decrease overwintering inoculum.

### **PREHARVEST INTERVAL**

Agrisure can be applied up to and including the day of harvest.

### **ORNAMENTAL APPLICATIONS:**(California registered uses are listed in a separate section.)

Agrisure is for use in the following use sites: Greenhouse, Lathhouse, Shadehouse, Commercial and Residential Landscapes, Interiorscapes, Nursery, Container Production and Field Applications.

### **ORNAMENTAL PLANT TYPES:**

Agrisure is for use on the following plant types: Annual and Perennial Flowers, Bedding Plants, Foliage

Plants, Ground Covers, Woody and Ornamental Shade Trees, Woody and Herbaceous Ornamentals,

and Ornamental Nut and Fruit Trees.

### FOR USE ON:

Ageratum	Crape Myrtle	Impatiens*	Plum-Ornamental
Ajuga	Crassula	Iris	Poinsettia*
Almond-Ornamental	Croton	Juniper	Poppy
Alyssum	Delphinium	Kalanchoe	Pothos
Andromeda	Dianthus	Lavender	Primrose
Aphelandra	Dizygotheca	Lemon balm	Privet
Apple-Ornamental	Dogwood	Lipstick balm	Protea
Apricot-Ornamental	English Ivy	Lipstick vine	Prune-Ornamental
Artemisia	Episcia	(Aeschynanthus)	Pulmonaria
Aster	Euonymus	Marigold	Pyracantha
Azalea	Fern	Monarda (Bee balm)	Rhododendron
Boxwood	Forsythia	Nectarine-Ornamental	Rose
Calendula	Gazania	Pachysandra	Rose-of-Sharon
Carnation	Geranium	Palm	Schefflera
Centaurea montana	Gladiolus	Pansy*	Snapdragon
Cherry-Ornamental	Gloxinia	Peach-Ornamental	Statice
Chrysanthemum	Gypsophila	Peperomia	Tree Ivy
Cineraria	Hawthorn	Periwinkle	Verbena
Citrus	Heliopsis scabra	Philodendron	Viburnum
Coleus	Hibiscus	Phlox	Violet
Columbine	Holly	Pilea	Zinnia
Coral Bells	Hoya	Pine	Photinia
(Heuchera)	Hydrangea	Pittosporum	Spruce
Begonia	Ginseng	Fir	India-
hawthorn			
Clematis	Daylily	Gerbera daisy	

### DISEASES CONTROLLED

Alternaria blight	Fusarium
Anthraxnose	Helminthosporium leaf spot
Ascochyta blight	Ink spot
Blackspot	Phomopsis blight
Botrytis blight	Powdery mildew
Botrytis gray mold	Ovulinia sp.
Cercospora leaf spot	Ramularia leaf spot
Corynespora leaf spot	Ray blight
Didymellina leaf spot	Scab
Diplodia tip blight	Septoria leaf spot
Entomosporium leaf spot	Monilinia fruit and blossom rot
Downy mildew	Penicillium spp.
Rust	



9/26

**\*SPECIAL INSTRUCTIONS:**

Poinsettia

For control of intense powdery mildew, apply weekly at a rate of 0.25 pounds per 4350 sq. ft. or 1/10 acre during vegetative growth prior to bract formation, and apply every 14 days at 0.5 pounds per 4350 sq. ft. thereafter. A rate of 0.5 pounds per 4350 sq. ft applied weekly during vegetative growth has the potential to cause marginal burn to newly-developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens

For New Guinea impatiens, apply at a rate of 0.25 pounds per 4350 sq. ft. or 1/10 acre during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy

Do not use on pansies at early stages of growth.

**AGRICULTURAL APPLICATIONS:** (California registered uses are listed in a separate section.)  
Agricure is for use in the following use sites: Greenhouse, Orchard, Nursery and Field Applications.

**FOR USE ON:**

Acorn squash	Citrus	Okra	Snow peas
Almonds	Crenshaw melon	Oregano	Soybeans
Apples	Cucumber	Peaches	Spinach
Apricots	Dry beans	Pears	Strawberries
Alfalfa	Eggplant	Peas	Sugar beets
Avocado	Endive	Pecans	Summer squash
Barley	Escarole	Peppers	Sweet potatoes
Basil	Grapes	Pistachio	Thyme
Blueberries	Honeydew	Plums	Tomatoes
Broccoli	Hops	Potatoes	Turnip
Cabbage	Kale	Prunes	Walnut
Caneberries	Lettuce	Pumpkin	Watermelon
Cantaloupe	Lima bean	Radicchio	Wheat
Cauliflower	Mango	Rosemary	Winter squash
Cherries	Muskmelon	Sage	Zucchini
Chicory	Mustard greens	Snap beans	Tobacco

**DISEASES CONTROLLED:**

Alternaria leaf spot	Fusarium
Anthraxnose	Phoma
Botrytis	Phytophthora
Downy mildew	Powdery mildew

Cercospora leaf spot	Fly speck
Penicillium spp.	Monilinia fruit and blossom rot
Phomopsis blight	Scab
Septoria leaf spot	

### **TURF APPLICATION**

Agricure is for use in the following use sites: Commercial and residential lawns, golf course and other commercial turf grass sites.

### **FOR USE ON:**

Bentgrass  
 Bermudagrass  
 Blue grass, Kentucky, annual  
 Centipedegrass  
 Fescue, fine, tall, chewings, red  
 Ryegrass, perennial, annual  
 St. Augustinegrass  
 Zoysiagrass

### **DISEASES CONTROLLED**

Alternaria leaf spot	Downy mildew
Anthrachnose	Helminthosporium leaf spot
Ascochyta blight	Powdery mildew
Cercospora leaf spot	Septoria leaf spot

### **FOR USE ONLY IN CALIFORNIA**

**FOR USE ON:** Acorn squash, Almonds, Apples, Apricots, Cantaloupe, Cherries, Chicory, Crenshaw melon, Cucumber, Endive, Escarole, Grapes, Honeydew, Lettuce, Muskmelon, Peaches, Pears, Peppers, Pistachio, Plums, Prunes, Pumpkin, Radicchio, Rose, Spinach, Strawberries, Summer squash, Tomatoes, Walnut, Watermelon, Winter squash, Zucchini

### **DISEASES CONTROLLED**

Downy mildew	Powdery mildew
--------------	----------------

### **NOTICE**

Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

### **IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY**

This product has been researched to provide necessary data to support its use on the ornamental, greenhouse and agricultural plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. Always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

**NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH**

11/26

**DIRECTIONS OR ESTABLISHED SAFE PRACTICE.** To the extent allowable by state law the exclusive remedy of the user or Buyer, and the limit of liability of H & I Agritech, Inc. or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of H & I Agritech Inc.

**SUBLABEL B - COMMERCIAL PRODUCTS: FOR COMMERCIAL APPLICATORS  
SUBJECT TO WORKER PROTECTION STANDARD - WATER SOLUBLE PACKAGING  
LABEL**

**AGRICURE WATER SOLUBLE PACKAGE**

**FOR ORGANIC PRODUCTION**

**A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER  
DISEASES ON TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS,  
GREENHOUSE AND AGRICULTURAL CROPS, AND TURF**

This package can treat up to \_\_\_\_\_ sq. ft. (acres) at the recommended maximum treatment rate.

**ACTIVE INGREDIENT:**

Potassium Bicarbonate.....85.00%

INERT INGREDIENTS.....15.00%

TOTAL.....100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side (or back) panel for First Aid Information.

Net weight:           1.0 oz. (28.35 g)  
                              4.0 oz. (113.4 g)  
                              8.0 oz. (227 g)  
                              2.5 lbs. (1.14 kg)

Produced For:

12/20

H & I AGRITECH, INC.  
95 BROWN ROAD BOX 1030  
ITHACA, NY 14850 - 1257

EPA REG. NO. 70870-1  
EPA EST. NO.

**PRECAUTIONARY STATEMENTS**  
**HAZARD TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing

**FIRST AID**

- If swallowed:
- Call poison control center or doctor immediately for treatment advice.
  - Have person drink several glasses of water.
  - Do not induce vomiting.
  - Do not give anything by mouth to an unconscious person.
- If in eyes:
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have a product label or container with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858 7378 between the hours of 6:30 a.m. to 4:30 p.m. Pacific time, seven days a week for emergency medical information.

**PERSONAL PROTECTIVE EQUIPMENT**

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

**ENVIRONMENTAL HAZARDS**

13/26

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters and rinsate.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

### **AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

There is a restricted entry of one (1) hour for this product.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeved shirt and short pants, waterproof gloves, shoes plus socks.

### **NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

### **STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in a cool, dry place. Prevent exposure to moisture. Keep container tightly closed and out of reach of children. Avoid contamination with other pesticides or fertilizers.

**PESTICIDE SPILL PROCEDURE:** In case of a spill, carefully sweep up the material to avoid generating dust. Store in a labeled container for disposal.

**PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

14/26

## GENERAL INFORMATION

Agricure is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental plants, and on greenhouse, nursery and agricultural crops, and turf. Agricure is applied using a sufficient volume of water to insure complete coverage of all stems and foliage.

**NOTICE TO USER:** Due to variability within species and in application techniques and equipment, and possible tank mix combinations, neither the manufacturer nor the seller has determined whether Agricure can be safely used on all varieties and species of ornamentals, greenhouse, turf and agricultural plants under all conditions. Therefore, the professional user should determine if Agricure can be used safely before broad usage by applying the prescribed use rate under the conditions expected to be encountered to a small test area. Any adverse effects should be visible within 14 days.

### (1 ounce package)

For curative disease control on sensitive crops apply at a rate of 1 ounce (1 package) of Agricure per 2.5 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 1 ounce per 500 sq. ft. per week.

### (4 ounce package)

For curative disease control on sensitive crops apply at a rate of 4 ounces (1 package) of Agricure per 10 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 4 ounces per 2175 sq. ft. per application .

### (8 ounce package)

For curative disease control on sensitive crops apply at a rate of 8 ounces (1 package) of Agricure per 20 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 8 ounces per 4350 sq. ft. or 1/10 acre per application .

### (2.5 pound package)

For curative disease control on sensitive crops apply at a rate of 2.5 pounds of Agricure per 100 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 2.5 pounds per 21,750 sq. ft. or 1/2 acre per application .

**IMPORTANT:** Read the entire label before using Agricure. Consult your State Agricultural Experiment Station or Extension Service Specialist for additional information on timing of applications, rates, and any additional requirements or restrictions.

## MIXING INSTRUCTIONS

Agricure must be diluted with water before applying. Carefully measure and mix the product and water.

15/20  
1

1. Fill tank with 1/2 to 3/4 of the selected volume of water prior to the addition Agricure. Start mechanical or hydraulic agitation.
2. Slowly add the required number of packages of Agricure to the tank.
3. Continue to fill the tank with water to the desired volume while agitating to promote uniform distribution and prevent settling.

DO NOT MIX WITH OTHER SOLUBLE PESTICIDES OR FERTILIZERS NOT COMPATIBLE WITH MILD ALKALINE SOLUTIONS. THE ADDITION OF SURFACTANT OR WETTING AGENT TO SPRAY IS NOT REQUIRED. PRODUCT IS PRE-FORMULATED WITH THESE PRODUCTS FOR BEST PERFORMANCE. DO NOT ADJUST SPRAY SOLUTION pH AFTER MIXING OF PRODUCT. ACIDIFICATION OF SOLUTION WILL CAUSE REDUCED PRODUCT PERFORMANCE.

#### **SPRAY APPLICATION METHOD**

Do not apply this product through any type of irrigation system. Do not apply directly to soil as a directed spray or drench. For best results, use a properly calibrated sprayer or similar equipment, and check equipment calibration frequently.

#### **CLEANING SPRAY EQUIPMENT**

Clean application equipment thoroughly using a detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

#### **FIELD, INDOOR, GREENHOUSE AND TURF APPLICATION**

(1 oz. package)

Apply at a rate of one to two packages of Agricure for 1000 sq. ft. For maximum treatment rate, use one package of Agricure for 500 sq. ft. Use sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of

development. Do not exceed a use rate of 1 ounce (1 package) of Agricure per 500 sq. ft., per application. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

See special instructions for use on poinsettia, impatiens and pansy.

(4 oz. package)

Apply at a rate of one to two packages of Agricure for 4350 sq/ft (1/10 acre). For maximum treatment rate, use one package of Agricure for 2175 sq. ft (1/20 acre). Use sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 4 ounces (1 package) of Agricure per 2175 sq. ft., per application. Concentration of spray solution and frequency of

16/26

application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

(8 oz. package)

Apply at a rate of one to two packages of Agricure for 8700 sq/ft (2/10 acre). For maximum treatment rate, use one package of Agricure for 4350 sq. ft (1/10 acre). Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the

foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 8 ounces (1 package) of Agricure per 4350 sq. ft., per application. Concentration of spray solution and frequency of

application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

(2.5 lb. package)

Apply at a rate of one to two packages of Agricure per acre. For maximum treatment rate use two packages of Agricure per one acre (43,560 sq. ft.). Use sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 5 pounds of Agricure per acre, per application. Concentration of spray solution and frequency of application will depend upon disease pressure. Do

not  
store unused spray solution. Dispose of unused solution on site.

### **APPLICATION TIMING**

Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, apply Agricure as a post-harvest treatment to remaining foliage to decrease overwintering inoculum.

### **PREHARVEST INTERVAL**

Agricure can be applied up to and including the day of harvest.

**ORNAMENTAL APPLICATIONS:** (California registered uses are listed in a separate section.) Agricure is for use in the following use sites: Greenhouse, Lathhouse, Shadehouse, Commercial and Residential Landscapes, Interiorscapes, Nursery, Container Production and Field Applications.

### **ORNAMENTAL PLANT TYPES**

Agricure is for use on the following plant types: Annual and Perennial Flowers, Bedding Plants, Foliage



Plants, Ground Covers, Woody and Ornamental Shade Trees, Woody and Herbaceous Ornamentals, and Ornamental Nut and Fruit trees

**FOR USE ON:**

Ageratum	Crape Myrtle	Impatiens*	Plum-Ornamental
Ajuga	Crassula	Iris	Poinsettia*
Almond-Ornamental	Croton	Juniper	Poppy
Alyssum	Delphinium	Kalanchoe	Pothos
Andromeda	Dianthus	Lavender	Primrose
Aphelandra	Dizygotheca	Lemon balm	Privet
Apple-Ornamental	Dogwood	Lipstick balm	Protea
Apricot-Ornamental	English Ivy	Lipstick vine	Prune-Ornamental
Artemisia	Episcia	(Aeschynanthus)	Pulmonaria
Aster	Euonymus	Marigold	Pyracantha
Azalea	Fern	Monarda (Bee balm)	Rhododendron
Boxwood	Forsythia	Nectarine-Ornamental	Rose
Calendula	Gazania	Pachysandra	Rose-of-Sharon
Carnation	Geranium	Palm	Schefflera
Centaurea montana	Gladiolus	Pansy*	Snapdragon
Cherry-Ornamental	Gloxinia	Peach-Ornamental	Statice
Chrysanthemum	Gypsophila	Peperomia	Tree Ivy
Cineraria	Hawthorn	Periwinkle	Verbena
Citrus	Heliopsis scabra	Philodendron	Viburnum
Coleus	Hibiscus	Phlox	Violet
Columbine	Holly	Pilea	Zinnia
Coral Bells	Hoya	Pine	Photinia
(Heuchera)	Hydrangea	Pittosporum	Spruce
Begonia	Ginseng	Fir	India-
hawthorn			
Clematis	Daylily	Gerbera daisy	

**DISEASES CONTROLLED:**

Alternaria blight	Fusarium
Anthraxnose	Helminthosporium leaf spot
Ascochyta blight	Ink spot
Blackspot	Phomopsis blight
Botrytis blight	Powdery mildew
Botrytis gray mold	Ovulinia sp.
Cercospora leaf spot	Ramularia leaf spot
Corynespora leaf spot	Ray blight
Didymellina leaf spot	Scab
Diplodia tip blight	Septoria leaf spot
Entomosporium leaf spot	Monilinia fruit and blossom rot
Downy mildew	Penicillium spp.
Rust	

**\*SPECIAL INSTRUCTIONS:**

(1 oz. package)

Poinsettia

For control of intense powdery mildew, apply weekly at a rate of 1 ounce (1 package) per 1000 sq. ft. during vegetative growth prior to bract formation, and apply every 14 days at 1 ounce per 500 sq. ft. thereafter. A rate of 1 ounce per 500 sq. ft applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens

For New Guinea impatiens, apply at a rate of 1 ounce (1 package) per 1000 sq. ft. during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy

Do not use on pansies at early stages of growth.

(4 oz. package)

Poinsettia

For control of intense powdery mildew, apply weekly at a rate of 4 ounces (1 package) per 4350 sq. ft. or 1/10 acre during vegetative growth prior to bract formation, and apply every 14 days at 4 ounces per 2175 sq. ft. thereafter. A rate of 4 ounces per 2175 sq. ft applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens

For New Guinea impatiens, apply at a rate of 4 ounces (1 package) per 4350 sq. ft. or 1/10 acre during vegetative growth; discontinue spraying when blossom buds are visible.

(8 ounce package)

Poinsettia

For control of intense powdery mildew, apply weekly at a rate of 8 ounces (1 package) per 8700 sq. ft. or 1/5 acre during vegetative growth prior to bract formation, and apply every 14 days at 8 ounces per 4350 sq. ft. thereafter. A rate of 8 ounces per 4350 sq. ft applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens

For New Guinea impatiens, apply at a rate of 8 ounces (1 package) per 8700 sq. ft. or 1/5 acre during vegetative growth; discontinue spraying when blossom buds are visible.

(2.5 lb. package)

Poinsettia

For control of intense powdery mildew, apply weekly at a rate of 2.5 pounds (1 package) per acre

19/26

during vegetative growth prior to bract formation, and apply every 14 days at 2.5 pounds per 21,750 sq. ft. or 1/2 acre thereafter. A rate of 2.5 pounds per 21,750 sq. ft applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

#### Impatiens

For New Guinea impatiens, apply at a rate of 2.5 pounds (1 package) per acre during vegetative growth; discontinue spraying when blossom buds are visible.

**AGRICULTURAL APPLICATIONS:** (California registered uses are listed in a separate section.) Agricure is for use in the following use sites: Greenhouse, Orchard, Nursery and Field Applications.

#### **FOR USE ON:**

Acorn squash	Citrus	Okra	Snow peas
Almonds	Crenshaw melon	Oregano	Soybeans
Apples	Cucumber	Peaches	Spinach
Apricots	Dry beans	Pears	Strawberries
Alfalfa	Eggplant	Peas	Sugar beets
Avocado	Endive	Pecans	Summer squash
Barley	Escarole	Peppers	Sweet potatoes
Basil	Grapes	Pistachio	Thyme
Blueberries	Honeydew	Plums	Tomatoes
Broccoli	Hops	Potatoes	Turnip
Cabbage	Kale	Prunes	Walnut
Caneberries	Lettuce	Pumpkin	Watermelon
Cantaloupe	Lima bean	Radicchio	Wheat
Cauliflower	Mango	Rosemary	Winter squash
Cherries	Muskmelon	Sage	Zucchini
Chicory	Mustard greens	Snap beans	Tobacco

#### **DISEASES CONTROLLED:**

Alternaria leaf spot	Fusarium	Scab
Anthraco nose	Phoma	
Botrytis	Phytophthora	
Downy mildew	Powdery mildew	
Penicillium spp.	Monilinia fruit and blossom rot	
Phomopsis blight	Fly speck	
Septoria leaf spot	Cercospora leaf spot	

#### **TURF APPLICATION:**

Agricure is for use in the following use sites: Commercial and Residential Lawns, Golf Course and Other Commercial Turf Grass Sites.

#### **FOR USE ON:**

Bentgrass

20/20

Bermudagrass  
Blue grass, Kentucky, annual  
Centipedegrass  
Fescue, fine, tall, chewings, red  
Ryegrass, perennial, annual  
St. Augustinegrass  
Zoysiagrass

#### **DISEASES CONTROLLED**

Alternaria leaf spot  
Anthracnose  
Ascochyta blight  
Cercospora leaf spot

Downy mildew  
Helminthosporium leaf spot  
Powdery mildew  
Septoria leaf spot

#### **FOR USE ONLY IN CALIFORNIA**

**FOR USE ON:** Acorn squash, Almonds, Apples, Apricots, Cantaloupe, Cherries, Chicory, Crenshaw melon, Cucumber, Endive, Escarole, Grapes, Honeydew, Lettuce, Muskmelon, Peaches, Pears, Peppers, Pistachio, Plums, Prunes, Pumpkin, Radicchio, Rose, Spinach, Strawberries, Summer squash, Tomatoes, Walnut, Watermelon, Winter squash, Zucchini

#### **DISEASES CONTROLLED**

Downy mildew

Powdery mildew

#### **NOTICE**

Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

#### **IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY**

This product has been researched to provide necessary data to support its use on the ornamental, greenhouse and agricultural plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. The user should always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

**NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE.** To the extent allowable by state law the exclusive remedy of the user or Buyer, and the limit of liability of H & I Agritech, Inc. or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of H & I Agritech, Inc.

21/26

**SUBLABEL C- HOUSEHOLD PRODUCTS: FOR NON-COMMERCIAL APPLICATORS  
(NOT SUBJECT TO WORKER PROTECTION STANDARD)**

**AGRICURE**

**FOR ORGANIC PRODUCTION**

**A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER  
DISEASES ON TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS,  
GREENHOUSE AND GARDEN CROPS, AND TURF**

**ACTIVE INGREDIENT:**

Potassium Bicarbonate.....85.00%

OTHER INGREDIENTS.....15.00%

TOTAL.....100.00%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See side (or back) panel for the First Aid Information.

Net weight:       6 oz. (170 g)  
                      8 oz. (227 g)  
                      1 lb. (454 g)  
                      2.5 lbs. (1.14 kg)

Produced For:

H & I AGRITECH, INC.  
95 BROWN ROAD BOX 1030  
ITHACA, NY 14850 - 1257

EPA REG. NO. 70870-1  
EPA EST. NO.

**PRECAUTIONARY STATEMENTS  
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION**

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing  
Wash thoroughly with soap and water after handling.

**FIRST AID**

- If swallowed:
- Call poison control center or doctor immediately for treatment advice.
  - Have person drink several glasses of water.
  - Do not induce vomiting.
  - Do not give anything by mouth to an unconscious person.
- If in eyes:
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.

**HOT LINE NUMBER**

Have a product label or container with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858 7378 between the hours of 6:30 a.m. to 4:30 p.m. Pacific time, seven days a week for emergency medical information.

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters or rinsate.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**If Empty:** Do not reuse this container. Place in trash or offer for recycling if available.

**If Partially Filled:** Call your local solid waste agency or (800) 858-7378 (National Pesticide Information Center) for disposal instructions. Never place unused product down any indoor or outdoor drain.

**GENERAL INFORMATION**

Agricure is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial

and indoor woody and herbaceous ornamental, greenhouse and garden plants and turf.

Agricure is applied with spray equipment using a sufficient volume of water to insure complete coverage of all stems and foliage.

**IMPORTANT:** Read the entire label before using Agricure.

### **MIXING INSTRUCTIONS**

Dilute Agricure with water before applying. Carefully measure and mix the stated amounts of product and water.

Mix 1 to 2 level tablespoons of Agricure per gallon of water. Do not exceed this use rate. Mix the solution thoroughly before use. One gallon of this solution is sufficient to treat approximately 450 square feet. Apply the solution to all exposed surfaces of the plant. Use a sufficient spray volume to obtain complete coverage of all foliage and stems. Uniform and complete coverage is essential for the most effective results.

### **DO NOT STORE UNUSED SPRAY SOLUTIONS**

Spray adjuvants are not required. Some tank mixes with other pesticide products and fertilizers may produce spray mixes that are incompatible. A "jar test" is recommended for unfamiliar tank mixes. As with any pesticide, the addition of spray adjuvants and other pesticides to the spray tank may cause phytotoxicity to the foliage and/or fruit of susceptible crops and plants.

### **APPLICATION TIMING**

Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, Agricure may also be applied post-harvest to remaining foliage to decrease overwintering inoculum.

**ORNAMENTAL APPLICATIONS:** (California registered uses are listed in a separate section.)

### **PLANT TYPES**

Agricure is for: Woody and Ornamental Shade Trees, Woody and Herbaceous Ornamentals, Ornamental Nut and Fruit Trees, Annual and Perennial Flowers, Foliage Plants, Ground Covers and Bedding Plants.

### **FOR USE ON:**

Ageratum	Crape Myrtle	Impatiens	Plum-Ornamental
Ajuga	Crassula	Iris	Poinsettia
Almond-Ornamental	Croton	Juniper	Poppy

24/26

Alyssum	Delphinium	Kalanchoe	Pothos
Andromeda	Dianthus	Lavender	Primrose
Aphelandra	Dizygotheca	Lemon balm	Privet
Apple-Ornamental	Dogwood	Lipstick balm	Protea
Apricot-Ornamental	English Ivy	Lipstick vine	Prune-Ornamental
Artemisia	Episcia	(Aeschynanthus)	Pulmonaria
Aster	Euonymus	Marigold	Pyracantha
Azalea	Fern	Monarda (Bee balm)	Rhododendron
Boxwood	Forsythia	Nectarine-Ornamental	Rose
Calendula	Gazania	Pachysandra	Rose-of-Sharon
Carnation	Geranium	Palm	Schefflera
Centaurea montana	Gladiolus	Pansy	Snapdragon
Cherry-Ornamental	Gloxinia	Peach-Ornamental	Statice
Chrysanthemum	Gypsophila	Peperomia	Tree Ivy
Cineraria	Hawthorn	Periwinkle	Verbena
Citrus	Heliopsis scabra	Philodendron	Viburnum
Coleus	Hibiscus	Phlox	Violet
Columbine	Holly	Pilea	Zinnia
Coral Bells	Hoya	Pine	Photinia
(Heuchera)	Hydrangea	Pittosporum	Spruce
Begonia	Ginseng	Fir	India-
hawthorn			
Clematis	Daylily	Gerbera daisy	

### DISEASES CONTROLLED

Alternaria blight	Fusarium
Anthraxnose	Helminthosporium leaf spot
Ascochyta blight	Ink spot
Blackspot	Phomopsis blight
Botrytis blight	Powdery mildew
Botrytis gray mold	Ovulinia sp.
Cercospora leaf spot	Ramularia leaf spot
Corynespora leaf spot	Ray blight
Didymellina leaf spot	Scab
Diplodia tip blight	Septoria leaf spot
Entomosporium leaf spot	Monilinia fruit and blossom rot
Downy mildew	Penicillium spp.
Rust	

**GARDEN APPLICATIONS:** (California registered uses are listed in a separate section.)

### FOR USE ON:

Acorn squash	Crenshaw melon	Okra	Sage
Almonds	Cucumber	Oregano	Snap beans
Apples	Dry Beans	Peaches	Snow peas



Apricots	Eggplant	Pears	Spinach
Avocado	Endive	Peas	Strawberries
Basil	Escarole	Pecans	Summer squash
Blueberries	Grapes	Peppers	Sweet potatoes
Broccoli	Honeydew	Pistachio	Thyme
Cabbage	Kale	Plums	Tomatoes
Caneberries	Lettuce	Potatoes	Turnip
Cantaloupe	Lima bean	Prunes	Walnut
Cauliflower	Mango	Pumpkin	Watermelon
Cherries	Muskmelon	Radicchio	Winter squash
Chicory	Mustard greens	Rosemary	Zucchini
Citrus			

#### **DISEASES CONTROLLED:**

Alternaria leaf spot	Fusarium
Anthraxnose	Phoma
Botrytis	Phytophthora
Downy mildew	Powdery mildew
Cercospora leaf spot	Fly speck
Penicillium spp.	Monilinia fruit and blossom rot
Phomopsis blight	Scab
Septoria leaf spot	

#### **TURF APPLICATION**

Agricure is for use in the following use sites: Residential lawns and other turf grass sites.

#### **FOR USE ON:**

Bentgrass  
 Bermudagrass  
 Blue grass, Kentucky, annual  
 Centipedegrass  
 Fescue, fine tall, chewings, red  
 Ryegrass, perennial, annual  
 St. Augustinegrass  
 Zoysiagrass

#### **DISEASES CONTROLLED**

Alternaria leaf spot	Downy mildew
Anthraxnose	Helminthosporium leaf spot
Ascochyta blight	Powdery mildew
Cercospora leaf spot	Septoria leaf spot

26/26

**FOR USE ONLY IN CALIFORNIA**

**FOR USE ON:** Acorn squash, Almonds, Apples, Apricots, Cantaloupe, Cherries, Chicory, Crenshaw melon, Cucumber, Endive, Escarole, Grapes, Honeydew, Lettuce, Muskmelon, Peaches, Pears, Peppers, Pistachio, Plums, Prunes, Pumpkin, Radicchio, Rose, Spinach, Strawberries, Summer squash, Tomatoes, Walnut, Watermelon, Winter squash, Zucchini

**DISEASES CONTROLLED**

Downy mildew

Powdery mildew

**NOTICE**

Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

**IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY**

This product has been researched to provide necessary data to support its use on the ornamental, greenhouse and agricultural plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. Always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

**NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE.** To the extent allowable by state law the exclusive remedy of the user or Buyer, and the limit of liability of H & I Agritech, Inc. or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of H & I Agritech, Inc.