



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

May 8, 2015

Leonard Hauptert
Director of Operations
H&I Agritech, Inc.
95 Brown Rd., Box 1030
Ithaca, NY 14850

Subject: Non-PRIA (Pesticide Registration Improvement Act) Labeling Amendment – Add the following commodities to Sub labels A, B and C: celery, crabapples, crabapple-ornamental, dry onion, green onion, hops, lilac, onion, peony, and scallion. Change the maximum app rate for electrostatic sprayers in sub label A from 1.25 lbs, (20 ounces) to 1.125 lbs. (18 ounces). Add the OMRI logo to sub labels A, B and C.
Product Name: Agricure
EPA Registration Number: 70870-1
Application Date: 01/19/2015
OPP Decision Number: 499451

Dear Mr. Hauptert:

The amended labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

At this time, the U.S. Environmental Protection Agency (EPA) is not requiring a change to the restricted-entry interval (REI) of 1 hour as it appears on your labeling in the Agricultural Use Requirements Box. The EPA is currently developing guidance and criteria for when it will allow a REI of less than 4 hours and might require a change to the REI for this product in the future. This approval does not affect any terms or conditions that were previously imposed on this registration. You continue to be subject to existing terms or conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one (1) copy of the final printed labeling before you release this product for shipment with the new labeling. In accordance with 40 CFR § 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR § 152.3.

Should you wish to add/retain a reference to your company's website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the U.S. Environmental Protection Agency (EPA). If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6.

If you have any questions, please contact Shenell Bolden of my team by phone at (703) 347-0120 or via email at bolden.shenell@epa.gov

Sincerely,

A handwritten signature in blue ink that reads "Andrew C. Bryceland". The signature is fluid and cursive, with the first name being the most prominent.

Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

MASTER LABEL

Agricure
FOR ORGANIC PRODUCTION

A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER DISEASES ON
TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS, GREENHOUSE AND AGRICULTURAL
CROPS, AND TURF.

ACTIVE INGREDIENT:

Potassium Bicarbonate.....85.00%
OTHER INGREDIENTS.....15.00%
TOTAL.....100.00%


A: Sublabel – Commercial Products: For Commercial Applicators Subject To Worker
Protection Standard

B: Sublabel – Commercial Products: For Commercial Applicators Subject To Worker
Protection Standard – Water Soluble Packaging Label

C: Sublabel – Household Products: For Non-Commercial Applicators (Not Subject To
Worker Protection Standard)

EPA Reg. No. 70870-1
EPA Establishment No. XXXXX-XX-XXX
Batch Code: ADXXXX

Produced For: H & I Agritech, Inc
95 Brown Rd., Box 1030
Ithaca, NY 14850-1257



Sublabel A-Commercial Products: For Commercial Applicators Subject to Worker Protection Standard

Agricure
(MilStop)

FOR ORGANIC PRODUCTION
A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER DISEASES ON
TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS, GREENHOUSE AND AGRICULTURAL
CROPS, AND TURF

5 pounds can treat up to 1 acre at the recommended treatment rate.

ACTIVE INGREDIENT:
Potassium Bicarbonate.....85.00%
OTHER INGREDIENTS.....15.00%
TOTAL.....100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side (or back) panel for the First Aid Information

Net weight: 5 lbs. (2.27 kg)
 10 lbs. (4.54 kg)
 25 lbs. (11.35 kg)
 50 lbs. (22.7 kg)
 100 lbs. (45.5 kg)

Produced For:
H & I Agritech, Inc.
95 Brown Rd., Box 1030
Ithaca, NY 14850-1257

EPA REG. NO. 70870-1
EPA EST NO.



**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have a product label or container with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 between the hours of 6:30 a.m. to 4:30 p.m. Pacific Time, seven days a week for emergency medical information.	

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters and rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

There is a restricted entry of one (1) hour for this product.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeved shirt and short pants, waterproof gloves and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place. Prevent exposure to moisture. Keep container tightly closed and out of reach of children. Avoid contamination with other pesticides or fertilizers.

PESTICIDE SPILL PROCEDURE: In case of a spill, carefully sweep up the material to avoid generating dust. Store in a labeled container for disposal.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: (For pails with liner) Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or dispose of in a sanitary landfill or by incineration. If pail is contaminated, triple rinse promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container $\frac{1}{4}$ full of water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

(For plastic bags) Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Then offer for recycling, if available, or dispose of empty bag in a sanitary landfill or by incineration. Do not burn unless allowed by state and local ordinances. If burned, stay out of smoke.

GENERAL INFORMATION: Agricure is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental plants, and on greenhouse, nursery and agricultural crops, and on turf. Agricure is applied using a sufficient volume of water to insure complete coverage of all stems and foliage.

NOTICE TO USER: Due to variability within species and in application techniques and equipment, and possible tank mix combinations, neither the manufacturer nor the seller has determined whether Agricure can be safely used on all varieties and species of ornamentals, greenhouse, turf and agricultural plants under all conditions. Therefore, the professional user should determine if Agricure can be used safely before broad usage by applying the prescribed use rate under the conditions expected to be encountered to a small test area. Any adverse effects should be visible within 14 days.

For curative disease control on sensitive crops apply at a rate of 2.5 pounds of Agricure per 100 gallons of water per week and increase the rate each week in 0.25 pound increments. Do not exceed a use rate of 5.0 pounds per acre per application.

IMPORTANT: Read the entire label before using Agricure. Consult your State Agricultural Experiment Station or Extension Service Specialist for additional information on timing of applications, rates, and any additional requirements or restrictions.

MIXING INSTRUCTIONS: Agricure must be diluted with water before applying. Carefully measure and mix the product and water.

1. Fill tank with 1/2 to 3/4 of the selected volume of water prior to the addition of Agricure. Start mechanical or hydraulic agitation.
2. Slowly add the required amount of Agricure to the tank.
3. Continue to fill the tank with water to the desired volume while agitating to promote uniform distribution and prevent settling.

DO NOT MIX WITH OTHER SOLUBLE PESTICIDES OR FERTILIZERS NOT COMPATIBLE WITH MILD ALKALINE SOLUTIONS. THE ADDITION OF SURFACTANT OR WETTING AGENT TO SPRAY IS NOT REQUIRED. PRODUCT IS PRE-FORMULATED WITH THESE PRODUCTS FOR BEST PERFORMANCE. DO NOT ADJUST SPRAY SOLUTION pH AFTER MIXING OF PRODUCT. ACIDIFICATION OF SOLUTION WILL CAUSE REDUCED PRODUCT PERFORMANCE.

SPRAY APPLICATION METHOD: Do not apply this product through any type of irrigation system. Do not apply directly to soil as a directed spray or drench. For best results, use a properly calibrated sprayer or similar equipment, and check equipment calibration frequently.

CLEANING SPRAY EQUIPMENT: Clean application equipment thoroughly using a detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

FIELD APPLICATION: Apply at a rate of 2.0 to 5.0 pounds of Agricure per acre. Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per acre of treatment can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 5 pounds of Agricure per acre, per application. Do not store unused spray solution. Dispose of unused solution on site.

GREENHOUSE APPLICATION: Apply at a rate of 1.25 to 5.0 pounds per 100 gallons of water. Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 0.5 pounds of MilStop per 4350 sq.ft. or 1.15 pounds per 10,000 sq. ft., per application. Do not store unused spray solution. Dispose of unused solution on site.

See special instructions for use on poinsettia, impatiens and pansy.

Small Volume Application Chart: Use the chart below to determine Agricure quantities for small area applications. Adjust to desired concentration for low disease pressure or as a curative treatment.

Chart Treatment Rate: 2.5 lbs. per 100 gallons

Amount of Agricure	Amount of Water	Square feet treated
1 Tbsp	1 gallon	500
2 Tbsp	2 gallons	1,000
4 Tbsp	4 gallons	2,000
1/8 cup	2 gallons	1,000
1/4 cup	4 gallons	2,000
1/2 cup	8 gallons	4,000
1 oz. (28 grams)	2.5 gallons	1,100
2 oz. (57 grams)	5 gallons	2,200
4 oz. (113 grams)	10 gallons	4,400

ELECTROSTATIC SPRAYERS: Apply at a rate of 0.625 pounds (10 ounces) to 1.125 pounds (18 ounces) per 2 gallons of water per 10,000 sq. ft. Do not exceed a rate of 1.5 pounds (24 ounces) per 2 gallons of water.

THERMAL FOGGERS: Apply at a rate of 0.38 pounds (6 ounces) to 0.75 pounds (12 ounces) per 1 gallon of water per 10,000 sq. ft. Use the 0.38 pounds (6 ounces) per 1 gallon of water rate for plants that are in flower. Do not exceed 0.75 pounds (12 ounces) per 1 gallon of water.

Turf Application: Apply at a rate of 2.5 to 5 pounds of Agrisure per acre. Use a sufficient volume of spray solution to obtain complete coverage of foliage. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per acre of treatment can vary depending upon stage of development. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not exceed a use rate of 5 pounds of Agrisure per acre, per application. Do not store unused spray solution. Dispose of unused solution of site.

APPLICATION TIMING: Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, apply Agrisure as a post-harvest treatment to remaining foliage to decrease overwintering inoculum.

PREHARVEST INTERVAL: Agrisure can be applied up to and including the day of harvest.

ORNAMENTAL APPLICATIONS: Agrisure is for use in the following use sites: Greenhouse, Lathhouse, Shadehouse, Commercial and Residential Landscapes, Interiorscapes, Nursery, Container Production and Field Applications.

ORNAMENTAL PLANT TYPES: Agrisure is for use on the following plant types: Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, Woody and Ornamental Shade Trees, Woody and Herbaceous Ornamentals, and Ornamental Nut and Fruit Trees.

FOR USE ON: Ageratum, Ajuga, Almond-Ornamental, Alyssum, Andromeda, Aphelandra, Apple-Ornamental, Apricot-Ornamental, Artemisia, Aster, Azalea, Begonia¹, Boxwood, Calendula, Carnation, Centaurea montana, Cherry-Ornamental, Chrysanthemum, Cineraria, Citrus, Coleus, Columbine, Coral Bells (Heuchera), Clematis¹, Crabapple-Ornamental¹, Crape Myrtle, Crassula, Croton, Daylily¹, Delphinium, Dianthus, Dizygotheca, Dogwood, English Ivy, Episcia, Euonymus, Fern, Fir¹, Forsythia, Gazania, Geranium, Gerbera daisy¹, Ginseng¹, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Heliopsis scabra, Hibiscus, Holly, Hoya, Hydrangea, Impatiens*, India hawthorn¹, Iris, Juniper, Kalanchoe, Lavender, Lemon balm, Lilac¹ Lipstick balm, Lipstick Vine (Aeschynanthus), Marigold, Monarda (Bee balm), Nectarine-Ornamental, Pachysandra, Palm, Pansy*, Peach-Ornamental, Peony¹, Peperomia, Periwinkle, Philodendron, Phlox, Photinia, Pilea, Pine, Pittosporum, Plum-Ornamental, Poinsettia*, Poppy, Pothos, Primrose, Privet, Protea, Prune-Ornamental, Pulmonaria, Pyracantha, Rhododendron, Rose, Rose-of-Sharon, Schefflera, Snapdragon, Spruce¹, Statice, Tree Ivy, Verbena, Viburnum, Violet, Zinnia.

¹Not For Use In California

DISEASES CONTROLLED: Alternaria blight, Anthracnose, Ascochyta blight, Blackspot, Botrytis blight, Botrytis gray mold, Cercospora leaf spot, Corynespora leaf spot, Didymellina leaf spot, Diplodia tip blight, Downy mildew¹, Entomosporium leaf spot, Fusarium, Helminthosporium leaf spot, Ink spot, Monilinia fruit and blossom rot¹, Ovulinia sp., Penicillium spp. ¹, Phomopsis blight, Powdery mildew, Ramularia leaf spot, Ray blight, Rust¹, Scab, Septoria leaf spot.

¹Not For Use In California

***SPECIAL INSTRUCTIONS:** Poinsettia - For control of intense powdery mildew, apply weekly at a rate of 0.25 pounds per 4350 sq. ft. or 1/10 acre during vegetative growth prior to bract formation, and apply

every 14 days at 0.5 pounds per 4350 sq. ft. thereafter. A rate of 0.5 pounds per 4350 sq. ft applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens - For New Guinea impatiens, apply at a rate of 0.25 pounds per 4350 sq. ft. or 1/10 acre during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy - Do not use on pansies at early stages of growth.

AGRICULTURAL APPLICATIONS: Agricure is for use in the following use sites: Greenhouse, Orchard, Nursery and Field Applications.

FOR USE ON: Acorn squash, Alfalfa, Almonds, Apples, Apricots, Avocado, Barley, Basil, Blueberries, Broccoli, Cabbage, Caneberries, Cantaloupe, Cauliflower, Celery¹, Cherries, Chicory, Citrus, Crabapple¹, Crenshaw melon, Cucumber, Dry Beans, Dry Onion¹, Eggplant, Endive, Escarole, Grapes, Green Onion¹, Honeydew, Hops, Kale, Lettuce, Lima bean, Mango, Muskmelon, Mustard greens, Okra, Onion¹, Oregano, Peaches, Pears, Peas, Pecans, Peppers, Pistachio, Plums, Potatoes, Prunes, Pumpkin, Radicchio, Rosemary, Sage, Scallion¹, Snap beans, Snow peas, Soybeans, Spinach, Strawberries, Sugar Beets, Summer squash, Sweet potatoes, Thyme, Tobacco¹, Tomatoes, Turnip, Walnut, Watermelon, Wheat, Winter squash, Zucchini.

¹Not For Use In California

DISEASES CONTROLLED: Alternaria leaf spot, Anthracnose, Botrytis, Cercospora leaf spot¹, Downy mildew, Fly speck¹, Fusarium, Monilinia fruit and blossom rot¹, Penicillium spp.¹, Phoma, Phomopsis blight¹, Phytophthora, Powdery mildew, Scab¹, Septoria leaf spot¹

¹Not For Use In California

Turf Application: (Not for use in California) Agricure is for use in the following use sites: Commercial and residential lawns, golf course and other commercial turf grass sites.

For Use On: Bentgrass, Bermudagrass, Bluegrass (Kentucky, annual), Centipedegrass, Fescue (fine, tall, chewings, red), Ryegrass (perennial, annual), St. Augustinegrass, Zoysiagrass

Diseases Controlled: Alternaria leaf spot, Anthracnose, Ascochyta blight, Brown blight, Cercospora leaf spot, Dollar spot, Downy mildew, Helminthosporium leaf spot, Powdery mildew, Rust, Septoria leaf spot.

NOTICE

Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY: This product has been researched to provide necessary data to support its use on the ornamental, greenhouse and agricultural plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. Always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. To the extent allowable by state law the exclusive remedy of the user or Buyer, and the limit of liability of H & I Agritech, Inc. or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of H & I Agritech, Inc.

Sublabel B - Commercial Products: For Commercial Applicators Subject to Worker Protection Standard – Water Soluble Packaging Label

Agricure Water Soluble Package

FOR ORGANIC PRODUCTION
A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER DISEASES ON
TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS, GREENHOUSE AND AGRICULTURAL
CROPS

This package can treat up to ___ sq. ft. (acres) at the recommended maximum treatment rate.

ACTIVE INGREDIENT:

Potassium Bicarbonate.....	85.00%
OTHER INGREDIENTS.....	15.00%
TOTAL.....	100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See side (or back) panel for the First Aid Information

Net weight:

1.0 oz.	(28.35 g)
4.0 oz.	(113.4 g)
8.0 oz.	(227 g)
2.5 lbs.	(1.14 kg)

Produced For:
H & I Agritech, Inc.
95 Brown Rd., Box 1030
Ithaca, NY 14850-1257

EPA REG. NO. 70870-1
EPA EST NO.



**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have a product label or container with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 between the hours of 6:30 a.m. to 4:30 p.m. Pacific Time, seven days a week for emergency medical information.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- shoes plus socks

Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters and rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

There is a restricted entry of one (1) hour for this product.

PPE requirement for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeved shirt and short pants, waterproof gloves and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product not within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treated areas until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place. Prevent exposure to moisture. Keep container tightly closed and out of reach of children. Avoid contamination with other pesticides or fertilizers.

PESTICIDE SPILL PROCEDURE: In case of a spill, carefully sweep up the material to avoid generating dust. Store in a labeled container for disposal.

PESTICIDE DISPOSAL: Wastes resulting from use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available.

GENERAL INFORMATION: Agricure is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental plants, and on greenhouse, nursery and agricultural crops, and turf. Agricure is applied using a sufficient volume of water to insure complete coverage of all stems and foliage.

NOTICE TO USER: Due to variability within species and in application techniques and equipment, and possible tank mix combinations, neither the manufacturer nor the seller has determined whether Agricure can be safely used on all varieties and species of ornamentals, greenhouse, and agricultural plants under all conditions. Therefore, the professional user should determine if Agricure can be used safely before broad usage by applying the prescribed use rate under the conditions expected to be encountered to a small test area. Any adverse effects should be visible within 14 days.

(1 ounce package)

For curative disease control on sensitive crops apply at a rate of 1 ounce (1 package) of Agricure per 2.5 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 1 ounce per 500 sq. ft. per week.

(4 ounce package)

For curative disease control on sensitive crops apply at a rate of 4 ounces (1 package) of Agricure per 10 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 4 ounces per 2175 sq. ft. per application.

(8 ounce package)

For curative disease control on sensitive crops apply at a rate of 8 ounces (1 package) of Agricure per 20 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 4 ounces per 4350 sq. ft. or 1/10 acre per application.

(2.5 pound package)

For curative disease control on sensitive crops apply at a rate of 2.5 pounds (1 package) of Agricure per 100 gallons of water per week and increase the rate for more complete control, if needed. Do not exceed a use rate of 2.5 pounds per 21,750 sq. ft. or 1/2 acre per application.

IMPORTANT: Read the entire label before using Agricure. Consult your State Agricultural Experiment Station or Extension Service Specialist for additional information on timing of applications, rates, and any additional requirements or restrictions.

MIXING INSTRUCTIONS: Agricure must be diluted with water before applying. Carefully measure and mix the product and water.

1. Fill tank with 1/2 to 3/4 of the selected volume of water prior to the addition of Agricure. Start mechanical or hydraulic agitation.
2. Slowly add the required number of packages of Agricure to the tank.
3. Continue to fill the tank with water to the desired volume while agitating to promote uniform distribution and prevent settling.

DO NOT MIX WITH OTHER SOLUBLE PESTICIDES OR FERTILIZERS NOT COMPATIBLE WITH MILD ALKALINE SOLUTIONS. THE ADDITION OF SURFACTANT OR WETTING AGENT TO SPRAY IS NOT REQUIRED. PRODUCT IS PRE-FORMULATED WITH THESE PRODUCTS FOR BEST PERFORMANCE. DO NOT ADJUST SPRAY SOLUTION pH AFTER MIXING OF PRODUCT. ACIDIFICATION OF SOLUTION WILL CAUSE REDUCED PRODUCT PERFORMANCE.

SPRAY APPLICATION METHOD: Do not apply this product through any type of irrigation system. Do not apply directly to soil as a directed spray or drench. For best results, use a properly calibrated sprayer or similar equipment, and check equipment calibration frequently.

CLEANING SPRAY EQUIPMENT: Clean application equipment thoroughly using a detergent or commercial sprayer cleaner according to the manufacturer's directions before and after applying this product.

FIELD, INDOOR AND TURF APPLICATION

(1 oz. package)

Apply at a rate of one to two packages of Agricure for 1000 sq. ft. For maximum treatment rate, use one package of Agricure for 500 sq. ft. Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 1 ounce (1 package) of Agricure per 500 sq. ft., per application. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

See special instructions for use on poinsettia, impatiens and pansy.

(4 oz. package)

Apply at a rate of one to two packages of Agricure for 4350 sq/ft (1/10 acre). For maximum treatment rate, use one package of Agricure for 2175 sq. ft. (1/20 acre). Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 4 ounces (1 package) of Agricure per 2175 sq. ft., per application. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

(8 oz. package)

Apply at a rate of one to two packages of Agricure for 8700 sq/ft (2/10 acre). For maximum treatment rate, use one package of Agricure for 4350 sq. ft. (1/10 acre). Use a sufficient volume of spray solution

to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 8 ounces (1 package) of Agricure per 4350 sq. ft., per application. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

(2.5 lb. package)

Apply at a rate of one to two packages of Agricure per acre. For maximum treatment rate, use two packages of Agricure for one acre (43,560 sq. ft.). Use a sufficient volume of spray solution to obtain complete coverage of foliage and stems. Uniform and complete coverage of the foliage is essential for the most effective results. Spray volume per unit of treated area can vary depending upon plant type and stage of development. Do not exceed a use rate of 5 pounds of Agricure per acre, per application. Concentration of spray solution and frequency of application will depend upon disease pressure. Do not store unused spray solution. Dispose of unused solution on site.

APPLICATION TIMING: Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, apply Agricure as a post-harvest treatment to remaining foliage to decrease overwintering inoculum.

PREHARVEST INTERVAL: Agricure can be applied up to and including the day of harvest.

ORNAMENTAL APPLICATIONS: Agricure is for use in the following use sites: Greenhouse, Lathhouse, Shadehouse, Commercial and Residential Landscapes, Interiorscapes, Nursery, Container Production and Field Applications.

ORNAMENTAL PLANT TYPES: Agricure is for use on the following plant types: Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, Woody and Ornamental Shade Trees, Woody and Herbaceous Ornamentals, and Ornamental Nut and Fruit Trees.

FOR USE ON: Ageratum, Ajuga, Almond-Ornamental, Alyssum, Andromeda, Aphelandra, Apple-Ornamental, Apricot-Ornamental, Artemisia, Aster, Azalea, Begonia¹, Boxwood, Calendula, Carnation, Centaurea montana, Cherry-Ornamental, Chrysanthemum, Cineraria, Citrus, Coleus, Columbine, Coral Bells (Heuchera), Clematis¹, Crabapple-Ornamental¹, Crape Myrtle, Crassula, Croton, Daylily¹, Delphinium, Dianthus, Dizygotheca, Dogwood, English Ivy, Episcia, Euonymus, Fern, Fir¹, Forsythia, Gazania, Geranium, Gerbera daisy¹, Ginseng¹, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Heliopsis scabra, Hibiscus, Holly, Hoya, Hydrangea, Impatiens*, India hawthorn¹, Iris, Juniper, Kalanchoe, Lavender, Lemon balm, Lilac¹, Lipstick balm, Lipstick Vine (Aeschynanthus), Marigold, Monarda (Bee balm), Nectarine-Ornamental, Pachysandra, Palm, Pansy*, Peach-Ornamental, Peony¹, Peperomia, Periwinkle, Philodendron, Phlox, Photinia, Pilea, Pine, Pittosporum, Plum-Ornamental, Poinsettia*, Poppy, Pothos, Primrose, Privet, Protea, Prune-Ornamental, Pulmonaria, Pyracantha, Rhododendron, Rose, Rose-of-Sharon, Schefflera, Snapdragon, Spruce¹, Statice, Tree Ivy, Verbena, Viburnum, Violet, Zinnia.

¹Not For Use In California

DISEASES CONTROLLED: Alternaria blight, Anthracnose, Ascochyta blight, Blackspot, Botrytis blight, Botrytis gray mold, Cercospora leaf spot, Corynespora leaf spot, Didymellina leaf spot, Diplodia tip blight, Downy mildew¹, Entomosporium leaf spot, Fusarium, Helminthosporium leaf spot, Ink spot, Monilinia fruit and blossom rot¹, Ovulinia sp., Penicillium spp. ¹, Phomopsis blight, Powdery mildew, Ramularia leaf spot, Ray blight, Rust¹, Scab, Septoria leaf spot.

¹Not For Use In California

***SPECIAL INSTRUCTIONS:**

(1 oz. package)

Poinsettia - For control of intense powdery mildew, apply weekly at a rate of 1 ounce (1 package) per 1000 sq. ft. during vegetative growth prior to bract formation, and apply every 14 days at 1 ounce per 500 sq. ft. thereafter. A rate of 1 ounce per 500 sq. ft. applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens - For New Guinea impatiens, apply at a rate of 1 ounce (1 package) per 1000 sq. ft. during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy - Do not use on pansies at early stages of growth.

(4 oz. package)

Poinsettia - For control of intense powdery mildew, apply weekly at a rate of 4 ounces (1 package) per 4350 sq. ft. or 1/10 acre during vegetative growth prior to bract formation, and apply every 14 days at 4 ounces per 2175 sq. ft. thereafter. A rate of 4 ounces per 2175 sq. ft. applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens - For New Guinea impatiens, apply at a rate of 4 ounces (1 package) per 4350 sq. ft. or 1/10 acre during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy - Do not use on pansies at early stages of growth.

(8 oz. package)

Poinsettia - For control of intense powdery mildew, apply weekly at a rate of 8 ounces (1 package) per 8700 sq. ft. or 1/5 acre during vegetative growth prior to bract formation, and apply every 14 days at 8 ounces per 4350 sq. ft. thereafter. A rate of 8 ounces per 4350 sq. ft. applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens - For New Guinea impatiens, apply at a rate of 8 ounces (1 package) per 8700 sq. ft. or 1/5 acre during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy - Do not use on pansies at early stages of growth.

(2.5 lb. package)

Poinsettia - For control of intense powdery mildew, apply weekly at a rate of 2.5 pounds (1 package) per acre during vegetative growth prior to bract formation, and apply every 14 days at 2.5 pounds per 21,750 sq. ft. or 1/2 acre thereafter. A rate of 2.5 pounds per 21,750 sq. ft. applied weekly during vegetative growth has the potential to cause marginal burn to newly developing leaves, but does not injure developing bracts when applied biweekly.

Impatiens - For New Guinea impatiens, apply at a rate of 2.5 pounds (1 package) per acre during vegetative growth; discontinue spraying when blossom buds are visible.

Pansy - Do not use on pansies at early stages of growth.

AGRICULTURAL APPLICATIONS: Agricure is for use in the following use sites: Greenhouse, Orchard, Nursery and Field Applications.

FOR USE ON: Acorn squash, Alfalfa, Almonds, Apples, Apricots, Avocado, Barley, Basil, Blueberries, Broccoli, Cabbage, Caneberries, Cantaloupe, Cauliflower, Celery¹, Cherries, Chicory, Citrus, Crabapple¹, Crenshaw melon, Cucumber, Dry Beans, Dry Onion¹, Eggplant, Endive, Escarole, Grapes, Green Onion¹, Honeydew, Hops, Kale, Lettuce, Lima bean, Mango, Muskmelon, Mustard greens, Okra, Onion¹, Oregano, Peaches, Pears, Peas, Pecans, Peppers, Pistachio, Plums, Potatoes, Prunes, Pumpkin, Radicchio, Rosemary, Sage, Scallion¹, Snap beans, Snow peas, Soybeans, Spinach, Strawberries, Sugar

Beets, Summer squash, Sweet potatoes, Thyme, Tobacco¹, Tomatoes, Turnip, Walnut, Watermelon, Wheat, Winter squash, Zucchini.

¹Not For Use In California

DISEASES CONTROLLED: Alternaria leaf spot, Anthracnose, Botrytis, Cercospora leaf spot¹, Downy mildew, Fly speck¹, Fusarium, Monilinia fruit and blossom rot¹, Penicillium spp.¹, Phoma, Phomopsis blight¹, Phytophthora, Powdery mildew, Scab¹, Septoria leaf spot¹

¹Not For Use In California

Turf Application: (Not for use in California) Agricure is for use in the following use sites: Commercial and residential lawns. Golf course and other commercial turf grass sites.

For Use On: Bentgrass, Bermudagrass, Bluegrass (Kentucky, annual), Centipedegrass, Fescue (fine, tall, chewings, red), Ryegrass (perennial, annual), St. Augustinegrass, Zoysiagrass

Diseases Controlled: Alternaria leaf spot, Anthracnose, Ascochyta blight, Brown blight, Cercospora leaf spot, Dollar spot, Downy mildew, Helminthosporium leaf spot, Powdery mildew, Rust, Septoria leaf spot.

NOTICE

Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY: This product has been researched to provide necessary data to support its use on the ornamental, greenhouse and agricultural plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. Always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. To the extent allowable by state law the exclusive remedy of the user or Buyer, and the limit of liability of H & I Agritech, Inc. or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of H & I Agritech, Inc.

Sublabel C – Household Products: For Non-Commercial Applicators (Not Subject to Worker Protection Standard)

Agricure
(Greencure)

FOR ORGANIC GARDENING

A FUNGICIDE FOR THE CONTROL OF POWDERY MILDEW AND OTHER DISEASES ON TERRESTRIAL AND INDOOR ORNAMENTAL PLANTS, GREENHOUSE AND GARDEN CROPS, AND TURF

ACTIVE INGREDIENT:

Potassium Bicarbonate.....	85.00%
OTHER INGREDIENTS.....	15.00%
TOTAL.....	100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

See side (or back) panel for the First Aid Information

Net weight:

6 oz.	(170 g)
8 oz.	(227 g) (Makes up to 16 gallons of spray)
1 lb.	(454 g)
2.5 lbs.	(22.7 kg)(Makes up tp 80 gallons of spray)

Produced For:
H & I Agritech, Inc.
95 Brown Rd., Box 1030
Ithaca, NY 14850-1257

EPA REG. NO. 70870-1
EPA EST NO.



**PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid	
If swallowed:	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting.• Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
HOT LINE NUMBER	
Have a product label or container with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 between the hours of 6:30 a.m. to 4:30 p.m. Pacific Time, seven days a week for emergency medical information.	

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water. Do not contaminate water when cleaning equipment or disposing of equipment washwaters and rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

If Empty: Nonrefillable container. Do not use or refill this container. Offer for recycling, if available.

If Partially Filled: Call your local solid waste agency or (800) 858-7378 (National Pesticide Information Center for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION: Agrisure is a broad spectrum, contact foliar fungicide that controls a variety of diseases on terrestrial and indoor woody and herbaceous ornamental, greenhouse and garden plants, and turf. Agrisure is applied using a sufficient volume of water to insure complete coverage of all stems and foliage.

IMPORTANT: Read the entire label before using Agrisure.

MIXING INSTRUCTIONS: Dilute Agrisure with water before applying. Carefully measure and mix the stated amounts of product and water.

Mix 1 to 2 level tablespoons of Agrisure per gallon of water. Do not exceed this rate. Mix the solution thoroughly before use. One gallon of this solution is sufficient to treat approximately 450 square feet. Apply the solution to all exposed surfaces of the plant. Use a sufficient spray volume to obtain complete coverage of all foliage and stems. Uniform and complete coverage is essential for the most effective results. Do not store unused spray solutions.

Spray adjuvants are not required. Some tank mixes with other pesticide products and fertilizers may produce spray mixes that are incompatible. A "jar test" is recommended for unfamiliar tank mixes. As with any pesticide, the addition of spray adjuvants and other pesticides to the spray tank may cause phytotoxicity to the foliage and/or fruit of susceptible crops and plants.

APPLICATION TIMING: Start application at first sign of disease. For best protection, repeat at one to two week intervals until conditions are no longer favorable for disease development. Shorten the interval during rainy weather or during periods of high relative humidity.

For appropriate crops, apply Agrisure as a post-harvest treatment to remaining foliage to decrease overwintering inoculum.

ORNAMENTAL APPLICATIONS: (California registered uses are listed in a separate section.)

PLANT TYPES: Agrisure is for: Woody and Ornamental Shade Trees, Woody and Herbaceous Ornamentals, Ornamental Nut and Fruit Trees, Annual and Perennial Flowers, Foliage Plants, Ground Covers and Bedding Plants.

FOR USE ON: Ageratum, Ajuga, Almond-Ornamental, Alyssum, Andromeda, Aphelandra, Apple-Ornamental, Apricot-Ornamental, Artemisia, Aster, Azalea, Begonia, Boxwood, Calendula, Carnation, Centaurea montana, Cherry-Ornamental, Chrysanthemum, Cineraria, Citrus, Coleus, Columbine, Coral Bells (Heuchera), Clematis, Crabapple-Ornamental¹, Crape Myrtle, Crassula, Croton, Daylily, Delphinium, Dianthus, Dizygotheca, Dogwood, English Ivy, Episcia, Euonymus, Fern, Fir, Forsythia, Gazania, Geranium, Gerbera daisy, Ginseng, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Heliopsis scabra, Hibiscus, Holly, Hoya, Hydrangea, Impatiens, India hawthorn, Iris, Juniper, Kalanchoe, Lavender, Lemon balm, Lilac¹, Lipstick balm, Lipstick Vine (Aeschynanthus), Marigold, Monarda (Bee balm), Nectarine-Ornamental, Pachysandra, Palm, Pansy, Peach-Ornamental, Peony¹, Peperomia, Periwinkle, Philodendron, Phlox, Photinia, Pilea, Pine, Pittosporum, Plum-Ornamental, Poinsettia, Poppy, Pothos, Primrose, Privet, Protea, Prune-Ornamental, Pulmonaria, Pyracantha, Rhododendron, Rose, Rose-of-Sharon, Schefflera, Snapdragon, Spruce, Statice, Tree Ivy, Verbena, Viburnum, Violet, Zinnia.

DISEASES CONTROLLED: Alternaria blight, Anthracnose, Ascochyta blight, Blackspot, Botrytis blight, Botrytis gray mold, Cercospora leaf spot, Corynespora leaf spot, Didymellina leaf spot, Diplodia tip blight, Downy mildew, Entomosporium leaf spot, Fusarium, Helminthosporium leaf spot, Ink spot, Monilinia fruit and blossom rot, Ovinia sp., Penicillium spp., Phomopsis blight, Powdery mildew, Ramularia leaf spot, Ray blight, Rust, Scab, Septoria leaf spot.

GARDEN APPLICATIONS: (California registered uses are listed in a separate section.)

FOR USE ON: Acorn squash, Almonds, Apples, Apricots, Avocado, Basil, Blueberries, Broccoli, Cabbage, Caneberries, Cantaloupe, Cauliflower, Celery, Cherries, Chicory, Citrus, Crabapple, Crenshaw melon, Cucumber, Dry Beans, Eggplant, Endive, Escarole, Grapes, Honeydew, Hops, Kale, Lettuce, Lima bean, Mango, Muskmelon, Mustard greens, Okra, Onion, Oregano, Peaches, Pears, Peas, Pecans, Peppers, Pistachio, Plums, Potatoes, Prunes, Pumpkin, Radicchio, Rosemary, Sage, Snap beans, Snow peas, Spinach, Strawberries, Summer squash, Sweet potatoes, Thyme, Tomatoes, Turnip, Walnut, Watermelon, Winter squash, Zucchini.

DISEASES CONTROLLED: Alternaria leaf spot, Anthracnose, Botrytis, Cercospora leaf spot, Downy mildew, Fly speck, Fusarium, Monilinia fruit and blossom rot, Penicillium spp., Phoma, Phomopsis blight, Phytophthora, Powdery mildew, Scab, Septoria leaf spot

Turf Application: Agrisure is for use in the following use sites: Residential lawns and other turf grass sites.

For Use On: Bentgrass, Bermudagrass, Bluegrass (Kentucky, annual), Centipedegrass, Fescue (fine, tall, chewings, red), Ryegrass (perennial, annual), St. Augustinegrass, Zoysiagrass

Diseases Controlled: Alternaria leaf spot, Anthracnose, Ascochyta blight, Brown blight, Cercospora leaf spot, Dollar spot, Downy mildew, Helminthosporium leaf spot, Powdery mildew, Rust, Septoria leaf spot.

FOR USE ONLY IN CALIFORNIA

FOR USE ON: Acorn squash, Almonds, Apples, Apricots, Cantaloupe, Cherries, Chicory, Crenshaw Melon, Cucumber, Endive, Escarole, Grapes, Honeydew, Lettuce, Muskmelon, Peaches, Pears, Peppers, Pistachio, Plums, Prunes, Pumpkin, Radicchio, Rose, Spinach, Strawberries, Summer squash, Tomatoes, Walnut, Watermelon, Winter squash, Zucchini

Diseases Controlled: Downy mildew, Powdery mildew

NOTICE

Buyer assumes all responsibility for safety if use is not in accordance with label instructions.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY: This product has been researched to provide necessary data to support its use on the ornamental, greenhouse and agricultural plants listed on the label. However, it is understood that tests may not have been carried out on all varieties or cultivars and under all growing conditions on all plants listed on the label. Always follow the label directions. Exercise care when using this product on a given variety until familiar with the results under your growing conditions.

NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. To the extent allowable by state law the exclusive remedy of the user or Buyer, and the limit of liability of H & I Agritech, Inc. or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of H & I Agritech, Inc.