

70907-12

11/18/2003

1/56



ORACLE DICAMBA

AGRICULTURAL HERBICIDE

FOR WEED CONTROL IN CORN, SORGHUM, SMALL GRAINS, PASTURE, HAY RANGELAND, GENERAL FARMSTEAD (Non-Cropland), RIGHTS-OF-WAY, PUBLIC UTILITY AND INDUSTRIAL AREAS, FALLOW, SUGARCANE, ASPARAGUS, TURF AND GRASS SEED CROPS.

Active Ingredient:

Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid)*.....49.77%

Inert Ingredients:.....50.23%

TOTAL 100.00%

* This product contains 41.35% 3,6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon (480 g/L)

KEEP OUT OF REACH OF CHILDREN

WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See side panel for additional precautionary statements.

EPA Reg. No. 70907-12
EPA Est. No. 2217-KS-1
Net Contents: 2.5 gal.

Gharda USA, Inc.
Newtown, PA 18940

ACCEPTED
In EPA Letter Dated

NOV 18 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

70907-12

PRECAUTIONARY STATEMENT
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING. Causes eye irritation. Do not get in eyes, on skin, or on clothing.
 Harmful if swallowed. Avoid breathing spray mist. Wash thoroughly after handling.

FIRST AID (Substituted Benzoic acid)	
If inhaled:	▪ Remove person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. ▪ Call a poison control center or doctor for further treatment advice.
If on Skin or clothing:	▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes. ▪ Call a poison control center or doctor for treatment advise.
If In eyes:	▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control center or doctor for treatment advise.
If swallowed:	▪ Call poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by the poison control center or doctor. ▪ Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER (Substituted Benzoic acid) Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency medical treatment information call: 1-(866)-359-5660	
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage.	

Personal Protective Equipment

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations:

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

Apply this product only as directed on label.

...

...

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

For any requirements specific to your State or tribe, consult the agency responsible for pesticide regulation. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks
- Protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons, including children and pets, out of the treated areas until sprays have dried.

Before applying ORACLE, read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury, or illegal residues.

GENERAL INFORMATION

The following directions apply to all uses of ORACLE. Additional precautions and restrictions will be found in each specific use section.

Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

Do not apply this product through any type of irrigation system.

MIXING AND APPLICATION

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF ORACLE. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

ORACLE is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (see COMPATIBILITY TEST on page _) should be made prior to tank mixing.

Ground or aerial application equipment which will give good spray coverage of weed foliage should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of a diluted spray per treated acre when using ground application equipment, or 1 to 10 gallons of a diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, ORACLE should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g. cultivating or mowing) treated areas for at least 7 days following application.

BEST STEWARDSHIP PRACTICES

ORACLE provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

GROUND AND SURFACE WATERS PROTECTION

1) Point source contamination - To prevent point source contamination, do not mix, load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment, or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or antisiphoning devices must be used on all mixing equipment.

2) Movement by surface runoff or through soil - Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow. To minimize the possibility of

ground water contamination, carefully follow application rate recommendations as affected by soil type in the general information section of this label.

3) Movement by water erosion of treated soil - Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

SENSITIVE CROP PRECAUTIONS

ORACLE may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes and other broadleaf plants when contacting their roots, stems, or foliage. These plants are most sensitive to ORACLE during their development or growing stage. FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING ORACLE.

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of ORACLE with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when spray particles may be carried by air currents to areas where sensitive crops and plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays to avoid potential herbicide drift. Select nozzles which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground applications are Delavan Raindrops, Spraying Systems XR flat fans or large capacity flood nozzles such as D10, TK10 or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 gpa, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult with your spray nozzle supplier concerning the choice of drift reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- Do not apply ORACLE adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.

- To avoid injury to desirable plants, equipment used to apply ORACLE should be thoroughly cleaned (see PROCEDURE FOR CLEANING SPRAY EQUIPMENT on page 9) before reusing to apply any other chemicals.

All crop uses of ORACLE are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. **Tank mix recommendations are for use only in states where the tank mix product and application site are registered.**

BAND TREATMENTS

ORACLE may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE treated acre} = \text{Band RATE per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per treated acre} = \text{Band VOLUME per treated acre}$$

COMPATIBILITY TEST

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

**Amount of Herbicide to Add to One Pint of Spray Carrier
(Assuming Volume is 25 Gallons per Acre)**

HERBICIDE FORMULATIONS	RATE PER ACRE	LEVEL TEASPOONS
Dry	1 lb.	1 1/2
Liquid	1 pt.	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Re-run the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

PROCEDURE FOR CLEANING SPRAY EQUIPMENT

The steps listed below are suggested for thorough cleaning of spray equipment following applications of ORACLE or tank mixes of ORACLE or tank mixes of ORACLE plus 2,4-D amine.

- 1) Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- 2) Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- 3) Flush the solution out of the spray tank through the boom.
- 4) Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply ORACLE as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. ORACLE tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

- 5) Complete step 1.
- 6) Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5

to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.

7) Flush the detergent solution out of the spray tank through the boom.

8) Repeat step 1, and follow with steps 2, 3, and 4.

GENERAL WEED LIST

This is a general list of weeds which may be treated with ORACLE in accordance with this label as recommended under the rates and timing sections of the Individual Use Headings. Proper usage of this product will give control or growth suppression of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species including:

ANNUALS

Amaranth, Spiny (Spiny Pigweed)

Aster, Slender

Bedstraw

Beggarweed, Florida

Broomweed, Common

Buckwheat, Wild

Buffalobur

Burclover, California

Burcucumber

Buttercup, Roughseed

Carpetweed

Catchfly, Nightflowering

Chamomile, Corn

Chickweed, Common

Clovers (Annual)

Cockle, Corn

Cockle, Cow

Cocklebur, Common

Croton, Tropic

Croton, Woolly

Daisy, English

Evening primrose, Cutleaf

Fleabane, Annual

Goosefoot, Nettleleaf

Henbit

Jimsonweed

Knotweed

Kochia

Ladysthumb
Lambsquarters, Common
Lambsquarters (triazine resistant)
Lettuce, Prickly
Mallow, Common
Mallow, Venice
Mayweed
Morningglory, Ivyleaf
Morningglory, Tall
Mustard, Tansy
Mustard, Wild
Mustard (Yellowtops)
Nightshade, Black
Pennycress, Field (Fanweed, Frenchweed, Stinkweed)
Pepperweed, Virginia (Peppergrass)
Pigweed, Prostrate
Pigweed, Redroot (Carelessweed)
Pigweed, Rough
Pigweed, Smooth
Pigweed (triazine resistant)
Pigweed, Tumble
Poorjoe
Puncturevine
Purslane, Common
Pusley, Florida
Radish, Wild
Ragweed, Common
Ragweed, Giant (Buffaloweed)
Ragweed Lance-Leaf
Rubberweed, Bitter (Bitterweed)
Sesbania, Hemp
Shepherdspurse
Sicklepod
Sida, Prickly (Teaweed)
Smartweed, Green
Smartweed, Pennsylvania
Sneezeweed, Bitter
Sowthistle, Annual
Sowthistle, Spiny
Spikeweed, Common
Spurge, Prostrate
Spurry, Corn
Starbur, Bristly
Sumpweed, Rough
Sunflower, Common (Wild)

12/56

Sunflower, volunteer
Thistle, Russian
Velvetleaf
Waterhemp
Waterprimrose, Winged
Wormwood, Annual

BIENNIALS

Burdock, Common
Carrot, Wild (Queen Anne's Lace)
Cockle, White
Evening primrose, Common
Geranium, Carolina
Gromwell
Knapweed, Diffuse
Knapweed, Spotted
Mallow, Dwarf
Plantain, Bracted
Ragwort, Tansy
Starthistle, Yellow
Sweetclover
Teasel
Thistle, Bull
Thistle, Milk
Thistle, Musk
Thistle, Plumeless

PERENNIALS

*Alfalfa
Artichoke, Jerusalem
Aster, Spiny
Waster, Whiteheath
Beadstraw, Smooth
Bindweed, Field
Bindweed, Hedge
Blueweed, Texas
*Bursage, (Bur Ragweed)(Lakeweed)(Povertyweed)
Bursage, Woollyleaf (Lakeweed)
Buttercup, Tall
Campion, Bladder
Chickweed, Field
Chickweed (Mouseear, Canada)

Chicory
*Clover, Hop
*Dandelion, Common
*Dock, Broadleaf (Bitterdock)
*Dock, Curly
Dogbane, Hemp
*Dogfennel (Cypressweed)
Fern, Bracken
Garlic, Wild
Goldenrod, Canada
Goldenrod, Missouri
Goldenweed, Common
Hawkweed
Horsenettle, Carolina
Ironweed
Knapweed, Black
Knapweed, Russian
Mare's Tail (Horseweed)
Milkweed, Climbing
Milkweed, Common
Milkweed, Honeyvine
Milkweed, Western Whorled
Nettle, Stinging
Nightshade, Silverleaf (White Horsenettle)
Onion, Wild
*Plantain, Broadleaf
Plantain, Buckhorn
Pokeweed
Ragweed, Western
Redvine
Sericia Lespedeza
Smartweed, Swamp
Snakeweed, Broom
*Sorrel, Red (Sheep Sorrel)
Sowthistle
Sowthistle, Perennial
Spurge, Leafy
Sundrop, Halfshrub (Eveningprimrose)
Thistle, Canada
Toadflex, Dalmation
Tropical Soda Apple
Trumpet creeper (Buckvine)
Vetch
Waterhemlock
Waterprimrose, creeping

*Woodsorrel, Creeping
 Common Yellow
 Wormwood, Common
 Wormwood, Louisiana
 *Yankeeweed
 Yarrow, Common

* Noted perennials may be controlled using ORACLE at rates lower than those recommended for other listed perennial weeds. (See APPLICATION RATES AND TIMINGS sections on pages 40_ and 41).

WOODY

Alder
 Ash
 Aspen
 Basswood
 Beech
 Birch
 *Blackberry
 *Blackgum
 *Cedar
 Cherry
 Chinquapin
 Cottonwood
 *Creosotebush
 Cucumbertree
 *Dewberry
 *Dogwood
 Elm
 Grape
 *Hawthorn (Thornapple)
 Hemlock
 Hickory
 Honeylocust
 Honeysuckle
 Hornbeam
 Huckleberry
 Huisache
 Ivy, Poison
 Kudzu
 Locust, Black
 Maple
 Mesquite
 Oak

Oak, Poison
 Olive, Russian
 Persimmon, Eastern
 Pine
 *Plum, Sand (Wild Plum)
 Poplar
 Rabbitbrush
 *Redcedar, Eastern
 *Rose, McCartney
 *Rose, Multiflora
 Sagebrush, Fringe
 Sassafras
 Serviceberry
 Spicebush
 Spruce
 Sumac
 *Sweetgum
 Sycamore
 Tarbush
 Willow
 Witchhazel
 *Yaupon
 *Yucca

* Growth Suppression

FIELD, SEED*, POPCORN* AND SILAGE CORN

Observe all PRECAUTIONS, MIXING and APPLICATION instructions on pages 5 as well as the following:

* Do not apply ORACLE to seed corn or popcorn without first verifying with your local seed corn company (supplier) the ORACLE selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.

ORACLE is not registered for use on sweet corn.

Direct contact of ORACLE with corn seed must be avoided. If corn seeds are less than 1 1/2 inches below the surface, delay application until corn has emerged.

Up to 2 applications of ORACLE may be made during a growing season. Do not exceed a total 1 1/2 pints of ORACLE per treated acre per crop year. Allow two weeks or more between applications of ORACLE. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.

16/50

Applications of ORACLE to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.

Agriculturally approved surfactants or sprayable fertilizers (1/2-1 gallon per acre of 28%, 30%, 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate*) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum-based oils after crop emergence or crop injury may result.

Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

Several synthetic pyrethroid insecticides are labeled for tankmix applications with ORACLE. Refer to their label for specific recommendations.

* Not for use in California

WEEDS CONTROLLED

ORACLE will control many ANNUAL broadleaf weeds or give growth suppression of many PERENNIAL broadleaf weeds commonly found in corn. (Refer to the GENERAL WEED LIST on pages 10 - 14).

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

PREPLANT/PREEMERGENCE IN NO-TILLAGE CORN

Applications of ORACLE may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply ORACLE at 1 pint per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 1/2 pint per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply ORACLE after 4-6 inches of regrowth has occurred.

PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN

ORACLE may be applied after planting and prior to corn emergence. Application at 1 pint per treated acre may be made to medium or fine textured soils, which contain 2% or greater organic matter. DO NOT apply to coarse textured soils (sand, sandy loam and loamy sand) until after crop emergence (see Early Postemergence uses below).

Preemergence application of ORACLE does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrate treated soil over seed furrow.

EARLY POSTEMERGENCE (ALL TILLAGE SYSTEMS)

Spike through 8 inch tall corn

ORACLE at 1 pint per treated acre may be applied during the period from corn emergence through the five leaf stage or 8 inches tall, which ever comes first. Reduce the rate to 1/2 pint per treated acre if corn is growing on coarse textured soils (sand, sandy loam, loamy sand). See Late Postemergence applications given below if the 6th true leaf is emerging from whorl or corn is greater than 8 inches tall.

LATE POSTEMERGENCE (ALL TILLAGE SYSTEMS)

8 to 36 inch tall corn

Application of ORACLE at 1/2 pint per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when: (1) corn leaves prevent proper spray coverage; (2) sensitive crops are growing nearby; (3) tank mixing with 2,4-D.

DO NOT apply ORACLE when soybeans are growing nearby if any of these conditions exist:

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

OVERLAY (SEQUENTIAL) TREATMENTS

ORACLE may be applied to ground previously treated with one or more of the following herbicides:

acetochlor (Surpass®, Harness® Plus)

alachlor (Lasso®, Lasso MT®, Partner®)
atrazine
Bicep®
Broadstrike® + Dual®
Broadstrike® plus
Bronco®
Bullet®
butylate (Sutan+®/Genate®)
Clarity®
cyanazine (Bladox®)
dimethenamid (Frontier®)
EPTC (Eradicane®)
Extrazine II®
Guardsman®
glyphosate (Roundup®)
halosulfuron (Battalion®, Permit®)
Lariat®
Marksman®
metolachlor (Dual®)
paraquat (Gramoxone®)
pendimethalin (Prowl®)
propachlor (Ramrod®)
simazine (Princep®)
Surpass® 100

Apply ORACLE at 1/2 pint per treated acre to ground previously treated with full rates of Clarity® or Marksman® herbicides. Allow at least 2 weeks between applications.

READ AND FOLLOW LABEL DIRECTIONS FOR EACH OF THE ABOVE PRODUCTS.

TANK MIX TREATMENTS FOR CORN

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. **Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, and other restrictions.**

RATES AND TIMINGS

ORACLE PLUS	PREPLANT/PRE EMERGENT (NO TILLAGE CORN) DIRECTIONS	PREEMERGENT (CONVENTIONAL OR REDUCED TILLAGE CORN)	EARLY POST-EMERGENT (ALL TILLAGE SYSTEMS)	LATE POSTEMERGENT (ALL TILLAGE SYSTEMS)	ADDITIONAL DIRECTIONS
ACCENT® Nicosulfuron	-	-	1/2-1 ounce ai/A	1/2-1 ounce ai/A (to improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall.)	Application may be made to emerged weeds before corn is greater than 24 inches tall. Use non-ionic surfactant at .25% (V/V) with this tank mixture.
Atrazine	1 1/4-2 lbs. ai/A	1 1/4-2 lbs. ai/A	1 1/4-2 lbs. ai/A Crop oil concentrates may be used with this mixture if corn is 5 inches or less in height.	1 1/4-2 lbs. ai/A Do not apply if corn is greater than 12 inches tall.	Application may be made before grasses are 1 1/2" tall. Follow all state and Federal restrictions pertaining to atrazine applications.
BEACON® primisulfuron	-	-	0.31-0.62 ounce ai/A	0.31-0.62 ounce ai/A (to improve spray coverage of weeds and reduce risk of corn injury.	Application may be made to emerged weeds when corn is 4 to 24 inches tall. Use non-ionic surfactant at .25% (V/V) with this

				use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall).	tank mixture.
BLADEX® cyanazine	1 1/4-4 lbs. ai/A	1 1/4-4 lbs. ai/A	1 1/4-2 lbs. ai/A (use the 90 DF formulation only, after corn emergence.)	-	Application may be made before grasses are 1 1/2 inches tall, and before corn is beyond the 4 leaf stage.
DUAL® metolachlor	1 1/2-3 lbs. ai/A	1 1/2-3 lbs. ai/A (use only on fine or medium soils with 2 1/2% or greater organic matter.)	1 1/2-3 lbs. ai/A	-	Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall.

FRONTIER® dimethenamid	13-25 fl. oz/A	13-25 fl. oz/A (use only on fine or medium textured soils with 2.5% or greater organic matter.)	13-25 fl. oz/A	-	Application may be made up to 8 inch tall corn. This treatment must be combined with a herbicide that provides postemergence control of grass weeds if they are greater than 1 inch tall at the time of application.
GRAMOXONE® paraquat	1/4-1 lb. ai/A	1/4-1 lb. ai/A	-	-	Application may be made to emerged weeds but prior to corn emergence.
HARNESS®PLUS or SURPASS® acetochlor	1 1/2-3 lbs. ai/A	1 1/2-3 lbs. ai/A Use only on fine or medium textured soils with 2.5% or greater organic matter.	-	-	Application should be made prior to corn emergence.
LASSO® alachlor	1 1/2-4 lbs. ai/A	1 1/2-4 lbs. ai/A (use only on fine textured soils with greater than 2 1/2% organic matter).	1 1/2-4 lbs. ai/A	-	Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall. If microencapsulated forms of alachlor are used (Lasso MT, Partner), applications must be made prior to grass emergence.
PRINCEP® simazine	2.0-3.0 lbs. ai/A	2.0-3.0 lbs. ai/A	-	-	Application may be made prior to corn or weed emergence.
PROWL® pendimethalin	-	3/4-1 1/2 lbs. ai/A (use only on fine or medium textured soils with 2 1/2% or greater organic matter.)	3/4-1 1/2 lbs. ai/A	-	Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2 leaf stage of growth.
ROUNDUP®	1.0-3.0 lbs. ai/A	1.0-3.0 lbs. ai/A	-	-	Application may be made

glyphosate					to emerged weeds but prior to corn emergence.
2,4-D	1/4-1/2 lbs. ai/A	1/4-1/2 lbs. ai/A	Not recommended	1/8 lbs. ai/A	Drop pipes are to be used when corn height is 8 inches or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of the weed foliage.

SORGHUM (Milo)

Observe all PRECAUTIONS on pages __, including the reference to crops growing under stress.

Read and follow mixing and application instructions on page 5.

Applications of ORACLE to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture or hay, refer to the pasture use section of this label. Do not apply ORACLE to sorghum grown for seed production.

Make no more than one application per growing season.

WEEDS CONTROLLED

ORACLE, when applied at the recommended rate for sorghum, will control many actively growing ANNUAL broadleaf weeds and will reduce competition from established PERENNIAL broadleaf weeds as well as control their seedlings. (Refer to GENERAL WEED LIST on pages 10 - 14).

RATES AND TIMINGS

ORACLE may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of ORACLE must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3-5 leaf stage and weeds are

small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

BROADCAST RATE PER TREATED ACRE: 1/2 pint (1/4 lb. a.i.)

TANK MIX TREATMENT

ORACLE plus Atrazine

For improved control of emerged, actively growing broadleaf weeds including triazine resistant species and added suppression of perennial broadleaf weeds, tank mix 1/2 pint ORACLE with 0.5 to 1.25 lbs. a.i. atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix 1/2 pint ORACLE with 2 lbs. a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3-8 inches tall and when broadleaf weeds are small (less than 6 inches tall).

Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all State and Federal restrictions pertaining to atrazine applications.

ORACLE plus Buctril®

For improved control of broadleaf weeds, tank mix 1/2 pint ORACLE with 1-1 1/2 pint Buctril Herbicide per treated acre. Make application at 4 leaf to 15 inch tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall.

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS AND OTHER RESTRICTIONS.

OVERLAY (SEQUENTIAL) TREATMENTS

ORACLE may be applied to ground previously treated with one or more of the following herbicides:

Herbicide	Maximum rate per treated acre (lbs. a.i.)
alachlor (Lasso®) (Screen®-treated seed)	4
atrazine ¹	2.5
metolachlor (Dual®) (Concep®-treated seed)	2.5
propachlor (Ramrod®)	5

¹ Maximum use rate for atrazine is determined by soil type, tillage practices used, surface residue, and state or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for atrazine.

PREHARVEST USES

For Use Only in the States of Texas and Oklahoma

ORACLE may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial applications use at least 2 gallons of water-based carrier per treated acre.

Delay harvest until 30 days after treatment.

BROADCAST RATE PER TREATED ACRE: 1/2 pint (1/4 lbs. a.i.)

SMALL GRAINS (WHEAT, BARLEY AND OATS)

Not Underseeded to Legumes

Important

Observe all PRECAUTIONS on pages 2. Read and follow CLEANING, MIXING and APPLICATION instructions on pages 5.

If small grains are used for pasture hay, the following restrictions apply:

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
- There is no waiting period between treatment and grazing for non-lactating dairy animals.
- Treated areas may not be grazed by lactating dairy animals before 7 days after treatment.
- Do not harvest hay from treated areas before 37 days after treatment.

NOTE: Observe all precautions and restrictions on the labels of products used in tank mix treatments.

WEEDS CONTROLLED

ORACLE, or combinations with listed tank mix partners, will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds, it is recommended that ORACLE be applied in a tank mix with other herbicides. Refer to specific crop tank mix options.

Alkanet¹
 Bedstraw, Catchweed¹
 Bindweed, Field²
 Buckwheat, Tartary
 Buckwheat, Wild
 Carpetweed¹
 Chamomile, Corn
 Chervil, Bur¹
 Chickweed, Common¹
 Cockle, Corn
 Cockle, Cow
 Cocklebur, Common
 Cornflower (Bachelorbutton)¹
 Dandelion, Common²
 Dock, Curly²
 Dragonhead, American¹
 Evening Primrose, Cutleaf¹
 Falseflax, Smallseeded¹
 Fiddleneck (Tarweed)¹
 Flixweed¹
 Fumitory¹
 Gromwell, Corn¹
 Groundsel, Common¹
 Hempnettle¹
 Henbit
 Jacobs Ladder¹
 Knawel (German Moss)
 Knotweed, Prostrate
 Kochia
 Ladysthumb
 Lambsquarters, Common
 Lettuce, Miners¹
 Lettuce, Prickly
 Mallow, Common
 Mayweed, Chamomile (Dogfennel)¹
 Mustard, Blue (Purple)¹
 Mustard, Tansy
 Mustard Treacle¹
 Mustard, Tumble (Jim Hill)¹
 Mustard, Wild¹

Nightshade, Black
 Nightshade, Cutleaf¹
 Nightshade, Silverleaf² (White Horsenettle)
 Pennycress, Field (Fanweed, Frenchweed, Stinkweed)
 Pepperweed, Peppergrass¹
 Pigweed, Redroot (Carelessweed)
 Pigweed, Rough
 Pigweed, Tumble
 Pineappleweed¹
 Plantain, Broadleaf²
 Poppy, Red Horned¹
 Puncturevine¹
 Purslane, Common¹
 Radish, Wild¹
 Ragweed, Common¹
 Ragweed, Giant (Buffaloweed)¹
 Rocket, London¹
 Rocket, Yellow¹
 Salsify (Goatsbeard)¹
 Sheperdspurse¹
 Smartweed, Green
 Smartweed, Pennsylvania
 Sorrel, Red (Sheep Sorrel)¹
 Sowthistle, Annual
 Starthistle, Yellow¹
 Sunflower, Common (Wild)
 Thistle, Canada²
 Thistle, Russian
 Yarrow, Common²
 Velvetleaf
 Vetch¹

¹ These weeds will be controlled with ORACLE tank mixtures. Refer to tank mix label for specific weeds controlled.

² ORACLE tank mixes will provide suppression of established broadleaf weeds and control of their seedlings.

RATES AND TIMINGS

Application of ORACLE may be made before, during or after planting small grains. For best performance, make applications when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Application of ORACLE to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use ORACLE at 2 to 4 fluid ounces per treated acre in wheat, fall seeded barley, and oats, and at 2 to 3 fluid ounces per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia, Russian thistle and prickly lettuce or dense vegetative growth.

ORACLE used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for ORACLE rate and application timing.

For applications prior to the emergence of weeds or when sulfonyleurea resistant weeds are present or suspected, use a minimum of 3 fluid ounces per treated acre of ORACLE with a tank mix herbicide. Non-sulfonyleurea herbicides such as 2,4-D or MCPA tank mixed with ORACLE will offer more consistent control of sulfonyleurea resistant weeds.

When tank mixing with sulfonyleurea herbicides, such as Ally®, Amber®, Express®, Finesse®, Glean® and Harmony® Extra, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1-4 pints/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

FALL AND SPRING SEEDED WHEAT

ORACLE MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE. APPLICATIONS TO SPRING SEEDED WHEAT MUST BE MADE BEFORE WHEAT EXCEEDS THE 5 LEAF STAGE. Early developing wheat varieties such as TAM 107, MADISON, or WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

TANK MIX TREATMENTS

ORACLE may be tank mixed with one or more, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE: Apply 2-4 fluid ounces of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8-12 fl oz. (.25-.375 lb ai/A) ¹

MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (.25-.375 lb ai/A) ¹
Ally®	metsulfuron-methyl	60% DF	1/10 oz.
Amber®	triasulfuron	75% DF	0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/6 oz.
Finesse®	chlorsulfuron + metsulfuron-methyl	75% DF	1/3 oz.
Glean®	chlorsulfuron	75% DF	1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/3 oz.
Buctril®	bromoxynil ²	2 lb/gal	1-1.5 pts.
Bronate®	bromoxynil + MCPA	4 lb/gal	1-2 pts.
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	2-2 2/3 pts.
Stinger®	clopyralid	3 lb/gal	1/4-1/3 pt.
Karmex® ³	diuron ²	80% DF	1/2-1.5 lbs.
Sencor® ³	metribuzin ²	75% DF	1-10 oz.
Dakota® ⁴	fenoxaprop-ethyl + MCPA	3.1 lb/gal	16 oz.
Tiller® ⁴	fenoxaprop-ethyl + MCPA + 2,4-D	2.7 lb/gal	1-1.7 pts.

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

² Herbicides with the same active ingredient and/or different formulation may be used.

³ Tank mixtures for fall seeded wheat only.

⁴ Use 2 fluid ounces of ORACLE only. Do not use if wild oats is the target weed. Do not use on Durum wheat.

SPECIAL USE TANK MIXES FOR SPRING AND FALL SEEDED WHEAT (See Footnotes for Applicable Uses)

Apply 3-4¹ fluid ounces of ORACLE with:

Product ²	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D or MCPA Amine	2,4-D or MCPA	4 lb/gal	1-2 pts. ³ (0.5-1.0 lb ai/A) ⁴

2,4-D or MCPA Ester	2,4-D or MCPA	4 lb/gal	1-1.5 pts. ³ (.5-.75 lb ai/A) ⁴
Ally®	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber®	triasulfuron	75% DF	0.14-0.28 oz.
Express®	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Finesse®	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz.
Glean®	chlorsulfuron	75% DF	1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz
Ally® + 2,4-D Amine or Ester ⁵	metsulfuron-methyl + 2,4-D	60% DF + 4 lb/gal	1/20-1/10 oz + 8 fl oz.
Amber® + 2,4-D Amine or Ester ⁵	triasulfuron + 2,4-D	75% DF + 4 lb/gal	0.14-0.28 oz + 8 fl oz.
Express® + 2,4-D Amine or Ester ⁵	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/12-1/6 oz + 8 fl oz.
Finesse® + 2,4-D Amine or Ester ⁵	(chlorsulfuron + metsulfuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/6-1/3 oz + 8 fl oz.
Glean® + 2,4-D or Ester ⁵	chlorsulfuron + 2,4-D	75% DF + 4 lb/gal	1/6 oz + 8 fl oz.
Harmony® Extra + 2,4-D Amine or Ester ⁵	(thifensulfuron + tribenuron-methyl) + 2,4-D	75% DF + 4 lb/gal	1/6-1/3 oz + 8 fl oz
Roundup® RT ⁶	glyphosate	3.0 lb/gal	12-16 fl oz.

¹ ORACLE may be used at 6 fluid ounces on fall seeded wheat in Western Oregon as a spring application only. In CO, KS, NM, OK and TX up to 8 fluid ounces of ORACLE may be applied on fall seeded wheat after it exceeds the 3 leaf stage for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. ORACLE may be tank mixed with 2,4-D amine at 8 fluid ounces after wheat begins to tiller. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For Fall applications only, do not use if the potential for crop injury is not acceptable.

² Do not use low rates of sulfonyleurea herbicides, such as Ally®, Amber®, Express®, Finesse®, Glean®, and Harmony® Extra on more mature weeds and/or on dense vegetative growth.

³ NOTE: For use on Fall Seeded Wheat Only. Do Not Use unless potential crop injury will be acceptable.

⁴ When using formulations other than 4 lb/gal use pounds active/acre listed.

⁵ Use for improved control of Russian thistle, flxweed, gromwell, mayweed and fiddleneck.

⁶ ORACLE may be applied at 2 fluid ounces with Roundup® RT as a preplant application to small grains with no waiting period prior to planting. Add 0.5% by volume of an agriculturally approved non-ionic surfactant.

FALL SEEDED BARLEY

ORACLE MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE.

NOTE: For spring barley varieties that are seeded during the winter months or later, follow the rates and timings given for spring seeded barley.

TANK MIX TREATMENTS

ORACLE may be tank mixed with one or more, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2-4 fluid ounces ORACLE with:

Product ¹	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8 fl oz. (.25 lb ai/A) ²
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (.25-.375 lb ai/A)
Ally®	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber®	triasulfuron	75% DF	0.14-0.28 oz.

Express®	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Finesse®	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz.
Glean®	chlorsulfuron	75% DF	1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz.
Sencor®	metribuzin ³	75% DF	1-10 oz.
Buctric®	bromoxynil	2 lb/gal	1-1 1/2 pts.
Bronate®	bromoxynil + MCPA	4 lb/gal	3/4 - 1 1/2 pts.

¹ Do not use low rates of sulfonyleureas (Ally®, Amber®, Express®, Finesse®, Glean®, and Harmony® Extra) on more mature weeds and/or on dense vegetative growth.

² When using formulations other than 4 lb/gal use pounds active/acre listed.

³ Herbicides with the same active ingredient and/or different formulations may be used.

SPRING SEEDED BARLEY

ORACLE MUST BE APPLIED BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4 LEAF STAGE.

TANK MIX TREATMENTS

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2-3 fluid ounces ORACLE with:

Product ¹	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fl oz. (.25-.375 lb ai/A) ²
Ally®	metsulfuron-methyl	60% DF	1/20-1/10 oz.
Amber®	triasulfuron	75% DF	0.14-0.28 oz.

Express®	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz.
Finesse®	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz.
Glean®	chlorsulfuron	75% DF	1/6 oz.
Harmony® Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz.
Sencor®	metribuzin ³	75% DF	1-10 oz.
Buctrik®	bromoxynil	2 lb/gal	1-1 1/2 pts.
Bronate®	bromoxynil + MCPA	4 lb/gal	3/4-1 1/2 pts.

¹ Do not use low rates of sulfonyleureas (Ally®, Amber®, Express®, Finesse®, Glean®, and Harmony® Extra) on more mature weeds and/or on dense vegetative growth.

² When using formulations other than 4 lb/gal use pounds active/acre listed.

³ Herbicides with the same active ingredient and/or different formulations may be used.

FALL AND SPRING SEEDED OATS

ORACLE MUST BE APPLIED BEFORE SPRING SEEDED OATS EXCEED THE 5 LEAF STAGE. APPLICATIONS TO FALL SEEDED OATS MUST BE MADE PRIOR TO THE JOINTING STAGE.

TANK MIX TREATMENTS

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 2-4 fluid ounces ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fluid oz. (.25-.375 lb ai/A) ¹

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

SUGARCANE

Observe all PRECAUTIONS on pages 2. Read and follow MIXING AND APPLICATION instructions on page 5.

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

WEEDS CONTROLLED

ORACLE, when applied at recommended rates, will control many ANNUAL, BIENNIAL and PERENNIAL broadleaf weeds commonly found in sugarcane. (Refer to GENERAL WEED LIST on pages 10 – 14.)

RATES AND TIMINGS

Application of ORACLE may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane. Application rates and timings of ORACLE are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage and Type	Amount Product	Broadcast Rate Per Treated Acre lbs. a.i.
Annual		
Small, actively growing	1/2-1 pt	1/4 - 1/2
Established weed growth	1-1 1/2 pts	1/2 - 3/4
Biennial	1-2 pts	1/2 - 1
Perennial	2-4 pts	1 - 2*

* Application made over the top of actively growing sugarcane may result in crop injury.

When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

Retreatments may be made as needed, however, do not exceed a total of 4 pints (2 lbs. a.i.) of ORACLE per treated acre during a growing season.

TANK MIX TREATMENTS

ORACLE may be tank mixed with one or more of, but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled, geographic and other restrictions.

Herbicide	Rates per treated acre (lbs. a.i.)
Ametryn (Evik®)	2/5 - 8
asulam (Asulox®)	2 - 3 1/3
atrazine	2/5 - 4
2,4-D	1/2 - 3*

* Application of ORACLE plus 2,4-D tank mix at the higher listed rate range may result in crop injury.

PASTURE, HAY, RANGELAND AND GENERAL FARMSTEAD (Non-Cropland), RIGHTS-OF-WAY, AND PUBLIC UTILITY AND INDUSTRIAL AREAS

ORACLE is recommended for use on pasture, hay, rangeland, general farmstead (non-cropland) (including fence rows and non-irrigation ditchbanks) for broadleaf weed and brush control, and for use on non-cropland areas such as rights-of-way (such as roadways, rest areas, utility, railroad, highway, pipeline, and rights-of-way that run through pasture and rangeland); public utility facilities (such as substations, pipelines, tankfarms, pumping stations, parking and storage areas, fencerows and nonirrigated ditchbanks); brush control for forest site preparation or maintenance. ORACLE may also be applied to non-cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatment of roadsides and highways, utilities, railroad and pipeline rights-of-way. Noxious weeds

must be recognized at the State level but programs may be administered at State, County or other levels.

Observe all PRECAUTIONS on pages 2. Read and follow MIXING AND APPLICATION instructions on page 5.

GENERAL FARMSTEAD

ORACLE uses described in this section also pertain to small grains (such as barley, forage, sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

NEWLY SEEDED AREAS, including small grains grown for pasture may be severely injured if rates of ORACLE greater than 1 pint/acre are applied.

ESTABLISHED GRASS CROPS growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied. Furthermore, rates of ORACLE in excess of 2 quarts (2 lbs. a.i.) per treated acre may cause temporary injury to many grass species.

Bentgrass, carpetgrass, buffalograss and St. Augustine grass may be injured at rates exceeding 1 pint ORACLE (1/2 lb. a.i.) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured.

Treatments will kill alfalfa, clovers, lespedeza, wild winter peas, vetch and other legumes.

ANIMALS CANNOT BE REMOVED FROM TREATED AREA FOR SLAUGHTER PRIOR TO 30 DAYS AFTER LAST APPLICATION.

THERE IS NO WAITING PERIOD BETWEEN TREATMENT AND GRAZING FOR NON-LACTATING ANIMALS.

TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT

ORACLE Rate per Treated Acre	Days Before Grazing	Days Before Hay Harvest
Up to 1 pint (1/2 lb. a.i.)	7 days	37 days
Up to 1 quart (1 lb. a.i.)	21 days	51 days
Up to 2 quarts (2 lbs. a.i.)	40 days	70 days

NOTE: Observe all precautions and restrictions on labels of products used in tank mixtures.

RIGHTS-OF-WAY

Oracle can be used to control many broadleaf weeds on rights-of-way. This use includes applications to roadside, roadway and highways; to areas along utilities such as cable and powerlines; railroad track and embankment; highways, highway medians, bridge abutments, pipelines, and rights-of-way that run through pasture and rangeland. Use controlled application techniques that minimize the risk of off-target movement.

PUBLIC UTILITY AND INDUSTRIAL AREAS

Oracle can be used to control many broadleaf weeds and brush in noncrop areas on or surrounding substations, pipelines, tank farms, pump stations, production facilities, and bareground situations. It may also be used on parking and storage areas (refer to Best Stewardship Practices to avoid direct runoff from impervious surfaces).

MIXING AND APPLICATION

Read and observe Sensitive Crop Precautions recommendations in this label.

ORACLE can be applied using water, oil in water emulsions (including invert systems), or sprayable fluid fertilizer as a carrier. A COMPATABILITY TEST (page of this booklet) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

ORACLE may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment apply 1 to 40 gallons of diluted spray per treated acre in a water-based carrier.

ORACLE may be applied to individual clumps or small areas (SPOT TREATMENT) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to run-off) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

WEEDS AND BRUSH CONTROLLED

ORACLE, when applied at recommended rates, will give control of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species commonly found in pasture, hay, rangeland, and general farmstead (non-cropland) areas. (Refer to GENERAL WEED LIST on pages 10-14.)

Noted (*) PERENNIAL weeds may be controlled with lower rates of either ORACLE or ORACLE plus 2,4-D. See RATES AND TIMINGS below.

RATES AND TIMINGS

Application rates and timing of ORACLE are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage & Type	Amount Product	Broadcast Rate per Treated Acre (lbs. a.i.)
Annual Small, actively growing Established weed growth	1/2 - 1 pt. 1 - 1 1/2 pts.	1/4 - 1/2 1/2 - 3/4
Biennial ¹ Rosette diameter Less than 3 inches 3 inches or more Bolting	1/2 - 1 pt. 1 - 2 pts. 2 - 3 pts.	1/4 - 1/2 1/2 - 1 1 - 1 1/2
Perennial Suppression or top growth control Noted (*) Perennials Other perennials	1/2 - 1 qt. 1 - 2 qts. 2 qts.	1/2 - 1 1 - 2 2
Woody Brush & Vines Top growth suppression Top growth control ² Stems and stem suppression	1/2 - 1 qt. 1 - 2 qts. 2 qts.	1/2 - 1 1 - 2 2

¹ For best performance, make application when BIENNIAL weeds are in the rosette stage.

² Species noted in GENERAL WEED LIST pages 10-14 will require tank mixtures for adequate control.

Retreatments may be made as needed; however, do not exceed a total of 2 quarts (2 lbs. a.i.) of ORACLE per treated acre during a growing season.

Due to the variations that may occur on formulated products and specific use ingredients (e.g. water supplies), a COMPATIBILITY TEST as described on pages is recommended prior to actual tank mixing.

CUT SURFACE TREE TREATMENTS

ORACLE may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. A mix of 1 part ORACLE with 1 to 3 parts water should be used in application. ~~Use the lower dilution when treating difficult-to-control species.~~

FRILL OR GIRDLE TREATMENTS: Make a continuous cut or a series of overlapping cuts using an ax to girdle tree trunk. Spray or paint cut surface with the ORACLE/water mix.

STUMP TREATMENTS: Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

NOTE: For more rapid foliar effects, 2,4-D may be added to the ORACLE/water mix.

DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE

ORACLE can be applied when plants are dormant as an undiluted SPOT-CONCENTRATE directly to the soil or as a LO-OIL BASAL BARK treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of ORACLE should be applied directly to the soil as close as possible to the root crown but within 6-8 inches of the crown. On sloping terrain, application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying ORACLE directly to the soil. The use rate of ORACLE is dependent on the canopy diameter of the multiflora rose. Examples: Use ORACLE at 1/4, 1 or 2 1/4 fluid ounces of product respectively, for 5, 10, or 15 feet canopy diameters. Do not exceed a total of 2 qts. ORACLE per acre per year.

LO-OIL BASAL BARK applications of ORACLE should be applied to the basal stem region from the ground up to a height of 12-18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying ORACLE to the ground line. Refer to Mixing and Applications above in this section for method of preparing oil-in-water emulsion. Example for making approximately 2 gallons of a Lo-Oil spray solution mixture: combine 1 1/2 gallons water plus 1 ounce emulsifier plus 1 pint ORACLE plus 2 1/2 pints of No. 2 diesel fuel. Adjust amounts of

materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

FOREST SITE PREPARATION

GENERAL INFORMATION

Oracle may be used for control of undesirable conifers as well as many broadleaf weeds, vines, brambles, hardwood brush, and trees in forest site preparation. Oracle may be applied as broadcast foliar sprays from ground or aerial equipment. Oracle is absorbed through the leaf surfaces quickly after spraying and will also be absorbed from the soil by the roots. Translocation through the leaves, stems, and roots provides control of undesirable young conifer and broadleaf species. Woody plants, brush, and trees may not display the full extent of herbicide efficacy until several months following treatment. Oracle provides application flexibility for extended windows of application and tank mix options (refer to **Mixing and Application Instructions and Tank Mix Options**).

MIXING AND APPLICATION INSTRUCTIONS

Ground Operated Spray Equipment

Thoroughly mix and apply the recommended amount of Oracle (2 qts./A maximum) in a minimum of 15 gals. of water per acre. Spray solution should uniformly cover undesirable foliage for best results. A suitable nonionic surfactant should be added to the spray solution to enhance foliage wetting, spreading, and solution absorption. Drift control and foam reducing agents may be added at recommended rates, if needed. Spray pattern indicator agents may also be added at recommended rates, if desired. **DO NOT** spray under windy or gusty conditions. Maintain proper buffer zones to ensure drift does not reach off-target vegetation.

Aerial Spray Equipment

Thoroughly mix the recommended amount of Oracle (2 qts./ maximum) in a minimum of 10 gals. of water per acre and uniformly apply with properly calibrated aerial equipment. A suitable nonionic surfactant should be added to the spray solution to enhance wetting, spreading, and solution absorption. All precautions should be taken to minimize or eliminate spray drift. Drift control and foam control agents may be added at recommended rates, if needed.

Tank Mix Options

For extended range of species control, tank mix Oracle with other forest site preparation products such as Arsenal, Garlon, Accord, etc. Observe all precautions and restrictions on the product labels. Always follow the most restrictive label in a tank mix.

CONSERVATION RESERVE PROGRAM (CRP) ACRE

ORACLE is recommended for use on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs.

Observe all PRECAUTIONS, MIXING AND APPLICATION directions on pages 5.

ORACLE treatment will cause injury or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.

Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

Do not use adjuvants containing penetrants such as petroleum based oils after grass emergence on newly seeded grasses.

NEWLY SEEDED AREAS

ORACLE may be applied either preplant or postemergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat, or other grain species grown as a cover crop. Postemergence applications may be made after seedling grasses exceed the 3 leaf stage. Rates of ORACLE greater than 1 pint per treated acre may severely injure newly seeded grasses. Preplant applications - injury to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of ORACLE per treated acre West of the Mississippi River or 20 days per pint East of the Mississippi River.

ESTABLISHED GRASS STANDS

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species: bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustine grass may be injured when treated with ORACLE at rates exceeding 1 pint per treated acre.

WEEDS CONTROLLED

ORACLE, when applied at recommended rates, will control many annual and biennial weeds and provide control and suppression of many perennial weeds. (Refer to GENERAL WEED LIST on pages 10-14.)

RATES AND TIMINGS

Application rates and timing of ORACLE treatment are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature

Weed Type* & Stage	Broadcast Rate Per Treated Acre	
	Amount of Formulated ORACLE	Equivalent lbs. a.i.
	pints	
Annuals		
Small actively growing	1/4 to 1	1/8 to 1/2
Established weed growth	1	1/2
Biennials**		
Rosette diameter		
a) less than 3 inches	1/2 to 1	1/4 to 1/2
b) 3 inches or greater	1 to 2	1/2 to 1
c) bolting biennial	2 to 3	1 to 1 1/2
Perennials**		
Suppression/Control	2 to 4	1 to 2

* For best results, treat Biennial weeds with ORACLE when they are in the rosette stage of growth. Retreatments may be made as needed; however, DO NOT EXCEED A TOTAL OF 2 QUARTS (2 lbs. a.i.) of ORACLE per treated acre during the growing season.

** Biennial and Perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

TANK MIX TREATMENTS

To control grasses and additional broadleaf weeds, ORACLE may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2,4-D, glyphosate (Roundup®), paraquat (Gramoxone®), metsulfuron (Ally®) and others.

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES, AND OTHER RESTRICTIONS.

ASPARAGUS

IMPORTANT

Observe all PRECAUTIONS on pages 2. Read and follow MIXING AND APPLICATION instructions on pages 5.

If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

Do not harvest prior to 24 hours after treatment.

Do not use in the Coachella Valley of California.

Multiple applications may be made per growing season. Do not exceed a total of 1 pint of ORACLE per treated acre per crop year.

RATES AND TIMINGS

Apply ORACLE to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

Weeds	Rate per Treated Acre
Mustard, Black Pigweed, Redroot (Carelessweed) Sowthistle, Annual *Thistle, Canada Thistle, Russian	1/2 to 1 pt. (1/4-1/2 lb. a.i.)
*Bindweed, Field Chickweed, Common Goosefoot, Nettleleaf Radish, Wild Thistle, Milk	1 pt. (1/2 lb. a.i.)

ORACLE may be applied in a tank mixture with either 2,4-D or Roundup Herbicide for improved control of noted (*) weeds. READ AND FOLLOW 2,4-D AND ROUNDUP HERBICIDE PRODUCT LABELING FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

TURF AND LAWNS**FOR USE IN GENERAL FARMSTEAD (Non-Cropland) AND SOD FARMS**

IMPORTANT

Observe all PRECAUTIONS on page 2. Read and follow MIXING AND APPLICATION instructions on pages 5.

To avoid injury to newly seeded grasses, application of ORACLE should be delayed until after the second mowing. Furthermore, application rates in excess of 1 pint (1/2 lb. a.i.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass, carpetgrass, buffalograss, and St. Augustine grass.

In areas where roots of sensitive plants extend, do not apply in excess of 1/4 pint (1/8 lb. a.i.) of ORACLE per treated acre on coarse textured (sandy-type) soils, or in excess of 1/2 pint (1/4 lb. a.i.) per treated acre on fine textured (clay-type) soils. Do not make repeat applications in these areas for 30 days and until previous applications of ORACLE have been activated in the soil by rain or irrigation.

WEEDS CONTROLLED

ORACLE, when applied at recommended rates, will give control of many ANNUAL, BIENNIAL, and noted (*) PERENNIAL broadleaf weeds commonly found in turf. ORACLE will also give growth suppression of many other listed PERENNIAL broadleaf weeds and WOODY brush and vine species. (Refer to GENERAL WEED LIST on pages 10-14.).

MIXING AND APPLICATION

Apply 30 to 200 gallons of diluted spray per treated acre (3 qts. to 4 1/4 gals. per 1,000 sq. ft.) depending on density or height of weeds treated and on the type of equipment used.

RATES AND TIMINGS

Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	ORACLE		
	Pints per treated acre	lbs. a.i. per treated acre	Teaspoons per 1,000 sq. ft.
Annual			
Small, actively growing	1/2 to 1	1/4 to 1/2	1 to 2 1/4
Established weed growth	1 to 1 1/2	1/2 to 3/4	2 1/4 to 3 1/4
Biennial Rosette diameter			
less than 3 inches	1/2 to 1	1/4 to 1/2	1 to 2 1/4
3 inches or more	1 to 2	1/2 to 1	2 1/4 to 4 1/2
Perennials and Woody			
Brush and Vines	1 to 2	1/2 to 1	2 1/4 to 4 1/2

For best performance, apply when weeds are emerged and actively growing.

Retreatments may be made as needed; however, do not exceed a total of 2 pints (1 lb. a.i.) ORACLE per treated acre during the growing season.

TANK MIX TREATMENTS

READ AND FOLLOW THE LABEL OF EACH TANK MIX PRODUCT USED FOR PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE, APPLICATION RATES AND TIMINGS, AND OTHER RESTRICTIONS.

Tank mix treatments of ORACLE may be made with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product.

Apply 1/5 to 1/2 pint (1/10-1/4 lb. a.i.) of ORACLE per treated acre with 1/2 to 1 1/2 lbs. acid equivalent of 2,4-D, MCPA, or MCPP, or with 3/8 to 1/2 lb. a.i. of bromoxynil. Use the higher level of the listed rate ranges when treating established weeds. Repeat treatments may be made as needed; however, do not exceed 2 pints (1 lb. a.i.) of ORACLE per treated acre during the growing season.

GRASS SEED CROPS
GRASSES GROWN FOR SEED SUCH AS BERMUDA GRASS, BLUEGRASS,
FESCUE AND RYEGRASS

IMPORTANT

Observe all PRECAUTIONS on pages 2. Read and follow MIXING AND APPLICATION instructions on pages 5.

Refer to PASTURE, HAY, RANGELAND, AND GENERAL FARMSTEAD (Non-Cropland) section (pages 33) for possible grazing and feeding restrictions.

Do not use on bentgrass unless possible crop injury can be tolerated.

WEEDS CONTROLLED

ORACLE will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, it is recommended that ORACLE be applied in a tank mix with other herbicides.

Alfalfa¹
 Bedstraw, Catchweed
 Bindweed, Field
 Buttercup, Corn
 Buttercup, Creeping
 Buttercup, Western Field
 Catchfly, Nightflowering
 Chamomile, Corn
 Chickweed, Common
 Chickweed, Mouseear
 Clover
 Cockle, White
 Dock, Broadleaf
 Dock, Curly
 Hemlock, Poison
 Knapweed, Russian¹
 Knawel
 Knotweed, Prostrate
 Kochia
 Ladysthumb

Lambsquarters, Common
 Lettuce, Prickly
 Mayweed (Dogfennel)
 Ragwort, Tansy
 Sorrel, Red (Sheep Sorrel)
 Sowthistle, Annual
 Starwort, Little
 Thistle, Canada¹

¹ Top growth only.

RATES AND TIMINGS

Apply 1/2 to 1 pint of ORACLE per treated acre on SEEDLING GRASS after the crop reaches the 3-5 leaf stage. Apply up to 2 pints of ORACLE on well-established Perennial grass. DO NOT APPLY AFTER THE GRASS SEED CROP BEGINS TO JOINT. For best performance, make applications when weeds are in the 2-4 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

TANK MIX TREATMENTS

For control of grasses or additional broadleaf weeds, ORACLE may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, and geographic and other restrictions.

BROADCAST RATE PER TREATED ACRE:

Apply 1/2 to 2 pints ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	1 - 4 pts. (.5 - 2.0 lb ai/A) ¹
MCPA Amine	MCPA	4 lb/gal	1 - 2 pts. (.5 - 1.0 lb ai/A) ¹
Buctril®	bromoxynil ²	2 lb/gal	1 - 2 pts.
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	1 3/4 - 4 pts.
Karmex®	diuron ²	80% DF	2 - 4 lbs.
Stinger®	clopyralid	3 lb/gal	1/4 - 1 pt.

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

² Herbicides with the same common name and/or different formulations may be.

ANNUAL GRASS CONTROL

For suppression of ANNUAL GRASS WEEDS such as:

Brome, Downy (Cheatgrass)
 Brome, Ripgut
 Fescue, Rattail
 Windgrass

Apply up to 4 pints of ORACLE per treated acre in the fall or late summer after harvest and burning of established grass seed crops. Applications should be made immediately following the first irrigation when the soil is moist and before weeds have more than 2 leaves.

PREPLANT DIRECTIONS

(POST HARVEST\FALLOW\CROP STUBBLE\SET-A-SIDE)

FOR BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS

IMPORTANT

Observe all precautions. Read and follow mixing and application instructions.

WEEDS CONTROLLED

ORACLE may be applied alone or in tank mix combinations with other herbicides registered for this use.

ORACLE can be applied either POST HARVEST in the fall, spring or summer during the FALLOW period or to CROP STUBBLE\SET-A-SIDE acres. ORACLE, when applied at the recommended rates, will control many ANNUAL broadleaf weeds, see the WEEDS CONTROLLED section under small grains. In addition, ORACLE will control or suppress the following BIENNIAL and PERENNIAL broadleaf weeds:

Alfalfa¹
 Artichoke, Jerusalem
 Bindweed, Field
 Bindweed, Hedge
 Blueweed, Texas

Bursage (Bur Ragweed, Povertyweed, Lakeweed)¹
 Dandelion, Common¹
 Dock, Curly¹
 Dogbane, Hemp
 Garlic, Wild²
 Horsenettle, Carolina
 Knapweed, Diffuse
 Knapweed, Spotted
 Nightshade, Silver
 Rodvino
 Smartweed, Swamp
 Sowthistle, Perennial¹
 Spurge, Leafy
 Thistle, Bull
 Thistle, Canada²
 Thistle, Milk
 Thistle, Musk
 Thistle, Plumeless
 Thistle, Scotch
 Trumpet creeper (Buckvine)

¹ Perennials may be controlled using ORACLE at rates lower than those recommended for other listed perennial weeds. (See RATES AND TIMINGS under this heading).

² See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control program for these weeds.

RATES AND TIMINGS

Apply ORACLE as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or cropland or crop stubble the following spring or summer. Agriculturally approved spray additives, such as surfactants or oils, may be used to enhance spray coverage and the herbicides penetration of weed foliage. See CROPPING RESTRICTIONS for recommended interval between application and planting to prevent crop injury.

For best performance, make application when ANNUAL weeds are less than 6 inches tall, when BIENNIAL weeds are in the rosette stage and to PERENNIAL weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright PERENNIAL broadleaf weeds, such as Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, such as field bindweed and hedge bindweed, are best controlled when weeds are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulblets, after the effective period for ORACLE. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of ORACLE see the RATES AND TIMINGS section under the SMALL GRAINS heading for details.

ORACLE RATES PER TREATED ACRE:

Weed Type	Amount of Product Per Treated Acre
Annual	1/2 - 1 pts. (8 to 16 fl oz)
Biennial	1 - 2 pts. (16 to 32 fl oz)
Perennial	1 - 4 pts. (16 to 64 fl oz)
Perennial suppression	1 - 2 pts. (16 to 32 fl oz)
Noted ⁽¹⁾ perennials	2 - 4 pts. (32 to 64 fl oz)
Other perennials	4 pts. (64 fl oz)

Retreatments may be made as needed; however, do not exceed a total of 4 pints of ORACLE per treated acre during any given fallow period.

TANK MIX TREATMENTS

ORACLE may be tank mixed with one or more of , but not limited to, the following herbicides for control of grasses or additional broadleaf weeds. Read and follow the label of each tank mix product used for precautionary statements, directions for use, rates and timings, weeds controlled and geographic or other restrictions.

ORACLE BROADCAST RATE PER TREATED ACRE FOR ANNUAL WEED CONTROL:

Apply 1/4 to 1 pint ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Aatrex® 4L ¹	atrazine	4 lb/gal	0.5 - 6 pts.
Aatrex® Nine-O ¹	atrazine	90% DF	0.5 - 3.3 lbs.
Amber® ²	trisulfuron	75% DF	0.28 - 0.35 oz.

Ally® ²	metsulfuron-methyl	75% DF	0.1 oz.
Bladex® ¹	cyanazine	90% DF	2.7 - 3.6 lbs.
Cyclone®	paraquat	2 lb/gal	1 - 2 pts.
Fallowmaster®	glyphosate + dicamba	1.6 lb/gal	22 - 44 fl. oz.
Finesse® ²	chlorsulfuron + metsulfuron-methyl	75% DF	0.2 oz.
Gramoxone® Extra	paraquat	2.5 lb/gal	1.5 pts.
Kerb® ¹	pronamide	50-W	0.5 - 1.0 lb.
Landmaster® BW	glyphosate + 2,4-D	2.4 lb/gal	27 - 54 fl oz.
Roundup® or Roundup® RT	glyphosate	3 lb/gal	8 - 48 fl oz.
Sencor® DF ¹	metribuzin	75% DF	0.5 - 1 lb.
Sencor® 4 ¹	metribuzin	4 lb/gal	0.75 - 1 1/2 pts.
2,4-D	2,4-D	4 lb/gal	1 - 2 pts. (0.5 - 1 lb ai/A) ³

¹ Tank mixes of ORACLE with these products may be subject to special restrictions. See the Product Label of the tank mix partner for intended use rates, restrictions and other precautions.

² When tank mixing with sulfonylurea herbicides refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1 - 2 quarts/100 gallons of spray or not more than 0.25 - 0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonylurea resistant weeds may not be controlled by tank mixes of ORACLE and a sulfonylurea. Refer to the ORACLE tank mix section for alternative tank mixes.

³ When using formulations other than 4 lb/gal, use pounds active/acre listed.

ORACLE BROADCAST RATE PER TREATED ACRE FOR BIENNIAL AND PERENNIAL WEED CONTROL:

Apply 1 to 4 pints of ORACLE with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	2 - 4 pts.
2,4-D	2,4-D	4 lb/gal	2 - 6 pts. (1.0 - 3 lb ai/A) ¹
Landmaster®BW	glyphosate + 2,4-D	2.4 lb/gal	54 fl oz.
Roundup®	glyphosate	3.0 lb/gal	1 - 5 qts.
Roundup® RT	glyphosate	3.0 lb/gal	1 - 5 qts.
Tordon® 22K	picloram	2 lb/gal	1/2 - 1 pt.

¹ When using formulations other than 4 lb/gal use pounds active/acre listed.

SPECIAL TANK MIX TREATMENTS

For suppression of perennial weeds, apply 1/2 - 1 pints of ORACLE with 8 to 16 fluid ounces of Roundup Herbicide or Roundup RT per treated acre.

For wild garlic control, apply 1 pint ORACLE with 3 pints of 2,4-D LV Ester (4 lb/gal) per treated acre. Apply when wild garlic is 4 to 8 inches tall.

For Canada thistle control, use ORACLE, ORACLE plus Curtail®, or ORACLE plus Roundup® Herbicide or Roundup® RT tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brome, jointed goatgrass, common rye and volunteer wheat when they are actively growing. Use 1 pint ORACLE with 1/2 to 1 lb Kerb® 50W. Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply ORACLE plus Landmaster® BW or Fallowmaster® Herbicide to give improved control of kochia, wild buckwheat, prickly lettuce, field bindweed and Canada thistle. Use 1/8 - 1/4 pint ORACLE plus 22 - 54 fluid ounces of Landmaster® BW or Fallowmaster® Herbicide for annual weed control or 1/4 - 1/2 pint ORACLE plus 22 - 54 fluid ounces of Landmaster® BW or Fallowmaster® Herbicide for perennial weed suppression.

CROPPING RESTRICTIONS

The following recommendations are based on ORACLE use rates up to 4 pints per treated acre.

CORN, SORGHUM and SOYBEANS may be planted in the spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, treated areas should be cultivated to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of ORACLE per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of ORACLE per treated acre. Exclude days when ground is frozen.

WHEAT may be planted in the fall or spring following applications. Also, spot applications may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

East of the Mississippi River, the interval is 20 days per pint of ORACLE per treated acre or 1.25 days per 1 ounce. Moisture is essential for ORACLE degradation. Exclude days when ground is frozen.

West of the Mississippi River, the interval is 45 days per pint of ORACLE per treated acre or 3 days per ounce. Moisture is essential for ORACLE degradation. Exclude days when ground is frozen.

Following a normal harvest of barley, oats, or wheat, any rotational crop may be planted. If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow up with the planting of a sensitive crop.

**CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND
(SPOT APPLICATION ONLY)
FOR USE ONLY IN THE STATES OF IDAHO, MONTANA, NEVADA, OREGON,
UTAH, AND WASHINGTON**

IMPORTANT

Observe all PRECAUTIONS on pages 2. Read and follow MIXING AND APPLICATION instructions on pages 5 .

Do not treat subirrigated cropland or areas where the soil remains saturated with water throughout the year.

Make only one application of ORACLE per year.

WEEDS CONTROLLED

ORACLE, when applied at recommended rates, will control many broadleaf weeds including:

Bindweed, Field
Dock, Broadleaf (Bitterdock)
Dock, Curly
Knapweed, Black
Knapweed, Russian
Ragwort, Tansy
Spurge, Leafy
Thistle, Canada

RATES AND TIMINGS

ORACLE may be applied at any time following a crop harvest to stubble fallow or other cropland. Application should be made when weeds are actively growing and prior to a killing frost.

Apply 2 qts. (2 lbs. a.i.) of ORACLE per treated acre. Application may be made up to one month prior to the planting of wheat.

NOTE: Do not use unless injury to wheat or rotated barley will be acceptable.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application. Do not plant broadleaf crops such as alfalfa, beans, peas, potatoes, or sugarbeets into treated areas until two years after application.

In most cases, treatments will not kill perennial weed seedlings which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

WIPER APPLICATION USES

IMPORTANT

Observe all PRECAUTIONS on pages 2.

ORACLE may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution containing 1 part ORACLE to 1 part water. Do not contact desirable vegetation with herbicide solution. Wiper application should only be made to crops (including pastures) and non-cropland areas described in this label with the exception of Grain Sorghum (Milo).

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage or disposal.

STORAGE

Store in original container in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

PESTICIDE DISPOSAL

Triple rinse pesticide from containers and use rinsates in the pesticide application. Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Plastic or Metal: After triple rinsing (or equivalent) offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. ~~If burned, stay out of smoke.~~

[BULK STORAGE AND DISPOSAL: to be printed on labeling for bulk use only. AGITATE BEFORE USE]

STORAGE AND DISPOSAL**PROHIBITIONS**

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. This product may not be mixed, loaded, or used with 50 feet of all wells including abandoned wells, drainage wells, and sinkholes.

STORAGE

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

PESTICIDE DISPOSAL

Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal and local procedures under Subtitle C or the Resource Conservation and Recovery Act.

BULK TANK MAINTENANCE

Follow clean-out directions in Dealer Bulk Handling Guide for LEGEND 4L listed under Bulk Storage Tank Requirements.

GENERAL

Consult Federal, State or local disposal authorities for approved alternative procedures, such as limited burning.

NOTICE OF WARRANTY AND DISCLAIMER

Seller warrants that at the time of delivery the product in this container conforms to its chemical description contained hereon and is reasonably fit for its intended purpose under normal conditions of use. This is the only warranty made on this product. Seller expressly disclaims any implied warranties of merchantability or fitness for any particular purpose and, except as set forth above, any other express or implied warranties. Any damages arising from breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid for this product by Buyer, and shall not include incidental or consequential damages such as, but not limited to, loss of profits or values. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of the Seller. In no case shall Seller be liable for the consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. Buyer acknowledges the use of its own independent skill and expertise in the selection and use of the product and does not rely on any oral or written statements or representations.

Registered Trademarks

Aatrex, Amber, Beacon, Bicep, Concep, Dual, Evik, and Princep are registered trademarks of Novartis.

Accent, Ally, Bladex, Express, Extrazine, Finesse, Glean, Harmony, and Karmex are registered trademarks of E.I. duPont de Nemours & Co., Inc.

Asulox is a registered trademark of Rhone-Poulenc Agriculture Ltd.

Bronate is a registered trademark of Rhone-Poulenc Agrochimie.

Buctril is a registered trademark of Rhone-Poulenc Ag Company.

Battalion, Bronco, Bullet, Harness, Landmaster, Lariat, Lasso, Partner, Permit, Ramrod, Roundup, and Screen are registered trademarks of Monsanto Company.

Broadstrike, Curtail, Stinger, and Tordon are registered trademarks of DowElanco.

Cyclone, Eradicane, Surpass, and Sutan are registered trademarks of Zeneca, Inc.

Garlon is a registered trademark of Dow Chemical Company.

Genate is a trademark of Chevron Chemical Company.

Gramoxone is a registered trademark of Zeneca Ltd.

Dakota is a registered trademark of AgrEvo USA Company.

Kerb is a registered trademark of Rohm and Haas Company.

Prowl is a registered trademark of American Cyanamid Company.

Sencor is a registered trademark of Bayer AG.

Tiller is a registered trademark of Hoechst Schering Agrevo GmbH.

Tough is a registered trademark of Novartis.

Banvel, Clarity, Fallowmaster, Frontier, Guardsman, and Marksman are registered trademarks of BASF Corporation.

Guardsman and Marksman are Restricted Use Pesticides.