

71085-20

05/11/2011

1/19



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

MAY 11 2011

Ms. Judy Smith
RiceCo LLC
5100 Poplar Avenue, Suite 2428
Memphis, TN 38137

Subject: Notification per PR Notice 98-10 – Add Alternate Brand Names
Propanil EC
EPA Reg. No. 71085-20
Application Dated – May 11, 2011

Alternate Brand Names:	Propanil 4 Herbicide	Herbax 48 EC
	Propanil 480	Propasint 4E
	Propanil 48 EC	Super Propanil 500
	Supernox 500	Propasint 48 EC
	Propanil 4	Propanil 48
	Propanex 500	

Dear Ms. Smith:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10 for the subject product.

The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action requested falls within the scope of PRN 98-10. The label submitted with the application has been date-stamped "Notification" and will be placed in our records.

If you have any questions regarding this letter, please contact Maggie Rudick at (703) 347-0257 or rudick.maggie@epa.gov.

Sincerely,

Kable Bo Davis, Product Manager 25
Herbicide Branch
Registration Division (7505P)

	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input checked="" type="checkbox"/> Amendment <input type="checkbox"/> Other	OPP Identifier Number
	Application for Pesticide - Section I		

1. Company/Product Number 71085-20	2. EPA Product Manager Kable Bo Davis	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) PROPANIL EC	PM# 25	
5. Name and Address of Applicant (Include ZIP Code) RiceCo LLC 5100 Poplar Avenue, Suite 2428 Memphis, TN 38137 <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

MAY 11 2011

Notification adding the following alternate names: Propanil 4 Herbicide, Propanil 480, Propanil 48 EC, Supemox 500, Propanil 4, Propanex 500, Herbax 48 EC, Propasint 4E, Super Propanil 500, Propasint 48 EC, Propanil 48. These alternate names to be transferred to 71085-20. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No		<input checked="" type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container		Glass	
		If "Yes" Package wgt. No. per container		Paper	
3. Location of Net Contents Information <input checked="" type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container 2.5, 5, 30, 55, bulk gallons		5. Location of Label Directions <input type="checkbox"/> on label	
6. Manner in Which Label is Affixed to Product <input checked="" type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled				<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Judy Smith	Title Regulatory Consultant	Telephone No. (Include Area Code) 901-684-5390
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 		
3. Title Regulatory Consultant		
4. Typed Name Judy Smith		5. Date May 11, 2011



8/19

May 11, 2011

Mr. Kable Davis PM 25
Registration Division (7505P)
U. S. Environmental Protection Agency
One Potomac Yard
2777 Crystal Drive
Arlington, VA 22202

**RE: ALTERNATE NAMES
PROPANIL EC – EPA REG. 71085-20**

Dear Mr. Davis:

Enclosed please find the following:

- 1 – 8570-1 Notification form
- 1 – Copy of Label

RiceCo LLC would like to add the below listed alternate names to the above referenced product:

- Propanil 4 Herbicide
- Propanil 480
- Propanil 48 EC
- Supernox 500
- Propanil 4
- Propanex 500
- Herbax 48 EC
- Propasint 4E
- Super Propanil 500
- Propasint 48 EC
- Propanil 48

RiceCo LLC recently cancelled the above list of alternate names from 71085-2 as that product's formulation is solvent free and Propanil EC is not solvent free. All of the alternate names listed are used in RiceCo's international business.

RiceCo would like to point out to the EPA that although the active ingredient label claim is 43.5% for Propanil EC, the "48 or 480" listed in the above names represents that the product contains 4 lbs. active ingredient per gallon. I was told this is common knowledge in foreign countries and as stated above the alternate names listed are used for our international business.

If additional information is required, please do not hesitate to contact me by telephone @ (901) 684-5390 or by email @ riceco.reg@ricecollc.com.

Sincerely,

RICECO LLC

Judy Smith
RiceCo Regulatory Department

"...from the paddy to the plate"

4/19

NOTIFICATION
MAY 11 2011



PROPANIL EC

Herbicide

ALTERNATE NAMES: Propanil 4 Herbicide, Propanil 480, Propanil 48 EC, Supernox 500, Propanil 4, Propanex 500, Herbax 48 EC, Propasint 4E, Super Propanil 500, Propasint 48 EC, Propanil 48

Active Ingredient:
Propanil: (3', 4'-dichloropropionanilide) 43.50%
Other Ingredients: 56.50%
TOTAL 100.00%
Equivalent to 4 lbs. active ingredient per gallon.

EPA Registration No. 71085-20

EPA Est. No. 34704-MS-01; 62171-MS-3

KEEP OUT OF REACH OF CHILDREN

WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se le explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
If Swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to by a poison control center or doctor. • Do not give anything by mouth to an unconscious person
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	

In Case of Chemical Spill, Leak, Exposure Call
Global Logistics @
(504) 439-3140 or (727) 374-5705

MANUFACTURED FOR:
RiceCo LLC
Memphis, TN 38137

NET CONTENTS: 30 GAL

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING

Causes skin irritation. Causes moderate eye irritation. Harmful if swallowed. Do not get in eyes, on skin, or clothing. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of barrier laminate or butyl rubber. If you want more options, follow the instructions for category B on an EPA chemical resistance category selection chart.

(This section is for formulations NOT packaged with Built-in Probes)

Mixers, loaders, ground applicators, and handlers cleaning up spills or equipment or otherwise exposed to the concentrate and handlers removing an unrinsed probe must wear the following:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves,
- Chemical-resistant footwear plus socks,
- Protective eyewear if the system operates under pressure, and,
- Chemical-resistant apron when mixing and loading.

Pilots and handlers removing a triple-rinsed probe must wear:

- Long-sleeved shirt,
- Long pants, and
- Shoes plus socks

See Engineering Controls for additional requirements.

(This section is for formulations packaged WITH built-in probes)

Some materials that are chemical-resistant to this product are made of barrier laminate or butyl rubber. If you want more options, follow the instructions for category B on an EPA chemical-resistance category selection chart.

Mixers, loaders, applicators and other handlers must wear the following:

- Long-sleeved shirt,
- Long pants
- Shoes plus socks, and
- Chemical-resistant gloves and chemical-resistant apron when mixing/loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

See Engineering Controls for additional requirements.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Note to EPA Reviewer: The following Engineering Controls will be used if product packaged with built in probe

ENGINEERING CONTROLS

Mixers and loaders must use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4)] for dermal protection and must:

- Wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders,
- Wear protective eyewear, if the system operates under pressure, and
- Chemical-resistant footwear and coveralls must be provided and be immediately available for use in an emergency, such as a broken package, spill, or equipment breakdown.

Note to EPA Review: The following Engineering Controls will be used if product packaged without built in probe.

ENGINEERING CONTROLS

Mixers and loaders must either:

- (1) use a closed system that meets the requirements listed in the Worker Protection Standard (WPS) for dermal protection of agricultural pesticides [40 CFR 170.240(d)(4)],

OR

- (2) Use the probe system described below:

PROBE SYSTEM

Specific requirements for use of the probe closed mixing/loading system:

- ✓ Remove plug from bung of drum containing this product only when drum is sitting on the ground or on a secure level platform, with the bung end of the drum pointed up.
- ✓ Do not pour this product from its drum.
- ✓ Transfer product from the drum to the mixing tank by use of suction hose connected at one end to the suction pump on the mixing tank and connected at the other end to a probe (dip tube) that is inserted through the bung opening into the drum.
- ✓ Do not handle the probe or bung in a manner that allows dripping or splattering of the product onto yourself or any other person.
- ✓ Do not touch the portion of the probe that has been in contact with this product until after the probe has been triple rinsed with water.
- ✓ If all of the product is removed from the drum, then triple rinse the probe while it remains inside the drum.

UN-RINSED PROBES

- ✓ If an un-rinsed probe must be removed from the drum, then use an anti-drip flange, and immediately transfer the probe into a container of rinse water. The anti-drip flange must be designed to remove excess propanil product from the probe as it is extracted from the drum.
- ✓ Take the following steps if the probe must be disconnected from the suction hose before both the probe and the hose have been triple rinsed:
 - (1) Equip the probe end of the hose with a shut off valve,
 - (2) Install a dry break coupling between the valve and the probe,
 - (3) Close the shut-off valve before disconnecting the probe.

PPE FOR ALL TRANSFER SYSTEMS

In addition, mixers and loaders using all systems must:

- wear the personal protective equipment required in the PPE section of this labeling for mixers and loaders,
- wear protective eyewear, if the system operates under pressure, and
- when using a system that meets the requirements in the WPS as a closed system or using a probe system when the probe is not removed, chemical-resistant footwear must be provided, be immediately available, and be used in an emergency, such as a broken package, spill, or equipment breakdown.

All systems must be capable of removing the pesticide from the shipping container and transferring it into mixing tanks and/or application equipment. At any disconnect point, the system must be equipped with a dry disconnect or dry couple shut-off device that is warranted by the manufacturer to minimize drippage.

Flaggers: Human flagging is prohibited. Flagging to support aerial application is limited to use of the Global Positioning System (GPS) or mechanical flaggers.

Enclosed Cabs for Aerial Applicators: Pilots must use an enclosed cockpit that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(6)].

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high watermark. Do not contaminate water when disposing of equipment

washwaters or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas;

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical prior to flooding may result in shallow groundwater contamination due to cracks in the subsoil of the rice paddy.

This product may contaminate water through runoff following rainfall events and by seepage through levees. This product has a high potential for runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Levees should be constructed with adequate time prior to chemical application so that they are compacted to reduce seepage and to hold a 3-6 inch flood.

PHYSICAL AND CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such a plants, soil or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof materials
- Chemical-resistant footwear plus socks
- Protective Eyewear

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Open dumping is prohibited. Do not store this product near fertilizers, seeds, insecticides, or fungicides. Store at temperature above 32°F. If product is allowed to freeze, warm to 50°F and agitate before using. Containers should not be stacked more than three (3) containers high. Reclose all partially used containers by thoroughly tightening screw cap. Damaged or leaking containers that contain product that cannot be used immediately should be transferred to suitable sound containers and properly marked. Any spilled material should be thoroughly absorbed with a suitable absorbent, swept up and transferred to a new or waste container for disposal as indicated under "Pesticide Disposal".

For safety and prevention of unauthorized use, all pesticides should be stored in locked facilities. To prevent accidental misuse, different pesticides should be stored in separate areas with enough distance between to provide clear identification.

Opened, partially used pesticides should be stored in original containers when possible. When transfer to another container is necessary because of leakage or damage, carefully mark and identify contents of the new container. Keep containers closed when not in use.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL:

Nonrefillable Container: Do not reuse or refill this container. Offer for recycling if available. Clean container promptly after emptying.

Nonrefillable container equal to or less than 5 gallons: Triple rinse as follows. Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Nonrefillable container 5 gallons to bulk: Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

REFILLABLE BULK CONTAINER: Refillable container. When the container is empty, replace the cap and seal all openings that have been opened during use; and return the container to the point of purchase, or to a designated location (specified by RiceCo LLC). This container must only be refilled with this pesticide product. **Do Not Reuse the Container for Any Other Purpose.** Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn-out threads and closure devices. Check for leaks after refilling and before transporting. Do not transport if this container is damaged or leaking. If the container is damaged, leaking or obsolete, contact RiceCo LLC at 1-888-835-1313. If not returned to the point of purchase or to a designated location dispose of the container. Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. Disposal of this container must be in compliance with state and local regulations.

STEPS TO BE TAKEN IN CASE MATERIAL IS RELEASED OR SPILLED: Eliminate ignition sources. Ventilate area. Avoid breathing vapors. Use MSHA/NIOSH self-contained breathing apparatus or air mask for large spills in confined areas. Dike the spill with inert material (sand, earth, fuller's earth, etc.) And, if appropriate, transfer the liquid and solid diking material to separate containers for recovery or disposal. Remove contaminated clothing promptly and wash affected skin areas with soap and water. Wash clothing before reuse. Keep out of all sewers and open holes bodies of water. REFER TO PRECAUTIONARY STATEMENTS.

PRECAUTIONS AND RESTRICTIONS

Do not apply to any crop other than rice. **PROPANIL EC** herbicide injures most crops except cereal grains and perennial grasses. Avoid drift or accidental application from turning aircraft on cotton, soybeans, corn, safflower, seedling legumes, vegetables, orchards, vineyards, gardens, shrubs and ornamentals. Once applied, it does not release fumes hazardous to nearby crops.

Do not apply this product through any type of irrigation system.

Do not contaminate water intended for irrigation or domestic purposes.

Do not apply when weather conditions favor drift from target area to be treated.

Water drained from treated rice fields must not be used to irrigate other crops or released within 1/2 miles upstream of a potable water intake in flowing water (e.g., river, stream, etc.) or within 1/2 miles of a potable water intake in a standing body of water, such as a lake, pond or reservoir.

Applications to fields where catfish farming is practiced and draining water from treated fields into areas where catfish farming is practiced is prohibited during 12 months following treatment.

Do Not fish or commercially grow fish, shellfish or crustaceans on treated fields during the 12 months following treatment.

Do Not apply when temperature exceeds 90°F.

Do not plant or transplant crops in the treated area for at least 60 days follow an application of this product.

Do not apply more than 6 quarts **PROPANIL EC** (6.0 lbs active ingredient) per acre per application. Do not apply more than 8 quarts **PROPANIL EC** (8.0 lbs. active ingredient) per acre per season.

Do not apply within 60 days of harvest.

PRODUCT INFORMATION

(FOR RICE GROWN IN THE SOUTHERN UNITED STATES ONLY)

PROPANIL EC is a selective postemergence herbicide for use in rice only for control of the following weeds:

- *BARNYARDGRASS (WATERGRASS)
- BEAKRUSH (SPEARHEAD)
- COCKSPUR, GULF
- CRABGRASS SPECIES
- CROTON, WOOLLY
- DOCK, CURLY
- FOXTAIL SPECIES
- GOOSEGRASS
- HOORAH GRASS
- MEXICANWEED
- PANICUM, TEXAS
- PARAGRASS
- PIGWEED, REDROOT
- REDWEED
- SESBANIA, HEMP (COFFEEBEAN)
- SIGNALGRASS, BROADLEAF
- SPIKERUSH (WIREGRASS)

- Echinochloa crus-gali, E. Colonom*
- Rhynchospora corniculata*
- Echinochloa cruz-pavonis*
- Digitaria spp.*
- Croton capitus*
- Rumex crispus*
- Setaria spp.*
- Eleusine indica*
- Fimbristylis miliaceae*
- Cyperonia castanaefolia*
- Panicum texanum*
- Panicum purpurascens*
- Amaranthus retroflexus*
- Melochia corchorifolia*
- Sesbania exaltata*
- Bracharia platyphytia*
- Eleocharis spp.*

*Biotypes of barnyardgrass may develop that cannot be effectively controlled by propanil alone. Where these biotypes are known or suspected to be present, and are found in a mixed weed population in which **PROPANIL EC** is effective, a tank mixture of **PROPANIL EC** at 4 quarts (4 pounds active) per acre with either Prowl® at 1.5 to 2 pints/A or Bolero® 8EC at 3 to 4 pints/A or Facet® at labeled rates is recommended to

control barnyardgrass (up to 3 leaf stage). These tank mixtures may reduce crop tolerance and are applied at the user's risk.

Read and observe all label directions before using. When tank mixing, always read all individual manufacturers labels. In interpreting all labels for the tank mixture, the most restrictive situations must apply.

PROPANIL EC is an emulsifiable concentrate containing 4 pounds active ingredient per U.S. gallon. **PROPANIL EC** is not a hormone-type herbicide, but kills susceptible weeds by direct contact action. For this reason, thorough coverage of emerged weeds is essential for best results. Only weeds that have emerged and are exposed at time of application will be controlled. **PROPANIL EC** has no pre-emergence or residual herbicidal activity. Apply **PROPANIL EC** herbicide only to fields that have been drained of floodwater. **PROPANIL EC** is most effective if applied when susceptible grasses and broadleaf weeds are small and growing actively under favorable soil moisture and weather conditions. Early weed control removes competition, saves moisture and generally contributes to increased yields.

APPLICATION TIMING AND DOSAGE

Treat grassy and weedy fields when a satisfactory stand of rice that will tolerate flooding is established. The amount of **PROPANIL EC** herbicide to apply depends primarily upon the stage and growth condition of the grasses. The growth stage of the rice is also a factor in dosage and timing limitations, so as to avoid the possibility of excessive residues. For best results apply **PROPANIL EC** herbicide at the rate of 3 to 4 quarts (3 to 4 pounds active) per acre when the grasses are actively growing in the 1 to early 4-leaf stage. This rate will also control many seedling broadleaf and aquatic weeds. Generally this will be 15 to 25 days after planting of the rice. In order to insure satisfactory weed control, do not apply less than 3 quarts of **PROPANIL EC** herbicide per acre in a single spray application.

Apply **PROPANIL EC** herbicide at the rate of 4 to 6 quarts (4 to 6 lbs. active) per acre to actively growing grasses in the 4 to 6 leaf and early tillering stage or when they are in the 2 to 4 leaf stage but stressed under dry soil conditions. Generally this will be 20 to 30 days after planting of the rice.

EMERGENCY TREATMENT: Apply **PROPANIL EC** herbicide at the rate of 5 to 6 quarts (5 to 6 lbs. active) in 15 gallons of spray per acre for emergency control of older tillering grass. Generally this will be 30 to 40 days after planting. If the field is already flooded, the water should be lowered or drained before spraying to expose more of the grass and weeds. Emergency treatment should be considered as a salvage operation only and cannot be relied upon for total control of grass and weeds.

TO AVOID EXCESSIVE RESIDUES AT HARVEST, DO NOT APPLY AFTER THE END OF TILLERING FOR THE RICE VARIETY BEING TREATED. DO NOT APPLY MORE THAN A MAXIMUM OF SIX POUNDS ACTIVE INGREDIENT PER ACRE IN A SINGLE APPLICATION OR EXCEED EIGHT POUNDS ACTIVE INGREDIENT PER ACRE TOTAL DOSAGE PER SEASON.

APPLICATION EQUIPMENT

Aircraft – Fixed wing aircraft or helicopters should have well-designed spray systems that produce a uniform pattern of medium-fine spray droplets. Apply **PROPANIL EC** herbicide on small grass in no less than 10 gallons of total spray per acre with boom-nozzle sprayers. Increase volume to 12 to 15 gallons per acre for larger or denser stands of grass or during periods of low humidity. The optimum effective spray swath width depends on operating conditions and type of aircraft being used. For uniform spray coverage with fixed-wing aircraft, do not exceed a spray swath width of 10 percent greater than the wingspan or the length of the boom in helicopters. Measure the swaths accurately for flagging. (See Spray Drift Management section for additional aerial application requirements.)

GROUND SPRAYERS – Use standard low-pressure herbicide sprayers equipped with boom and flat fan nozzles. Use nozzle sizes that deliver a medium-fine droplet in 15 to 20 gallons total spray per acre at 40 to 50 psi and at ground speeds not in excess of 3 to 4 mph. Avoid raising boom too high. Spray patterns should meet uniformly. Flush all equipment with clear water after each day's use. Clean all equipment, including nurse tanks used for **PROPANIL EC** herbicide, with detergent wash followed by a water rinse, **BEFORE AND AFTER** spraying other pesticides or other crops. (See Spray Drift Management section for additional ground application requirements.)

CROP TOLERANCE AND GROWING CONDITIONS

All leading commercial varieties of rice are exceptionally tolerant to **PROPANIL EC** herbicide. A temporary yellowing or tip burn may be noted after treatment, but new growth is normal. Severe leaf burn and partial killing of rice may occur if the product is applied when rice is under stress and in a weakened growth condition due to disease or insect infestations, excessive soil salts, over watering, or prolonged drought and extremely hot weather. Growers are cautioned not to spray under such conditions and/or when maximum daily temperatures have been or are expected to go above 100°F.

EFFECT OF CLIMATIC CONDITIONS AND CULTURAL PRACTICES ON WEED CONTROL

Field and Seedbed Preparation

Fields should be accurately leveled and contoured and have well-prepared seedbeds free of clods. This encourages uniform and rapid emergence of rice, grass and broadleaf weeds and permits better timing and coverage of **PROPANIL EC** herbicide sprays resulting in optimum weed control.

Water Management

Before application of **PROPANIL EC** herbicide, drained or dry planted fields should be flushed as often as needed to prevent drying and crusting. Flushing encourages uniform emergence and vigorous growth of grass, broadleaf weeds and rice which is essential for best results. Flush fields in sufficient time so that weeds and rice are actively growing at time of treatment. Make sure the field is drained prior to treatment so that

grasses and broadleaf weeds are fully exposed. Weeds that are partially submerged in standing water at time of application will not be satisfactorily controlled. **Preflood Spray Treatment:** After treatment, treated fields should always be flooded before a second infestation of grass has a chance to develop. To prevent more grass from germinating after treatment, fields should be flooded as soon as possible after 24 hours.

Temperature

The temperature a few days before and after applying **PROPANIL EC** has an important bearing on the weed-killing activity. The activity increases as daily maximum temperatures increase above 75°F and decreases as the daily maximum temperatures decline below 75°F. Do not apply **PROPANIL EC** when maximum temperatures have been or are expected to stay below 65°F or to go above 100°F. Low temperatures at time of application are not so important as long as it warms up later during the day.

Relative Humidity and Rain

Grasses and weeds are more responsive to **PROPANIL EC** herbicide during periods of high humidity when the foliage is moist or covered by dew. When the humidity is very low increase spray volume to 12 to 15 gallons per acre for best results. Do not spray when rains threatens within eight hours, to avoid loss of the spray deposit before adsorption by the grass.

Wind

Avoid applications when the wind speed exceeds 10 mph because of drift hazard to sensitive crops and the possibility of uneven (streaked) applications.

COMPATIBILITY WITH OTHER CHEMICALS

Tank-mix applications of **PROPANIL EC** herbicide with other herbicides, insecticides, or liquid fertilizers may reduce crop tolerance and/or weed control or impair mixing properties. Use of these products in tank-mix application with **PROPANIL EC** herbicide is done at the user's risks.

SPRAY DRIFT MANAGEMENT

Avoiding spray drift at the application site is the responsibility of the applicator and the grower. The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions. The following drift management requirements must be followed to avoid off-target drift movement from the aerial applications to agricultural field crops.

Apply only when the wind speed is less than or equal to 10 mph at the application site. Apply as a medium or coarser spray (ASAE standard 572).

Additional requirements for ground applications:

Apply using a nozzle height of no more than 4 feet above the ground or crop canopy.

Importance of Droplet Size

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (see Wind, Temperature and Humidity, and Temperature Inversion sections of this label.)

Controlling Droplet Size

Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy protection. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles – Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation – Orienting nozzles so that the spray is released backwards, parallel to the airstream will produce larger droplets than other orientations. Significant deflection from the horizontal will reduce droplet size and increase drift potential.

Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles. Solid stream nozzles oriented straight back produce larger droplets than other nozzle types.

Boom Length - The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter.

Application Height – Do not release spray at a height greater than 10 feet above the ground or crop canopy. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

Swath Adjustment

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind. Swath adjustment distance should increase with increasing drift potential (higher wind, smaller drops, etc.)

Wind

Drift potential is lowest between wind speeds of 2-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 2 mph due to variable wind direction and high inversion potential. Note: local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect drift.

Temperature and Humidity

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

Temperature Inversions

Do not make any type of application into temperature inversions

Temperature inversions restrict vertical air mixing, which causes small, suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves literally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upwards and rapidly dissipates indicates good vertical air mixing.

Sensitive Areas

Do not apply by air if drift can occur to sensitive nontarget crops or plants that are within 100 feet of the application site. Sensitive areas include, but are not limited to, residential areas, bodies of water, known habitat for threatened or endangered species, and non-target crops.

EMERGENCY RELEASE PROVISION

Water holding (discharge) intervals for flood water following propanil application in all states:

For delayed flood (water-seeded) rice grown south of Interstate Highway 10 from the Texas/Louisiana border to Houston and east of State Highway 35 from Houston to Port Lavaca – Flood water must be held for 10 days after application, unless excessive rainfall completely submerges the rice crop and forces premature release. For Texas rice grown in areas north or west of these boundaries, the water holding interval will be 7 days.

For delayed flood (water-seeded) rice in Southern Louisiana south of Highway 14 – Flood water must be held for 15 days after propanil application unless excessive rainfall completely submerges the rice crop and forces premature release. Delayed flood (water-seeded) rice in Louisiana, north of Highway 14 boundary, is subject to the 7-day water holding interval provisions.

For rice grown in California and all other parts of the US not mentioned above – Flood water must be held for 7 days after application, unless excessive rainfall completely submerges the rice crop and forces premature release.

Insecticides

Severe injury or kill of rice plants may result from tank-mix combinations or separate sprays of **PROPANIL EC** herbicide and certain insecticides. Do not combine **PROPANIL EC** herbicide with carbamates insecticides, such as carbaryl (Sevin, etc.), Methomyl (Lannate, Nudrin, etc.), or organophosphate insecticides such as parathion, methyl parathion, Guthion, malathion, Systox, WPN, Phosphamidron, etc. Do not apply any of the above insecticides to rice fields within 14 days before or after **PROPANIL EC** herbicide. Do not use carbamates or systemic organophosphorus insecticides on rice fields to be treated with **PROPANIL EC** herbicide. Do not apply to rice fields that were planted with rice seed treated with bird repellents containing methiocarb such as Mesurol, Borderland Red, etc. Consult local Extension specialist for current recommendations of approved insecticides on rice.

SPRAY MIXTURE PREPARATION

Wet Spray Application

Thoroughly mix **PROPANIL EC** with clean water (water that is free of sediment and agricultural chemicals) in the spray tank. Do not use water from paddies. Only approved drift control agents may be used with **PROPANIL EC**. Do not use any other additives except as directed by this label.

To ensure uniform mixing and application, agitate the mixture before application. If the mixture is not sprayed immediately after agitation, re-agitate it before application. Always apply **PROPANIL EC** spray preparation within 24 hours of product mixing, or the product may degrade.

Do not store **PROPANIL EC** in nurse tanks or any other tanks used to store or transport clean water. Install one-way valves (anti-siphoning devices) on lines and hoses of mixing/loading equipment to prevent contamination of nurse tanks or other clean water sources.

Mixing and application equipment exposed to **PROPANIL EC** cannot be used for anything other than rice applications until it has been cleaned according to the procedures in the Sprayer Cleanup section of this label.

Additional Mixing Instructions (wet spray)

1. Fill the tank 1/4 to 1/3 full of clean water.
2. While agitating, add the required amount of **PROPANIL EC**.
3. Continue agitation until the product is fully dispersed, at least 5 minutes.
4. Once the **PROPANIL EC** is fully dispersed, maintain agitation and continue filling the tank with water. The product should be thoroughly mixed with water before adding any other material.
5. As the tank is filling, add the required tank mix partner (other labeled rice herbicides, adjuvants, drift control agents, etc.).
6. If the mixture is not continuously agitated, settling may occur. If settling occurs, thoroughly re-agitate before using.

- 7. Apply **PROPANIL EC** spray preparations within 24 hours of product mixing, or the product may degrade.
- 8. If **PROPANIL EC** and a tank mix partner are to be applied in multiple loads, pre-slurry the product in clean water prior to adding to the tank. This will prevent the tank mix partner from interfering with the dissolution of the **PROPANIL EC**.

SPRAYER CLEANUP

Before using equipment exposed to **PROPANIL EC** to treat another crop, clean the sprayer and any other equipment (loading hoses, batch tanks, etc.) using the following procedure:

- 1. Steam-clean tank using a non-chlorine-based detergent, taking care to remove all physical residues.
- 2. Thoroughly rinse sprayer, tanks, boom, and hoses with clean water (free of sediment and agricultural chemicals).
- 3. Fill the tank one-half full with clean water and add Nutrasol at 32 oz. per 100 gallons water. Fill the tank to capacity with clean water. Flush the nozzles, boom, and hoses, and agitate (and recirculate, if possible) the sprayer for 15 minutes. Drain the equipment, taking care to flush the boom and hoses thoroughly.
- 4. Rinse tanks, hoses and nozzles with clean water to remove Nutrasol.
- 5. Fill the tank one-half full with clean water and add 1 gal. 21% ammonia or 7 gallons 3% ammonia per 100 gallons water. Fill the tank to capacity with clean water. Flush the nozzles, boom, and hoses and agitate (and recirculate, if possible) the sprayer for 15 minutes. Drain the equipment, taking care to flush the boom and hoses thoroughly.
- 6. Remove nozzles, screens, and strainers, and clean them separately.
- 7. Rinse tanks, booms, and hoses with clean water.
- 8. Repeat steps 5 and 7 an additional 3 times.
- 9. Rinse tanks, booms, and hoses to remove all traces of ammonia.
- 10. Water rinses may be applied to rice fields. Dispose of bleach rinses at an approved waste disposal facility.

NOTE: When applying multiple loads of **PROPANIL EC** several days in a row, the following procedure must be performed at the end of each day; partially fill the tank with fresh water, flush the boom and hoses, and allow to sit overnight.

ATTENTION: Do not use chlorine bleach with ammonia. All traces of liquid fertilizer containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed from the mixing and application equipment using water before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odor that can cause eye, nose, and throat and lung irritation. Do not clean equipment in an enclosed area.

Perform cleanup procedures on batch tanks and any other mixing equipment separately from aircraft hoppers. Take care to clean loading hoses and any other equipment or surfaces exposed to **PROPANIL EC**.

CONDITIONS OF SALE AND WARRANTY

RiceCo LLC warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label only when used in accordance with label directions under normal conditions of use. To the extent permitted by applicable law, RICECO LLC MAKE NO OTHER EXPRESS OR IMPLIED WARRANTIES EITHER OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR USE. Handling, storage and use of the product by Buyer or User are beyond the control of RiceCo LLC and Seller. Risks such as crop injury, ineffectiveness or other unintended consequences resulting from, but not limited to weather or soil condition, presence of other materials, disease, pests, drift to other crops or property or failure to follow label directions will be assumed by the Buyer or User. TO THE EXTENT PERMITTED BY APPLICABLE LAW, IN NO CASE SHALL RICECO OR SELLER BE HELD LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE HANDLING, STORAGE OR USE OF THIS PRODUCT

Bolero is a registered trademark of Kumiai Chemical Industry LTD
Facet and Prowl are registered trademarks of BASF Company



**5100 POPLAR AVENUE, SUITE 2428
MEMPHIS, TENNESSEE 38137 USA**

“....from the paddy to the plate”

Stamped Accepted 3/24/10